



Lepidoptera of South Ossetia (Northern Transcaucasia). Part I. Introduction and Superfamily Pyraloidea Latreille, 1809

Aleksandr N. Streltzov

Herzen State Pedagogical University of Russia, 48 Moika Emb., Saint Petersburg 191186, Russia

Petr Ya. Ustjuzhanin

Altai State University, 61 Lenina Ave., Barnaul 656049, Russia; Biological Institute, Tomsk State University, 36 Lenina Ave., Tomsk 634050, Russia;
<https://orcid.org/0000-0002-5222-2241>

Roman V. Yakovlev

Altai State University, 61 Lenina Ave., Barnaul 656049, Russia; Biological Institute, Tomsk State University, 36 Lenina Ave., Tomsk 634050, Russia;
<https://orcid.org/0000-0001-9512-8709>

In the first part of the publication, we present the faunal list of Pyraloidea of South Ossetia, including 111 species of 64 genera, belonging to two families. For the first time, 96 species are reported for South Ossetia. *Scoparia manifestella* (Herrich-Schäffer, 1848) and *Scoparia perplexella* (Zeller, 1839) are found in the Caucasus for the first time, and the new species *Scoparia sinevi* Streltzov, 2022 is currently indicated as an endemic of South Ossetia.

Acta Biologica Sibirica 8: 281–296 (2022)

doi: 10.14258/abs.v8.e18

<http://journal.asu.ru>

Corresponding author: Aleksandr N. Streltzov (streltzov@mail.ru)

Academic editor: A. Matsyura

Received 22 July 2022 | Accepted 30 August 2022 | Published 9 September 2022

<http://zoobank.org/8C8BBF94-29DA-4F4B-98A6-B9F1E21DC616>

Citation: Streltzov AN, Ustjuzhanin PYa, Yakovlev RV (2022) Lepidoptera of South Ossetia (Northern Transcaucasia). Part I. Introduction and Superfamily Pyraloidea Latreille, 1809. Acta Biologica Sibirica 8: 281–296. <https://doi.org/10.14258/abs.v8.e18>

Keywords

Biodiversity, Caucasus, species richness, fauna, Pyralidae, Crambidae

Introduction

South Ossetia (Fig. 1) is a compact region in Northern Transcaucasia, located at the southern slope of the Central Caucasus at the foothills of the Inner Kartli Plain. Almost 90 % of the territory of the republic is located at the altitudes higher than 1000 m above sea level. The highest point of South

Ossetia is mountain Khalatsa (3938 m). The mountainous portion of the republic is presented by the Dvaletskiy ridge on the border with Russia and the transverse ridges – Rachinsky, Likhsky, Kesheltsky, Mashkharsky, Dzausky, Gudissky, Kharulsky, Lomissky and Mtuleti. The foothill area is located in the extreme south of the region, where the valleys of the rivers Prone, Greater and Lesser Liakhva, Mejuda and Ksan go to the northern outskirts of the Inner Kartli Plain.



Figure 1. South Ossetia on the map of Caucasus

The climate of South Ossetia changes with the altitude from the moderately dry and warm, with a cool winter and hot summer (in the south-eastern part) to the humid, with a cold long winter and a cool summer at 2000 m, and to the humid and cold during all the year at 3000 m (in the north). The average temperature of January is -1°C in Tskhinvali (867 m), at 2000 m – from -3 to -8°C, higher – from -12 to -14°C, and of July – 20,6°C, 8-13°C, 4-2°C respectively. The annual amount of precipitation goes up from 600 mm on the plain to 1000-1800 mm in the highlands. The snow border lies at 3200-3300 m. The current glaciation is minor, the largest glacier, Lagzzigi, has an area about 1,8 km².

At the foothills, the secondary shrub steppes are common, with *Rosa* L., *Crataegus* Tourn. Ex L. (Rosaceae), and *Paliurus spina-christi* Mill. (Rhamnaceae) on the site of cut forests. In the extreme south, there are portions of Phrygana. In the valleys, there are trees and shrubs on the alluvial soils. Trees and shrubs take about half of the territory. In the lower belt of the forest (600-700 m a.s.l.) the predominating are *Quercus* and *Quercus-Caprinus* forests with shrubs (*Mesmilus* L. (Rosaceae), *Cornus* L. (Cornaceae), and *Berberis* L. (Berberidaceae)) and fruit trees (*Malussylvestris* (L.) Mill., *Pyrus communis* L., and *Prunus avium* (L.) L. (Rosaceae)). At the altitude

of 1000–1100 m, they are replaced by *Fagus* L. forests with *Caprinus* L. (Betulaceae) and *Acer* L. (Sapindaceae). Higher than 1500–1600 m, there are mixed *Fagus-Picea* forests, interspersed with *Picea* Forests, in some places with *Abies* Mill. and *Pinus* L. (Pinaceae). In the west forests, some Colchisian elements are found (in the evergreen under-bush – *Prunus laurocerasus* L. (Rosaceae), *Ilex* L. (Aquifoliaceae), *Ruscus* Tour. & L.); the epiphytes are numerous. In gorges, *Taxus baccata* L. (Taxaceae) is preserved in some places. At the upper border of the forests, the crooked woods of *Fagus* L. (Fagaceae), *Acer trautvetteri* Medw. (Sapindaceae), and *Betula raddeana* Trautv. (Betulaceae) are often found; beech accounts for about 80% of the total forested area. Higher than 2200–2350 m there are sub-alpine (with *Rhododendron caucasicum* Pall. (Ericaceae) and alpine meadows (Dzagoev 2003).

The faunal diversity of South Ossetia is very poorly studied. Only two articles on the spider and amphibian fauna completely illustrate the richness of the animal world of South Ossetia (Ponomarev and Komarov 2015; Tuniev et al. 2017). The information on the rare and endangered species of flora and fauna of South Ossetia is provided in the beautifully published Red Book (Bazaev et al. 2017). The articles on the Lepidoptera fauna are rather fragmentary (Pospelov et al. 1986; Pukhaev et al. 1987; Pukhaev and Pukhaeva 1989, 1991; Pukhaeva 1995; Didmanidze 2004; Komarov 2013; Dobronosov and Komarov 2014, 2015).

This series of articles is devoted to the Lepidoptera fauna of South Ossetia. The superfamily Pyraloidea Latreille, 1809 of South Ossetia are practically not studied. In the single article on this theme, they give data on the distribution of 15 Pyraloidea species (Dobronosov and Komarov 2014), collected on the territory and environs of Tskhinval and Atriskhev cities. The authors report the following for this region: *Agriphila straminea* (Denis & Schiffermüller, 1775), *Chilo phragmitella* (Hübner, 1805), *Chrysocrambus craterellus* (Scopoli, 1763), *Evergestis frumentalis* (Linnaeus, 1761), *Cynaedadentalis* (Denis & Schiffermüller, 1775), *Loxostege sticticalis* (Linnaeus, 1761), *Scoparia subfusca* Haworth, 1811, *Diasemia reticularis* (Linnaeus, 1761), *Nomophila noctuella* (Denis & Schiffermüller, 1775), *Paratalanta pandalis* (Hübner, 1825), *Aphomia sociella* (Linnaeus, 1758), *Endotricha flammealis* (Denis & Schiffermüller, 1775), *Pyralis farinalis* (Linnaeus, 1758), and *Selagia argyrella* (Denis & Schiffermüller, 1775).

Additionally, Dobronosov (2017) and Tuniev et al. (2018) reported the important invasive species *Cydalima perspectalis* (Walker, 1859) for the village of Nagutni of Znaur district, and for the balneary "Nyfs" of South Ossetia. This species is widely distributed in the Eastern Palaearctic (Streltzov 2008) and also found in Russian Far East (Kirpichnikova 2005). In XXI, it penetrated to Europe (Kruger 2008). From Italy it was introduced to the Caucasus and then spread all over the Caucasian region (Shchurov et al. 2015; Dobronosov 2017; Tuniev et al. 2018; Ustjuzhanin et al. 2022).

The new species of Crambidae, *Scoparia sinevi* Streltzov, 2022 was described by us (Streltzov et al. 2022) from Rachinsky ridge in the north-west of South Ossetia.

Material and methods

The specimens were collected in South Ossetia in seven localities. The list of collection sites was chosen by Alexandr Fomichev, Barnaul, for the most complete overview of the altitudinal belts and plant communities of South Ossetia.

The collections were carried out by manual collection during the daytime and at dusk, as well as on light screens Naturaliste-150 and Naturaliste-180 (using lamps OSRAM-160, 250 W), powered by the inverter generator Honda EU10i (Fig. 2) and autonomous light traps ENTOSPHINX lamp UV LED 12 V/19,2W (equipped with diodes 240 UV LED) (Fig. 3). Deadening of the specimens was carried out using ethyl acetate. The material was mounted on entomological pins.



Figure 2. Light screens Naturaliste-180 (photo by R. Yakovlev)



Figure 3. Autonomous light traps (photo by A. Fomichev)

The examined material is kept in the collections:

ASSP collection of Alexandr Streltzov (Saint-Petersburg, Russia);

CUK collection of Petr Ustjuzhanin and Vasiliy Kovtunovich (Novosibirsk, Moscow, Russia);

PMM collection of Pavel Morozov (Moscow, Russia);

RMBH Russian Museum of Biodiversity Hotspots, N. Laverov Federal Center for Integrated Arctic Research of the Ural Branch of the Russian Academy of Sciences (Arkhangelsk, Russia);

RYB collection of Roman Yakovlev (Barnaul, Russia);

ZISP Zoological Institute of the Russian Academy of Sciences (Saint-Petersburg, Russia).

List of collecting localities (Fig. 4)

1. South Ossetia, Tskhinval Distr., 2 km NW Grom, 42°10'6"N / 44°11'53"E, 930 m, 22–25.06.2021, A. Streltzov, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 5);

2. South Ossetia, Leningor Distr., 4 km E Leningor, 42°08'45"N, 44°30'55"E / 1200 m, 26–27.06.2021, A. Streltzov, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 6);
3. South Ossetia, Dzaus Distr., 4 km NNE Kvaisa, Koz lake, 42°33'32"N / 43°37'59"E, 1580 m, 28–30.06.2021, A. Streltzov, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 7);
4. South Ossetia, Dzaus Distr., Rachinsky Range, near Dodtota, 42°27'25"N / 43°43'18"E, 1750 m, 1–2.07.2021, A. Streltzov, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 8);
5. South Ossetia, Dzaus Distr., Dvalet Range, near Kherusel't, 42°32'37"N / 43°47'32"E, 1760 m, 3–5.07.2021, A. Streltzov, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 9);
6. South Ossetia, Dzaus Distr., Mtiolet Range, near Erman, 42°31'2"N / 44°14'10"E, 2140 m, 7–9.07.2021, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 10);
7. South Ossetia, Znaur Distr., 2 km W Dzagina, 42°14'34"N / 43°43'11"E, 1100 m, 11–12.07.2021, P. Ustjuzhanin & R. Yakovlev leg. (Fig. 11).




Figure 4. Map of South Ossetia with collecting localities**Figure 5.** Tskhinval Distr., 2 km NW Grom, 42°10'6"N / 44°11'53"E (photo by R. Yakovlev)



Figure 6. Leningor Distr., 4 km E Leningor, $42^{\circ}08'45''N$, $44^{\circ}30'55''E$ (photo by R. Yakovlev)



Figure 7. Dzaus Distr., 4 km NNE Kvaisa, Koz lake, $42^{\circ}33'32''N / 43^{\circ}37'59''E$ (photo by R. Yakovlev)



Figure 8. Dzaus Distr., Rachinsky Range, near Dodtota, $42^{\circ}27'25''N / 43^{\circ}43'18''E$ (photo by R. Yakovlev)



Figure 9. Dzaus Distr., Dvalet Range, near Kherusel't, $42^{\circ}32'37''N / 43^{\circ}47'32''E$ (photo by R. Yakovlev)



Figure 10. Dzaus Distr., Mtiulet Range, near Erman, 42°31'2"N / 44°14'10"E (photo by R. Yakovlev)



Figure 11. Znaur Distr., 2 km W Dzagina, 42°14'34"N / 43°43'11"E (photo by R. Yakovlev)

Results

#	Taxa	Localities						
		1	2	3	4	5	6	7
PYRALIDAE								
1	<i>Aphomia sociella</i> (Linnaeus, 1758)	-	-	-	+	-	-	-
2	<i>Endotricha flammealis</i> ([Denis & Schiffmüller], 1775)	+	+	-	-	-	-	+
3	<i>Synaphe antennalis</i> (Fabricius, 1794)	+	-	-	-	-	-	-
4	<i>Hypsopygia glauccinali</i>	+	+	-	-	-	-	-

	<i>s</i> (Linnaeus, 1758)								
5	<i>Pyralis farinalis</i> (Linnaeus, 1758)	-	+	-	-	-	-	-	-
6	<i>Pyralis</i> (Christoph, 1893)	-	-	-	-	-	-	-	+
7	<i>Elegia similella</i> (Zincken, 1818)	+	+	-	-	-	-	-	+
8	<i>Pempeliella ornatella</i> ([Denis & Schiffermüller], 1775)	-	+	+	+	-	-	-	-
9	<i>Delplanqueia inscripella</i> (Duponchel, 1836)	-	+	-	-	+	-	-	-
10	<i>Catastia marginata</i> ([Denis & Schiffermüller], 1775)	-	-	+	-	-	-	-	-
11	<i>Sciota rheinella</i> (Zincken, 1818)	-	-	-	-	-	-	-	+
12	<i>Psorosanucleolella</i> (Möschler, 1866)	+	+	-	-	-	-	-	-
13	<i>Selagia argyrella</i> ([Denis & Schiffermüller], 1775)	-	+	-	-	-	-	-	-
14	<i>Selagia spadicella</i> (Hübner, 1796)	-	+	-	-	-	-	-	+
15	<i>Etiella zincicella</i> (Treitschke, 1832)	-	+	-	-	-	-	-	-
16		+	-	-	-	-	-	-	-

	<i>Merulempista cingillella</i> (Zeller, 1846)							
17	<i>Oncocera</i> (Scopoli, 1763)	+	+	+	+	-	-	+
18	<i>Laodamia faecella</i> (Zeller, 1839)	-	-	-	-	+	-	-
19	<i>Pempelia albariella</i> (Zeller, 1839)	-	+	-	-	-	-	-
20	<i>Pempelia palumbella</i> ([Denis & Schiffermüller], 1775)	-	+	-	-	-	-	-
21	<i>Moitrelia obductella</i> (Zeller, 1839)	-	-	-	-	-	-	+
22	<i>Rhodophaea formosa</i> (Haworth, 1811)	+	-	-	-	-	-	-
23	<i>Dioryctria abietella</i> ([Denis & Schiffermüller], 1775)	-	-	+	+	-	-	+
24	<i>Phycita roborella</i> ([Denis & Schiffermüller], 1775)	-	+	-	-	-	-	+
25	<i>Hypochalcia ahenella</i> ([Denis & Schiffermüller], 1775)	-	-	-	+	+	-	-
26	<i>Hypochalcia subrubiginella</i> (Ragonot, 1887)	-	+	-	-	+	-	-
27	<i>Nephopterix angustella</i> (Hübner, 1796)	+	+	-	-	-	-	-
28	<i>Acrobasis advenella</i> (Zincken, 1818)	-	+	-	-	-	-	+
29	<i>Acrobasis</i>	-	+	-	-	-	-	-

	<i>c on soci ella</i> (Hübner, 1813)							
30	<i>Acrobasis obtusella</i> (Hübner, 1796)	-	+	-	-	-	-	+
31	<i>Acrobasis sua vella</i> (Zincken, 1818)	-	+	-	-	-	-	+
32	<i>Acrobasis tumidana</i> ([Denis & Schiffermüller], 1775)	-	+	-	-	-	-	+
33	<i>Eccopisa effractella</i> (Zeller, 1848)	+	-	-	-	-	-	-
34	<i>Episcythrasti s tabidella</i> (Mann, 1864)	+	-	-	-	-	-	-
35	<i>Eurhodope rosella</i> (Scopoli, 1763)	+	+	-	-	-	-	+
36	<i>Euzophera bigella</i> (Zeller, 1848)	+	+	-	-	-	-	-
37	<i>Euzophera</i> (Heinemann, 1865)	+	+	+	-	-	-	-
38	<i>Nyctegretis linea na</i> (Scopoli, 1786)	+	-	-	-	-	-	-
39	<i>Homoeosoma nebulosa</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	-
40	<i>Homoeosoma sinu ella</i> (Fabricius, 1794)	+	+	-	-	-	-	+
41	<i>Phycitodes in quinate</i>	-	-	-	-	-	-	+

	<i>lla</i> (Ragonot, 1887)							
42	<i>Phycitodes nigrilimbella</i> (Ragonot, 1887)	-	-	-	+	-	-	+
43	<i>Phycitodes saxicola</i> (Vaughan, 1870)	-	+	-	-	-	-	+
44	<i>Ephestia elutella</i> (Hübner, 1796)	+	+	-	-	-	-	+
45	<i>Cadra furcatella</i> (Herrich-Schäffer, 1849)	+	+	-	-	-	-	+
CRAMBIDAE								
46	<i>Euchromius bella</i> (Hübner, 1796)	-	-	-	-	-	-	+
47	<i>Euchromius ocellea</i> (Haworth, 1811)	-	+	-	-	-	-	-
48	<i>Euchromius superbellus</i> (Zeller, 1849)	+	-	-	-	-	-	-
49	<i>Chrysoteuchia culmella</i> (Linnaeus, 1758)	+	+	+	+	+	-	+
50	<i>Crambus lathonie llus</i> (Zincken, 1817)	-	-	-	+	-	-	-
51	<i>Crambus pascuellus</i> (Linnaeus, 1758)	+	+	+	+	+	-	+
52	<i>Crambus perrella</i> (Scopoli, 1763)	-	-	-	+	-	-	-
53	<i>Agriphila stramineella</i> ([Denis & Schifferm	-	-	-	-	+	-	-

	üller], 1775)								
54	<i>Catoptria</i> (Lederer, 1870)	+	+	-	-	-	-	-	-
55	<i>Catoptria confusella</i> (Staudinger, 1881)	-	-	-	+	+	-	-	-
56	<i>Catoptria laevigatellus</i> (Lederer, 1870)	-	+	-	+	+	+	-	-
57	<i>Catoptria pinella</i> (Linnaeus, 1758)	-	+	-	-	-	-	-	+
58	<i>Catoptria verellus</i> (Zincken, 1817)	-	-	+	-	-	-	-	-
59	<i>Metacrambus carectellus</i> (Zeller, 1847)	-	-	-	-	-	-	-	+
60	<i>Chrysocrambus craterellus</i> (Scopoli, 1763)	+	-	-	-	-	-	-	-
61	<i>Chrysocrambus linetellus</i> (Fabricius, 1781)	+	+	-	-	-	-	-	+
62	<i>Thisanotia chrysonuchella</i> (Scopoli, 1763)	+	-	-	-	-	-	-	-
63	<i>Pediasia contaminella</i> (Hübner, 1796)	-	+	-	+	-	-	-	-
64	<i>Platytes cerussella</i> ([Denis & Schiffermüller], 1775)	+	+	-	-	-	-	-	-
65	<i>Scoparia</i> (Herrich-Schäffer, 1848)	+	-	-	-	-	-	-	-
66	<i>Scoparia perplexella</i>	+	-	-	-	-	-	-	-

	(Zeller, 1839)							
67	<i>Scoparia in grat ella</i> (Zeller, 1846)	+	+	-	+	+	-	-
68	<i>Scoparia sinevi</i> (Streltzov, 2022)		-	-	+	-	-	-
69	<i>Eudonia lacustrata</i> (Panzer, 1804)	+	+	-	-	-	-	-
70	<i>Eudonia mur ana</i> (Curtis, 1827)	+	+	-	-	-	-	-
71	<i>Eudonia</i> (Stainton, 1849)	+	+	-	-	-	-	-
72	<i>Cynaeda dent alis</i> ([Denis & Schifferm üller], 1775)	+	+	-	-	-	-	+
73	<i>Ephelis cruentalis</i> (Geyer, 1832)	-	-	-	-	-	-	+
74	<i>Evergestis forf icalis</i> (Linnaeus, 1758)	-	-	+	-	-	-	-
75	<i>Hellula unda lis</i> (Fabricius, 1781)	+	-	-	-	-	-	-
76	<i>Loxostege sticticalis</i> (Linnaeus, 1761)	+	+	-	-	-	-	-
77	<i>Paratalanta cultralis</i> (Staudinger, 1867)	-	-	-	-	-	-	+
78	<i>Paratalanta hyalinalis</i> (Hübner, 1796)	+	+	-	+	+	-	+
79	<i>Pyrausta aurata</i> (Scopoli, 1763)	-	+	-	-	-	-	+
80	<i>Pyrausta cingulata</i> (Linnaeus, 1758)	+	+	-	-	-	-	+

81	<i>Pyrausta despicata</i> (Scopoli, 1763)	+	+	+	-	+	-	+
82	<i>Pyrausta ost rinalis</i> (Hübner, 1796)	+	+	-	-	+	-	+
83	<i>Pyrausta rectefascialis</i> (Toll, 1936)	+	+	-	-	-	-	+
84	<i>Pyrausta aere alis</i> (Hübner, 1793)	+	+	-	+	+	-	-
85	<i>Sitochroa palealis</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	-
86	<i>Sitochroa verticalis</i> (Linnaeus, 1758)	+	-	-	+	-	-	+
87	<i>Psammotis pulveralis</i> (Hübner, 1796)	+	+	-	-	-	-	-
88	<i>Anania coronata</i> (Hufnagel, 1767)	-	+	+	+	-	-	-
89	<i>Anania crocealis</i> (Hübner, 1796)	+	+	+	+	+	-	-
90	<i>Anania funebris</i> (Ström, 1768)	-	+	+	-	-	-	-
91	<i>Anania lancealis</i> ([Denis & Schiffermüller], 1775)	-	+	+	-	-	-	-
92	<i>Anania sta chydalis</i> (Germar, 1821)	-	+	+	+	-	-	-
93	<i>Anania ver basalis</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	+

94	<i>Anania hortulata</i> (Linnaeus, 1758)	+	+	+	-	-	-	-
95	<i>Anania fuscalis</i> ([Denis & Schiffermüller], 1775)	-	+	+	+	+	-	-
96	<i>Patania ruralis</i> (Scopoli, 1763)	-	-	-	-	-	-	+
97	<i>Mecyna flavalis</i> ([Denis & Schiffermüller], 1775)	+	+	-	-	-	-	-
98	<i>Mecyna trinalis</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	-
99	<i>Udea</i> (Herrich-Schäffer, 1851)	-	-	-	+	+	-	-
100	<i>Udea lutealis</i> (Hübner, 1809)	-	-	-	-	+	-	-
101	<i>Udea olivalis</i> ([Denis & Schiffermüller], 1775)	-	-	+	+	+	-	-
102	<i>Udea vastalis</i> (Christoph, 1887)	+	-	-	-	-	-	-
103	<i>Nomophila noctuella</i> ([Denis & Schiffermüller], 1775)	+	+	+	-	-	-	-
104	<i>Agroteraneomorialis</i> (Scopoli, 1763)	+	-	-	-	-	-	-
105	<i>Cydalima</i> (Walker, 1859)	+	-	-	-	-	-	-
106	<i>Dolicharthria punctalis</i> ([Denis & Schiffermüller]	+	-	-	-	-	-	-

[, 1775)

Table 1. Distribution of Pyraloidea of South Ossetia

Discussion

Most of the found species are new for the fauna of South Ossetia. Only five of the species known before (Dobronosov and Komarov 2014) are absent in our collection - *Chilo phragmitella* (Hübner, 1805), *Evergestis frumentalis* (Linnaeus, 1761), *Scoparia subfusca* Haworth, 1811, *Diasemia reticularis* (Linnaeus, 1761), *Paratalanta pandalis* (Hübner, 1825). Thus, the known fauna of Pyraloidea of South Ossetia includes 111 species of 2 families and 64 genera. In the species composition, Pyraloidea of South Ossetia are very close to the fauna of the Northern Caucasus (Sinev and Streltzov 2019; Sinev et al. 2019; Ustyuzhanin et al. 2022; Yakovlev et al. 2022), however, there is a Transcaucasian specificity, for example *Phycitodes inquinatella* (Ragonot, 1887), *Phycitodes nigrilimbella* (Ragonot, 1887), *Catoptria confusella* (Staudinger, 1881) and *Udea vastalis* (Christoph, 1887) are not found in Russia and accordingly, in the Northern Caucasus. Such species as *Scoparia manifestella* (Herrich-Schäffer, 1848) and *Scoparia perplexella* (Zeller, 1839) were found in the Caucasus for the first time, and the new species *Scoparia sinevi* (Streltzov, 2022) can currently be considered an endemic of South Ossetia.

Acknowledgments

The authors express their gratitude to R. Bakhanov (Gorno-Altaisk), A. Dzhussoev, F. Dzagoev, V. Tedeev, and V. Gabaev (all Tskhinvali) for organizing the field studies in the South Ossetia in June–July of 2021. For finding rare publications, we are grateful to Yu. Komarov and R. Pukhaev (Vladikavkaz). In conclusion, we would like to thank all the people of South Ossetia, whom we met during the expedition, for their help and politeness towards us and our work. Research by Roman Yakovlev and Petr Ustyuzhanin was supported by a grant from Altai State University No.15/22-VG(11 April 2022) "Lepidoptera of South Ossetia".

References

- Bazaev FB, Bestaev AZ, Busarova NV, Butaeva FG, Vaniev AG, Gabaev VN, Dzhioeva TsG, Doroshina GYa, Kabulov AZ, Kabulov ZE, Kabulov ID, Kachmazov DG, Kokoev TI, Komarov YuE, Lavrinenko YuV, Lotiev KYu, Nikolaev IA, Pukhaev RV, Pukhaeva ZA, Timuthin IN, Tuniev BS, Tuniev SB, Khetagurov KhA, Tskhovrebova SS, Tskhovrebova NI, Chibirova AKh, Shikov EV, Yusupov ZM (2017) Red data book of South Ossetia. Poligraphservis & T., Nalchik, 304 pp. [In Russian]
- Didmanidze EA (2004) Annotated list of Diurnal Butterflies (Lepidoptera: Rhopalocera) of Georgia and adjacent territory from Southern Caucasus. Proceedings of the Institute of Zoology Georgia 22: 197–226.
- Dobronosov VV (2017) New data on a box pyralid of *Cydalima perspectalis* (Walker, 1859) in the Central Caucasus. Aekonomika: ekonomika i selskoe hozajstvo 10 (22): 1–14. [In Russian]
- Dobronosov VV, Komarov YE (2014) To the knowledge of the fauna of Grass-Moths Microlepidoptera (Microlepidoptera, Pyraloidea) of the Republic of South Ossetia. Actual problems of ecology and biodiversity conservation in Russia and neighboring countries 10: 86–89. [In Russian]
- Dobronosov VV, Komarov YE (2015) To the knowledge of the fauna of Macrolepidoptera (Lepidoptera, Heterocera) of the Republic of South Ossetia. Bulletin of the Krasnodar Regional

Branch of the Russian Geographical Society 8: 179–186. [In Russian]

Dzagoev AD (2003) Geography of South Ossetia. Olimp, Vladikavkaz, 224 pp. [In Russian]

Kirpichnikova VA (2005) Addition: 49. Fam. Pyralidae. Key to insects of the Russian Far East. Vol. V. Caddisflies and Lepidoptera. P. 5. Dalnauka, Vladivostok, 526–539. [In Russian]

Komarov YE (2013) To the fauna of nocturnal Lepidoptera of the Republic of South Ossetia. Biodiversity and rational use of natural resources (Materials of the All-Russian Conference with International Participation, Makhachkala, 27–28 March 2013): 107–109. [In Russian]

Kruger EO (2008) *Glyphodes perspectalis* (Walker, 1859) – neu furdie Fauna Europas (Lepidoptera, Crambidae). Deutsche Entomologische Zeitschrift 118 (2): 81–83.

Ponomarev AV, Komarov YE (2015) The Spiders (Aranei) of Republic of Southern Ossetia. The South of Russia: ecology, development 10 (1): 116–147. [In Russian]

Pospelov SM, Pukhaev RV, Pukhaeva ZA (1986) Rare Noctuid species (Lepidoptera, Noctuidae) of Liakhva Reserve. Abstracts of the 1st Transcaucasian Conference on Entomology (Erevan): 153–154. [In Russian]

Pukhaev RV, Pukhaeva ZA (1989) On the fauna of Noctuid-Moths (Lepidoptera, Noctuidae) of South-Ossetian autonomous region of Georgian SSR. Abstracts of the 2st Transcaucasian Conference on Entomology (Tbilisi): 150–160. [In Russian]

Pukhaev RV, Pukhaeva ZA (1991) Noctuid-Moths of the forests of South Ossetia. Abstracts of the 6st Conference of MLI (Moscow) 1: 43–44. [In Russian]

Pukhaev RV, Pukhaeva ZA, Kabulov ZE (1987) On the fauna and biology of Noctuid-Moths of Liakhva Reserve. Reserves of Georgia (Tbilisi) 6: 234–247. [In Russian]

Pukhaeva ZA (1995) Noctuid-Moths of the southern slopes of Central Caucasus and biological rationale for pest control. Thesis of dissertation. Saint-Petersburg, 21 p. [In Russian]

Shchurov VI, Kuchmistaya EV, Vibe EN, Bondarenko AS, Skvortsov MM (2015) *Cydalima perspectalis* (Walker, 1859) – the real threat to biological diversity of natural forests of the northwest caucasus. Proceedings of the Kuban State Agrarian University 2 (53): 178–190. [In Russian]

Sinev SYu, Streltzov AN (2019) Crambidae. In: Sinev SYu (ed.) Catalogue of the Lepidoptera of Russia. Edition 2. Zoological Institute RAS, St. Petersburg, 178–196. [In Russian]

Sinev SYu, Streltzov AN, Trofimova TA (2019) Pyralidae. In: Sinev SYu (ed.) Catalogue of the Lepidoptera of Russia. Edition 2. Zoological Institute RAS, St. Petersburg, 165–178. [In Russian]

Strelitzov AN (2008) A new genus for *Glyphodes perspectalis* (Walker, 1859) (Pyraloidea, Crambidae, Pyraustinae). Eurasian Entomological Journal 7 (4): 369–372. [In Russian]

Strelitzov AN, Ustjuzhanin PYa, Yakovlev RV (2022) A new species of the Genus *Scoparia* Haworth, 1811 (Lepidoptera: Pyraloidea, Crambidae) from the Transcaucasia. Far Eastern Entomologist 457: 1–6. <https://doi.org/10.25221/fee.457.1>

Tuniyev BS, Aliev KhU, Timukhin IN (2018) *Cydalima perspectalis* (Walker, 1859) (Lepidoptera, Crambidae) in South Ossetia. Botanical Gerald of North Caucasus 1: 30–36. [In Russian]



Tuniyev BS, Lotiev KYu, Tuniyev SB, Gabaev VN, Kidov AA (2017) Amphibians and Reptiles of South Ossetia. *Nature Conservation Research* 2 (2): 1-23. [In Russian]

Ustjuzhanin PYa, Teimurov AA, Anikin VV, Matov AYu, Naydenov AE, Streltsov AN, Yakovlev RV (2022) Materials on the Lepidoptera fauna of the Dagestan Republic (Northeastern Caucasus, Russia): autumn aspect (Insecta: Lepidoptera). *SHILAP Revista de lepidopterología* 50 (198): 213–228.

Yakovlev RV, Teymurov AA, Kurbanova NS, Anikin VV, Matov AYu., Morozov PS, Naydenov AE., Spitsyn VM., Streltsov AN, Ustjuzhanin PYa (2022) Materials on the Lepidoptera fauna of the Dagestan Republic (Northeastern Caucasus, Russia): spring aspect. Families Coleophoridae, Pterophoridae, Pyralidae, Crambidae, Drepanidae, Geometridae, Sphingidae, Saturniidae, Notodontidae, Erebidae & Noctuidae. *South of Russia: ecology, development* 17 (2): 19–27.
<https://doi.org/10.18470/1992-1098-2022-2-19-27>