

Annotated checklist of the Cerambycidae (Coleoptera) of Mali

Robert H.-T. Beck	Eversbuschstrasse 83b, Munich, D-80999, Germany
Alexey M. Prozorov	University of Sciences, Techniques and Technology of Bamako, Bamako, BP 1805, Mali
Tatiana A. Prozorova	Karl Marx st., 41-14, Ulyanovsk, 432001, Russia
Mohamed M. Traore	University of Sciences, Techniques and Technology of Bamako, Bamako, BP 1805, Mali
Sékou F. Traoré	University of Sciences, Techniques and Technology of Bamako, Bamako, BP 1805, Mali
Aidas Saldaitis	Nature Research Centre, Akademijos str. 2, Vilnius, L-08412, Lithuania
Gergely Petrányi	University of Sciences, Techniques and Technology of Bamako, Bamako, BP 1805, Mali
Julia S. Volkova	Ulyanovsk State University, Universitetskaya Naberezhnaya str. 1, Ulyanovsk, 432063, Russia
Roman V. Yakovlev	Altai State University, Lenina pr. 61, Barnaul, 656049, Russia; Tomsk State University, Laboratory of Biodiversity and Ecology, Lenina pr. 36, Tomsk, 634050, Russia; Paleo Data Lab., Institute of Archaeology and Ethnography SB RAS, Kutateladze str. 7/3, Novosibirsk, 630090, Russia
Edita E. Revay	University of Sciences, Techniques and Technology of Bamako, Bamako, BP 1805, Mali
Günter C. Müller	University of Sciences, Techniques and Technology of Bamako, Bamako, BP 1805, Mali; Kuvin Center for the Study of Infectious and Tropical Diseases, Hadassah Medical School, The Hebrew University, Jerusalem, 91120, Israel

Fifty-three species of Cerambycidae were collected in Mali between 2014 and 2020 as by-catches within NIH and IVCC projects for malaria vector ecology and control, 42 of them are new records for the country. An updated list of 89 species of cerambycid beetles of Mali is presented. Two species, *Niphotragulus occidentalis* Breuning, 1977 and *Sophronica sudanica* Breuning, 1962, are so far only known from Mali.

<http://zoobank.org/4B4C8FB8-00B6-4F39-9A0F-3F8D07AC4F48>

Corresponding author: Robert H.-T. Beck (beck.robert@arcor.de)

Academic editor: M. Arnedo | Received 18 August 2022 | Accepted 2 September 2022 | Published 10 September 2022

Citation: Beck RH-T, Prozorov AM, Prozorova TA, Traore MM, Traoré SF, Saldaitis A, Petrányi G, Volkova JuS, Yakovlev RV, Revay E, Müller GC (2022) Annotated checklist of the Cerambycidae (Coleoptera) of Mali. Acta Biologica Sibirica 8: 317–365. <https://doi.org/10.14258/abs.v8.e20>

Keywords

climate change, deforestation, faunistic list, vegetation zones, West Africa

Introduction

The Republic of Mali is a fast-developing West African country with an area of 1.2 million km² and a quickly growing population of approximately 20 million (UN 2022). The territory stretches from north to south for a 1600 km distance and occupies five main ecological climate and vegetation zones (Rian et al. 2009; Coulibaly et al. 2016; Tandina et al. 2018; Sylla 2020): Saharan, Sahelian, Sudanian, Sudano-Guinean and Guinean zones (Fig. 1).



Figure 1. Vegetation zones and annual precipitation with the collecting sites (red for the original data, blue for the references) on a satellite view of Mali (Google, 2022). Locus “Kangaba+” includes closely situated Déguéla, Kangaba, Kenieroba and Ouronina

The vast northern areas include extreme desert, semidesert, and some oasis habitats with a very low population density of nomadic and semi-nomadic herders. Here the climate is typical of the Sahara Desert, and July is the hottest month. It rarely rains, though every few years some sporadic and local downpours can occur. More to the south, in the center of the country, in July, August, and September, the temperature decreases steadily because of the regular showers fed by the African summer monsoon, which brings humid currents from the Atlantic Ocean. Here the climate is also tropical in winter, and the hottest months are April, May, and June, when the maximum

temperature often exceeds 40°C, with peaks of 48°C, this area is a part of the Sahel. In the southern region, rainfall exceeds 500 mm per year (south of a line along the border of Senegal to Mopti), and in the extreme south (in a line from Bamako to Sikasso), it exceeds 1,000 mm with the rainy season lasting from June to November (WMO 2021).

The country as a whole suffers from desertification slowly moving from north to south and intensive human-caused accelerating deforestation, especially in the southern regions (FAO UN / UN EP 2020). The climate change is pushing on an annual basis the Sahara Desert up to 48 km towards the south in the Sahel, and once reliable rainy seasons are now fluctuating in time and total annual amounts of precipitation significantly, resulting in more intensive use of remaining ecosystems suitable for agriculture (Nicholson 2000; Nicholson 2001; Thomas et Nigam 2018).

In Mali deforestation is mainly driven by replacing woodlands with agricultural areas, but also logging for timber and clearing bushland and remaining forests for charcoal production in areas that can later not be used for agriculture like steep slopes and rocky areas is common practice (Mensah et al. 2020). The degradation of the remaining forests is further driven by overgrazing of cattle, sheep, and goats and burning annually the undergrowth at the end of the dry season to enhance fresh growth for grazing (Kiyani et al. 2017). In Malian villages firewood is still the main energy source for cooking and forests are accordingly cleaned out of any type of dead wood (Morton 2007). Accordingly, insect groups that are dependent on mature woodlands, especially fallen trees are strongly impacted and their habitats are quickly vanishing. One of the vulnerable groups is Cerambycidae, especially species that are xylophagous and develop inside the wood and roots of large trees.

Very little was so far known about the Cerambycidae of Mali – only one article was devoted to the fauna of the country by Villiers (1962) mentioning 31 taxa. Other records from Mali are sporadic without an attempt to summarize the local fauna (Pic 1898; Pic 1932; Lepesme 1952; Lepesme et Breuning 1958; Villiers 1962; Breuning 1962b; Breuning 1977; Breuning 1963; Quentin et Villiers 1971; Quentin et Villiers 1972; Sudre et al. 2007; Juhel et Bentanachs 2009; Teocchi et al. 2016; Juhel 2017; Bjørnstad 2019).

Material and methods

The material of the present study was obtained from by-catches from long-term malaria research conducted from 2008 to 2021 by the University of Sciences, Techniques, and Technologies of Bamako (USTTB). Adults were collected with sweep nets, Malaise traps, UV-CDC traps, and different types of larger UV traps (Kline et al. 2011; Sheikh et al. 2016) and by other miscellaneous methods by the local and visiting entomologists around the following localities (Fig. 1): Bamako, Déguéla, Kangaba, Kenieroba, Mopti, Ouronina, Sikasso. The collected material was identified and stored partially in the first author's collection and collection of the USTTB.

Species list

Subfamily *Philinae*, Tribe *Philini*

1. *Doesus telephoroides* Pascoe, 1862

Distribution in Africa: Benin, Cameroon, Central African Republic, Congo, Ethiopia, Mali, Nigeria, Senegal, Sudan; also recorded from India and Laos (Pascoe 1862; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo, 12.VI.1960 (Villiers 1962).

Subfamily *Prioninae*, Tribe *Prionini*

2. *Polyarthron pectinicornis* (Fabricius, 1793)

Synonymy follows Tavakilian 2022:

= *Polyarthron Faure-Bigueti* Pic, 1898

= *Prionus pectinicornis* Gaillardii Lameere, 1912

Distribution in Africa: Chad, Mali, Mauritania, Morocco, Niger, Senegal, Western Sahara (Fabricius 1793; Pic 1898; Lameere 1912a; Rungs 1947; Quentin 1956; Villiers 1961; Löbl et Smetana 2010; Tavakilian 2022).

Collection site in Mali: Timbuktu (Pic 1898).

Subfamily Prioninae, Tribe Macrotomini**3. *Mallodon downesi* Hope, 1843**

Synonymy follows Tavakilian 2022:

= *Mallodon laevipenne* White, 1853

= *Mallodon costipenne* White, 1853

= *Mallodon plagiatum* Thomson, 1867

= *Mallodon proximum* Thomson, 1867

Distribution in Africa: Angola, Benin, Burundi, Cameroon, Central African Republic, Congo, DR Congo, Equatorial Guinea, Ethiopia, Gabon, the Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast, Kenya, Liberia, Malawi, Mali, Mozambique, Namibia, Nigeria, Rwanda, São Tomé and Príncipe, Senegal, Sierra Leone, South Africa, Sudan, Tanzania, Togo, Uganda, Zambia, Zimbabwe; also recorded from Comoros, Costa Rica, Madagascar (Hope 1843; White 1853; Thomson 1867; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo, 2.VII.1956 (Villiers 1962), Bamako, 20.VIII.2018 (original data).

4. *Macrotoma palmata* (Fabricius, 1792)

Synonymy follows Tavakilian 2022:

= *Prionus senegalensis* Olivier, 1795

= *Prionus spinipes* Illiger, 1805

= *Macrotoma humeralis* White, 1853

= *Macrotoma Valida* Thomson, 1877

= *Macrotoma palmata* var. *rugulosa* Kolbe, 1894

= *Macrotoma palmata* var. *brevipes* Kolbe, 1894

= *Macrotoma Böhmi* Reitter, 1903

Distribution in Africa: Algeria, Angola, Benin, Botswana, Burundi, Cameroon, Central African Republic, Chad, Congo, Djibouti, DR Congo, Egypt, Equatorial Guinea, Eritrea, Gabon, Gambia, Ghana, Guinea, Guinea-Bissau, Ivory Coast, Kenya, Malawi, Mali, Mauritania, Morocco, Mozambique, Namibia, Nigeria, Rwanda, Senegal, Sierra Leone, Somalia, South Africa, Sudan, Ethiopia, Tanzania, Togo, Uganda, Zambia, Zimbabwe; also recorded from Mauritius, Saudi Arabia, Yemen (Fabricius 1792; Olivier 1795; Illiger 1805; White 1853; Thomson 1877; Kolbe 1894; Reitter 1903; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo; Niafunke, 5.VII.1960 (Villiers 1962); Déguéla, 12.VI.2016 (original data).

5. *Aulacopus reticulatus* (Audinet-Serville, 1832)

Synonymy follows Tavakilian 2022:

= *Aulacopus natalensis* White, 1853

= *Aulacopus natalensis* var. *impressicollis* Kolbe, 1898

Distribution in Africa: Angola, Botswana, Central African Republic, Congo, DR Congo, Ethiopia, Ghana, Guinea, Kenya, Mali, Mozambique, Senegal, Somalia, South Africa, Tanzania, Togo, Uganda, Zambia, Zimbabwe (Audinet-Serville 1832; White 1853; Kolbe 1898; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962).



Figure 2. Figures 2–4. Habitats near collection sites: **2.** Open rocky savanna nearby Ouronina. **3.** Woody lowland near Ouronina. **4.** Collecting on a white screen in the middle of a riverine forest nearby Déguéla.

6 . *Navosomopsis feisthameli* (Buquet, 1860)

Synonymy follows Tavakilian 2022:

= *Macrotoma novemcostata* Quedenfeldt, 1882

= *Macrotoma (Navosomopsis) ivoriensis* Lepesme, 1953

= *Macrotoma (Navosomopsis) ebororae* Gilmour, 1956

Distribution in Africa: Angola, Benin, Cameroon, Central African Republic, Congo, DR Congo, Equatorial Guinea, Gabon, Ghana, Guinea, Ivory Coast, Nigeria, Tanzania, Togo (Buquet 1860; Quedenfeldt 1882; Lepesme 1953b; Gilmour 1956b; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Kangaba, 3–7.VII.2016.

Subfamily Prioninae, Tribe Acanthophorini

7 . *Tithoes confinis* (Lapote de Castelnau, 1840)

Synonymy follows Tavakilian 2022:

= *Tithoes mandibularis* Thomson, 1877

= *Tithoes Intermedius* Thomson, 1877

= *Tithoës crassipes* Quedenfeldt, 1882

= *Tithoes falcatus* Kolbe, 1898

= *Tithoes gularis* Kolbe, 1898

= *Tithoes gnatho* Kolbe, 1898

= *Tithoes longicornis* Kolbe, 1898

Distribution in Africa: Angola, Botswana, Cameroon, Central African Republic, Chad, Djibouti, DR Congo, Eritrea, Ethiopia, Gabon, Gambia, Guinea, Kenya, Malawi, Mali, Mauritania, Mozambique, Namibia, Senegal, Somalia, South Africa, Sudan, Tanzania, Uganda, Zambia, Zimbabwe; also recorded from Saudi Arabia (Laporte de Castelnau 1840; Thomson 1877; Quedenfeldt 1882; Kolbe 1898; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962); Kangaba 5.vii.2016 (original data).

Subfamily Prioninae, Tribe Cantharocnemini

8 . *Cantharocnemis* (*Cantharoplatys*) *plicipennis* Fairmaire, 1887

Distribution in Africa: Cameroon, Central African Republic, Congo, DR Congo, Gabon, Ghana, Ivory Coast, Sierra Leone, Tanzania (Fairmaire 1887b; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, 5.VI.2016.

9 . *Cantharocnemis spondylioides* Audinet-Serville, 1832

Synonymy follows Tavakilian 2022:

= *Hoploscelis lucanoides* Audinet-Serville, 1832

= *Cantharocnemis variolosus* Fairmaire, 1882

= *Cantharocnemis latibula* Fairmaire, 1882

= *Cantharocnemis obockianus* Fairmaire, 1890

= *Cantharocnemis modestus* Fairmaire, 1897

= *Cantharocnemis (Cantharocnemis) Gahani* Lameere, 1902

= *Cantharocnemis Grandidieri* Lameere, 1912

= *Cantharocnemis (Cantharofodius) migsi* Gilmour, 1956

= *Cantharocnemis arabicus* Fuchs, 1969

Distribution in Africa: Angola, Botswana, Burkina Faso, Djibouti, Egypt, Eritrea, Ethiopia, Guinea, Kenya, Mali, Mauritania, Namibia, Niger, Senegal, Somalia, South Africa, Sudan, Tanzania, Uganda, Yemen, Zimbabwe; also recorded from Oman, Saudi Arabia and Yemen (Audinet-Serville 1832; Fairmaire 1882a; Fairmaire 1882b; Fairmaire 1890; Fairmaire 1897; Lameere 1902; Lameere 1912b; Gilmour 1956b; Fuchs 1969; Quentin et Simonetta 1992; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Unknown (Quentin et Simonetta 1992; Adlbauer et Beck 2015); Sikasso, VIII.2015 (original data).

Subfamily C erambycinae, Tribe Oemini**10 . *Calybistum lugubre* (Olivier, 1790)**

Synonymy follows Tavakilian 2022:

= *Calybistum Fuliginosum* Thomson, 1878

Distribution in Africa: Central African Republic, Ethiopia, the Gambia, Ghana, Ivory Coast, Kenya, Mali, Nigeria, Senegal, Sudan, Uganda (Olivier 1790; Thomson 1878; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 7.VIII.1956; Dogo (Villiers 1962); Ouronina, VIII.2020 (original data).

11 . *Enicoeme krelli* Adlbauer, 2003

Distribution in Africa: Ivory Coast, Nigeria (Adlbauer 2003; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Mopti, VI.2014.

12 . *Paroeme flava* (Thomson, 1858)

Synonymy follows Tavakilian 2022:

= *Paroeme laevicollis* Aurivillius, 1927

Distribution in Africa: Cameroon, Congo, Gabon, Mali, Nigeria (Thomson 1858; Aurivillius 1927; Breuning 1962c; Villiers 1962; Tavakilian 2022).

Collection site in Mali: Farimaké – Dioura, 19.VII.1954 (Villiers 1962).

13. *Hypoeshrus* (*Tibestia*) *dallonii* Peyerimhoff, 1936

Synonymy follows Tavakilian 2022:

= *Hypoeshrus strigosus* v. *rungsi* Lepesme et Breuning, 1955

= *Hypoeshrus strigosus* v. *peyerimoffi* Lepesme et Breuning, 1955

= *Hypoeshrus abyssinicus* v. *gyllenhali* Lepesme et Breuning, 1955

= *Hypoeshrus abyssinicus* sbsp. *dallonii* v. *mirei* Lepesme et Breuning, 1955

= *Hypoeshrus abyssinicus* sbsp. *dallonii* v. *wittei* Lepesme et Breuning, 1955

= *Hypoeshrus strigosus* v. *pallidipes* Quentin, 1956

Distribution in Africa: Algeria, Burkina Faso, Chad, DR Congo, Ethiopia, Kenya, Libya, Mali, Mauritania, Morocco, Niger, Senegal, Sudan, Uganda (Peyerimhoff 1936; Lepesme et Breuning 1955c; Quentin 1956; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Unknown (Adlbauer et Beck 2015).

Subfamily Cerambycinae, Tribe O브리ini

14. *Ossibia fuscata* (Chevrolat, 1856)

Synonymy follows Tavakilian 2022:

= *Adiaphorus crinitus* Fåhraeus, 1872

= *Obriaccum Senegalense* Thomson, 1878

= *Ossibia fuscata* v. *rubra* Quentin, 1956

Distribution in Africa: Chad, Djibouti, Eswatini, Ethiopia, Guinea-Bissau, Kenya, Malawi, Mali, Mauritania, Mozambique, Namibia, Niger, Nigeria, Senegal, Sierra Leone, Somalia, South Africa, Sudan, Tanzania, Zimbabwe; also recorded from Yemen (Chevrolat 1856; Fåhraeus 1872; Thomson 1878; Lepesme 1952a; Quentin 1956; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962).

15. *Ossibia murina* (Gerstäcker, 1855)

Distribution in Africa: Chad, Ethiopia, Ivory Coast, Kenya, Mali, Mozambique, Namibia, Niger, South Africa, Sudan, Tanzania, Zimbabwe (Gerstäcker 1855; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 13.VII.1956 (Villiers 1962).

16. *Oxilus terminatus* Buquet, 1859

Synonymy follows Tavakilian 2022:

= *Oxilus terminatus* v. *miréi* Quentin, 1956

= *Oxilus terminatus* var. *abyssinicus* Breuning, 1957

Distribution in Africa: Chad, Ethiopia, Kenya, Mali, Mozambique, Senegal, Sudan, Uganda (Buquet 1859; Quentin 1956; Breuning 1957a; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 7.VII.1956(Villiers 1962).

Subfamily Cerambycinae, Tribe Xystrocerini

17. *Xystrocera dispar* (Fåhraeus, 1872)

Synonymy follows Tavakilian 2022:

= *Xystrocera curticollis* Fairmaire, 1882

= *Xystrocera nitidiventris* Fairmaire, 1887

= *Xystrocera parvicollis* Fairmaire, 1892

Distribution in Africa: Botswana, Burkina Faso, Central African Republic, Chad, Djibouti, DR Congo, Ethiopia, Kenya, Mozambique, Namibia, Somalia, South Africa, Sudan, Tanzania, Uganda, Zimbabwe and Mali (**new record**); also recorded from Saudi Arabia (Fåhraeus 1872; Fairmaire 1882; Fairmaire 1887a; Fairmaire 1892b; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Ouronina, 5.VI.2016.

Taxonomic note: Externally similar to 19. *Xystrocera vittata* (Fabricius, 1792), species relation should be reconsidered.

18. *Xystrocera nigrita* Audinet-Serville, 1834

Distribution in Africa: Angola, Benin, Burundi, Cameroon, Central African Republic, Chad, Congo, DR Congo, Ethiopia, Gabon, Ghana, Guinea, Guinea-Bissau, Ivory Coast, Kenya, Liberia, Mali, Mozambique, Namibia, Niger, Nigeria, Rwanda, Senegal, Sierra Leone, Sudan, Tanzania, Togo, Uganda (Audinet-Serville 1834; Villiers 1962; Martins 1980; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962); Ouronina, 5.VI.2016, VIII.2020 (original data).

19. *Xystrocera vittata* (Fabricius, 1792)

Synonymy follows Tavakilian 2022:

= *Xystrocera senegalensis* Klug, 1835

Distribution in Africa: Angola, Cameroon, Congo, Ghana, Guinea, Guinea-Bissau, Ivory Coast,

Kenya, Liberia, Malawi, Mali, Mozambique, Senegal, Sierra Leone, Sudan, Togo (Fabricius 1792; Klug 1835; Villiers 1962; Quentin et Villiers 1979; Martins 1980; Joly et al. 2008; Tavakilian 2022).

Collection site in Mali: Farimaké - Dioura, 6.VII.1954 (Villiers 1962).

Taxonomic note: Externally similar to 17. *Xystrocera dispar* (Fahraeus, 1872), species relations should be reconsidered.

Subfamily Cerambycinae, Tribe Cerambycini

20 . *Taurotagus griseus* (Guérin-Méneville , 1844)

Synonymy follows Tavakilian 2022:

= *Taurotagus Auberti* Fairmaire, 1892

= *Taurotagus Greenfieldi* Gahan, 1894

Distribution in Africa: DR Congo, Djibouti, Egypt, Ethiopia, Ivory Coast, Kenya, Mali, Senegal, Somalia, Sudan, Tanzania; also recorded from Yemen (Guérin-Méneville 1844; Fairmaire 1892; Gahan 1894; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962).

21 . *Taurotagus impressus* Duffy, 1955

Distribution in Africa: Rwanda (Duffy 1955; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015; Ouronina, VIII.2020.

22 . *Neoplocaederus glabricollis* (Hope, 1843)

Distribution in Africa: Central African Republic, Liberia (Hope 1843; White 1853; Teocchi 1993; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015.

23 . *Neoplocaederus viridescens* (Atkinson, 1953)

Distribution in Africa: DR Congo, the Gambia, Ghana, Ivory Coast, Senegal, Sierra Leone (Atkinson 1953; Adlbauer 1993; Adlbauer 2000; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VII.2016.

24 . *Neoplocaederus denticornis* (Fabricius, 1801)

Synonymy follows Tavakilian 2022:

= *Plocaederus cribrithorax* Kolbe, 1897

= *Plocaederus atlanticus* Rungs, 1953

Distribution in Africa: Angola, Cameroon, Chad, DR Congo, Djibouti, Ethiopia, the Gambia, Ghana, Guinea, Ivory Coast, Kenya, Mali, Morocco, Mozambique, Nigeria, Senegal, Sierra Leone,

Somalia, South Africa, Sudan, Tanzania, Togo, Zimbabwe; also recorded from Oman, Saudi Arabia and Yemen (Fabricius 1801; Kolbe 1897; Kotán et Sama 2011; Rungs 1953; Villiers 1962; Adlbauer et Beck 2015; Trócoli 2019, Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962); Ouronina, VIII.2020.

25. *Neoplocaederus cyanipennis* (Thomson, 1861)

Distribution in Africa: Angola, DR Congo, Ethiopia, the Gambia, Ivory Coast, Niger, Senegal (Thomson 1861; Adlbauer et Beck 2015; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

26. *Derolus arciferus* (Gahan, 1891)

Synonymy follows Tavakilian 2022:

= *Derolus* (s. s.) *nigritulus* Breuning, 1978

= *Derolus* (s. s.) *girardi* Breuning, 1978

Distribution in Africa: Angola, Central African Republic, DR Congo, Equatorial Guinea, Ethiopia, Ghana, Guinea, Ivory Coast, Malawi, Mozambique, Niger, Senegal, South Africa, Sudan, Tanzania, Togo. (Gahan 1891; Breuning 1978a; Adlbauer et Beck 2015; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

27. *Derolus subaureus* (Jordan, 1894)

Synonymy follows Tavakilian 2022:

= *Derolus* (*Derolus*) *pseudoreus* Lapesme et Breuning, 1958

= *Derolus* (s. str.) *togoensis* Breuning, 1974

Distribution in Africa: Benin, Burkina Faso, Central African Republic, Ethiopia, Ghana, Ivory Coast, Kenya, Mali, Niger, Senegal, Sudan, Togo (Jordan 1894; Lapesme et Breuning 1958; Villiers 1962; Breuning 1974b; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Kouloba (Lapesme et Breuning 1958); Kara, 16.V.1959 (Villiers 1962); Ouronina 7.VI.2016 (original data).

28. *Dissaporus cachani* (Lapesme et Breuning, 1958)

Distribution in Africa: Benin, Cameroon, Gabon, Ghana, Ivory Coast, Togo (Lapesme et Breuning 1958; Adlbauer 2006b; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VI.2020.

29. *Sudreana rugosa* Adlbauer, 2006

Distribution in Africa: Burkina Faso, Ivory Coast (Adlbauer 2006a; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

Subfamily Cerambycinae, Tribe Hesperophanini

30 . *Tropicophanes fasciatus* (Billberg, 1817)

Distribution in Africa: Angola, Ivory Coast, Sierra Leone, Tanzania, Zambia (Schönherr 1817a; Villiers 1959; Mourglia et Teocci 1994; Adlbauer 2000; Delahaye 2009; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VI.2020.

Subfamily Cerambycinae, Tribe Plectogasterini

31 . *Plectogaster jordani* Heath, 1905

Distribution in Africa: Benin, Burkina Faso, Cameroon, Central African Republic, Congo, DR Congo, Gabon, Ghana, Guinea, Ivory Coast, Liberia, Nigeria, Togo, Sierra Leone (Heath 1905; Adlbauer et Delahaye 2006; Meunier 2007; Bouyer 2017; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, 15.VI.2016; Sikasso, VIII.2015.

Subfamily Cerambycinae, Tribe Phoracanthini

32 . *Cordylomera spinicornis nitidipennis* Audinet-Serville, 1834

Distribution in Africa: the Gambia, Ghana, Ivory Coast, Mali, Nigeria, Senegal, Sudan (Audinet-Serville 1833; Duffy 1952; Villiers 1962; Tavakilian 2022).

Collection site in Mali: Kara, 30.II.1957 (Villiers 1962).

Subfamily Cerambycinae, Tribe Callichromatini

33 . *Philematium festivum* (Fabricius, 1775)

Distribution in Africa: Benin, Cameroon, Central African Republic, DR Congo, Ethiopia, Gabon, Ghana, Guinea-Bissau, Ivory Coast, Liberia, São Tomé and Príncipe, Senegal, Sierra Leone, Togo, Uganda and Mali (**new record**); also recorded from Guadeloupe (Fabricius 1775; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Kenieroba, VI.2016.

34 . *Hinziellus plagiatus* (Dalman, 1817)

Synonymy follows Tavakilian 2022:

= *Prosyne tenellus* Bates, 1879

Distribution in Africa: Ivory Coast, Guinea, Nigeria, Sierra Leone, Togo (Schönherr 1817; Bates 1879; Juhel 2020; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015; Ouronina, VIII.2020.

35 . *Guitelia villeti* Oberthür, 1911

Synonymy follows Tavakilian 2022:

= *Locustipes saltator* Heller, 1919

Distribution in Africa: Ghana, Mali, Sudan, Togo (Oberthür 1911; Heller 1919; Quentin et Villiers 1971; Quentin et Villiers 1972).

Collection site in Mali: Unknown (Quentin et Villiers 1971; Quentin et Villiers 1972).

3 6 . *Helymaeus tricolor* (Guérin-Méneville, 1840)

Synonymy follows Tavakilian 2022:

= *Closteromerus Raffrayi* Fairmaire, 1882

= *Helymaeus pallidiventrtris* Jordan, 1894

= *Helymaeus erlangeri* Schmidt, 1922

= *Helymaeus testaceiventrtris* subsp. *rufescens* Schmidt, 1923

= *Helymaeus Bayeri* Burgeon, 1931

Distribution in Africa: Burkina Faso, Eritrea, Ethiopia, Kenya, Mali, Senegal, Somali, South Africa, Tanzania (Guérin-Méneville 1840; Fairmaire 1882c; Jordan 1894; Schmidt 1922; Schmidt 1923; Burgeon 1931; Juhel et Bentanachs 2009; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Unknown (Juhel et Bentanachs 2009; Adlbauer et Beck 2015).

3 7 . *Oxyprosopus coeruleus* (Olivier, 1790)

Synonymy follows Tavakilian 2022:

= *Promeces jucundus* Guérin-Méneville, 1840

= *Cerambyx Fabricii* Dalman, 1817

= *Oxyprosopus Jucundus* Thomson, 1864

Distribution in Africa: Guinea, Liberia, Mali, Senegal, Sierra Leone (Olivier 1790; Dalman 1817; Guérin-Méneville 1840; Thomson 1864; Schmidt 1922; Juhel 2017; Tavakilian 2022).

Collection site in Mali: Unknown (Juhel 2017).

Subfamily Cerambycinae, Tribe Achrysonini

3 8 . *Allogaster geniculatus* Thomson, 1864

Distribution in Africa: Ivory Coast, Senegal (Thomson 1864; Adlbauer 2000; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Kenieroba, VI.2016.

Subfamily Cerambycinae, Tribe Trachyderini

39 . *Purpuricenus decorus* (Olivier, 1800)

Synonymy follows Tavakilian 2022:

= *Acanthopterus tripunctatus* Gory, 1844

= *Philagathes Duchaussoyi* Théry, 1893

= *Purpuricenus decorus* var. *Theryi* Lepesme, 1948

= *Purpuricenus decorus* var. *Lafertei* Lepesme, 1948

= *Purpuricenus decorus* var. *Lhotei* Lepesme, 1948

Distribution in Africa: Cameroon, Guinea, Ivory Coast, Mali, Nigeria, Senegal (Olivier 1800; Gory 1844; Théry 1893; Aurivillius 1912; Lepesme 1948; Bjørnstad 2019; Tavakilian 2022).

Collection site in Mali: Unknown (Bjørnstad 2019).

Subfamily L amiinae , Tribe Sternotomini**40 . *Zographus regalis* (Brown, 1776)**

Synonymy follows Tavakilian 2022:

= *Zographus regalis* ab. *cuprea* Breuning, 1935

= *Zographus regalis* s.-sp. *Poleti* Le Moul, 1939

= *Zographus regalis* s.-sp. *Poleti* ab. *viridisparsus* Le Moul, 1939

= *Zographus regalis* s.-sp. *Poleti* ab. *viridiventris* Le Moul, 1939

= *Zographus regalis* s.-sp. *Poleti* ab. *viridicollis* Le Moul, 1939

= *Zographus regalis* s.-sp. *Dyoti* Le Moul, 1939

= *Zographus regalis* s.-sp. *Boni* Le Moul, 1939

= *Zographus regalis* s.-sp. *sangaensis* Le Moul, 1939

= *Zographus regalis* s.-sp. *Marquei* Le Moul, 1939

= *Zographus regalis* s.-sp. *dahomeyensis* Le Moul, 1939

= *Zographus regalis* s.-sp. *Favareli* Le Moul, 1939

= *Zographus regalis* s.-sp. *Favareli* ab. *pseudocuprea* Le Moul, 1939

= *Zographus regalis* s.-sp. *Favareli* ab. *viridimarginatus* Le Moul, 1939

= *Zographus regalis* s.-sp. *Favareli* ab. *viridiventris* Le Moul, 1939

= *Zographus regalis centralis* Allard, 1993

Distribution in Africa: Benin, Central African Republic, Congo, DR Congo, Gabon, Ghana, Ivory Coast, Senegal, Sierra Leone, Zambia (Brown 1776; Le Moulton 1939; Breuning 1959; Breuning 1965a; Allard 1993; Mourglia et Teocchi 1994; Adlbauer et Mourglia 1999; Delahaye 2009; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015.

41 . *Freadelpha eremita* (Westwood, 1845)

Synonymy follows Tavakilian 2022:

= *Freadelpha humeralis* Thomson, 1858

Distribution in Africa: Angola, Benin, Cameroon, Central African Republic, DR Congo, Ivory Coast, Senegal, Sierra Leone, Togo, Zambia (Westwood 1845; Thomson 1858; Allard 1993; Delahaye 2009; Teocchi et al. 2015; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, 15.VI.2016; Sikasso, VIII.2015.

Subfamily Lamiinae , Tribe Phrynetini

42 . *Eurysops esau* Chevrolat, 1855

Synonymy follows Tavakilian 2022:

= *Phryneta bupthalmus* White, 1858

Distribution in Africa: Cameroon, DR Congo, Nigeria, Sierra Leone, (Chevrolat 1855; White 1858; Breuning 1937; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

43 . *Phryneta spinator* Fabricius, 1792

Synonymy follows Tavakilian 2022:

= *Cerambix obscurus* Olivier, 1795

= *Lamia Sterilis* Schönherr, 1817

= *Phryneta tristis* Thomson, 1878

= *Phryneta obscura* var. *Ugandae* Aurivillius, 1914

Distribution in Africa: Angola, Benin, Burundi, Congo, DR Congo, Ethiopia, Ghana, Ivory Coast, Kenya, Liberia, Malawi, Mozambique, Niger, Rwanda, Senegal, Sierra Leone, Somalia, South Africa, Tanzania, Togo, Uganda, Zambia, Zimbabwe and Mali (**new record**); also recorded from Madagascar (Fabricius 1792; Olivier 1795; Schönherr 1817b; Thomson 1878; Aurivillius 1914; Teocchi et al. 2013; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Ouronina, VIII.2020.

44 . *Stenophryneta variegata* Aurivillius, 1907

Synonymy follows Tavakilian 2022:

= *Stenophrynetta cinerea* Aurivillius, 1907

Distribution in Africa: Angola, Cameroon, Ghana, Ivory Coast, Malawi, Mozambique, Senegal, Tanzania, Zambia, Zimbabwe (Aurivillius 1907; Teocchi et Sudre 2002; Delahaye 2009; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

Subfamily Lamiinae , Tribe Prosopocerini

45 . *Bangalaia nebulosa* (Quedenfeldt, 1887)

Synonymy follows Tavakilian 2022:

= *Anybostetha Quedenfeldti* Duvivier, 1892

= *Agnitogaster variegatus* Jordan, 1894

= *Prosopocera (Dalterus) Vaneyeni* Breuning, 1951

Distribution in Africa: Cameroon, Central African Republic, DR Congo, Equatorial Guinea, Gabon, Guinea, Ivory Coast, (Quedenfeldt 1887; Duvivier 1892b; Jordan 1894; Breuning 1951a; Lapesme et Breuning 1956; Breuning et Teocchi 1975b; Cools 1993; Teocchi et al. 2009; Teocchi et al. 2010; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

46 . *Prosopocera (Dalterus) inermis* Aurivillius, 1891

Synonymy follows Tavakilian 2022:

= *Prosopocera lutulenta* Gahan, 1894

= *Prosopocera minor* Hintz, 1909

= *Prosopocera (Timoreticus) inermis* ab. *viridescens* Breuning, 1936

= *Prosopocera (Dalterus) madagascariensis* Breuning, 1965

= *Prosopocera (Alphitopola) leucomarmorata* Breuning, 1966

= *Prosopocera (Dalterus) alboampliata* Breuning, 1981

= *Prosopocera* (s.s.) *serowensis* Breuning, 1986

Distribution in Africa: Angola, Botswana, Chad, DC Congo, Eswatini, Ethiopia, Kenya, Malawi, Mali, Mozambique, Namibia, Somalia, South Africa, Tanzania, Uganda, Zambia, Zimbabwe; also recorded from Madagascar (Aurivillius 1891; Gahan 1894; Hintz 1909; Breuning 1936a; Villiers 1962; Breuning 1965b; Breuning 1966; Breuning 1981b; Breuning 1986; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 16.VII.1956 (Villiers 1962).

47 . *Prosopocera anten na ta* Gahan, 1890

Synonymy follows Tavakilian 2022:

= *Prosopocera falcata* Distant, 1898

= *Prosopocera (Prosopocera) antennata* m. *quadrimaculata* Breuning, 1936

= *Prosopocera tricornis* Breuning, 1960

= *Prosopocera* (s. str.) *allardi* Breuning, 1964

= *Prosopocera* (Dalterus) *tchadensis* Breuning, 1967

= *Prosopocera* (s. s.) *paratchadensis* Breuning, 1967

= *Prosopocera (Parapocera) decelliana* Breuning, 1968

= *Prosopocera* (s. s.) *hintzi* m. *expressior* Breuning, 1969

= *Prosopocera (Prosopocera) pseudotchadensis* Breuning, 1981

= *Prosopocera* (s.s.) *antennata* ssp. *orientalis* Breuning, 1986

= *Prosopocera* (s.s.) *forchhammeri* Breuning, 1986

Distribution in Africa: Cameroon, Chad, Central African Republic, DR Congo, Ethiopia, Ivory Coast, Ghana, Guinea, Namibia, Nigeria, Senegal, Sierra Leone, South Africa, Tanzania, Togo, Zambia (Gahan 1890; Distant 1898; Breuning 1936a; Breuning 1960; Breuning 1964; Breuning 1967b; Breuning 1968a; Breuning 1969; Breuning 1981a; Forchhammer et Breuning 1986; Adlbauer et Mourglia 1999; Adlbauer et Beck 2015; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015.

48 . *Prosopocera aemilii* Aurivillius, 1907

Synonymy follows Tavakilian 2022:

= *Prosopocera Aemilii* Aurivillius, 1907

= *Prosopocera (Alphitopola) aemilii* ab. *viridissima* Breuning, 1936

= *Prosopocera (Alphitopola) aemilii* ab. *ochrescens* Breuning, 1936

= *Prosopocera (Alphitopola) aemilii* ab. *albida* Breuning, 1936

= *Prosopocera (Alphitopola) aemilii* ab. *annulata* Breuning, 1936

= *Prosopocera (Alphitopola) parajeanneli* Breuning, 1978

Distribution in Africa: Cameroon, Central African Republic, Congo, Benin, DR Congo, Equatorial Guinea, Gabon, Ghana, Ivory Coast (Aurivillius 1907; Breuning 1936a; Breuning 1978b; Teocchi 2000b; Teocchi et al. 2004; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VI.2016.

49 . *Prosopocera nivosa* (Fairmaire, 1897)

Synonymy follows Tavakilian 2022:

= *Alphitopola assimilis* Gahan, 1898

Distribution in Africa: Kenya, Somali, Tanzania (Fairmaire 1897; Gahan 1898; Teocchi 1989; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

Taxonomic note: Uncertain determination.

50 . *Prosopocera lactator* (Fabricius, 1801)

Synonymy follows Tavakilian 2022:

= *Lamia Radiata* Gory, 1835

= *Prosopocera plagiatrix* Kolbe, 1893

= *Prosopocera (Prosopocera) lactator* ab. *Duponti* Aurivillius, 1922

= *Prosopocera (Prosopocera) lactator* m. *femoralis* Breuning, 1936

= *Prosopocera lactator* v. *posticereducta* Lepesme, 1952

= *Prosopocera* (s. s.) *lactator* v. *lundae* Lepesme, 1953

= *Prosopocera* (s. str.) *lactator* v. *intermedia* Gilmour, 1956

Distribution in Africa: Angola, Benin, Botswana, Central African Republic, Congo, DR Congo Ethiopia, Guinea, Ivory Coast, Kenya, Malawi, Mali, Mozambique, Nigeria, Senegal, Sierra Leone, Somalia, Tanzania, Uganda, Zambia, Zimbabwe (Fabricius 1801; Gory 1835; Kolbe 1893b; Aurivillius 1922; Breuning 1936a; Lepesme 1952b; Lepesme 1953a; Gilmour 1956a; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 1956 (Villiers 1962).

Subfamily Lamiinae , Tribe Ceroplesini

51 . *Ceroplesis aestuans* (Olivier, 1800)

Synonymy follows Tavakilian 2022:

= *Ceroplesis aestuans* sbsp. *guineensis* Hintz, 1920

= *Ceroplesis aestuans* sbsp. *ornata* Hintz, 1920

= *Ceroplesis aestuans senegalensis* Fiedler, 1938

= *Ceroplesis aestuans dakarensis* Fiedler, 1938

= *Ceroplesis aestuans nigerica* Fiedler, 1938

= *Ceroplesis aestuans ubangiensis* Fiedler, 1938

= *Ceroplesis aestuans* v. *aegyptiacus* Gilmour, 1956

Distribution in Africa: Benin, Cameroon, Chad, Central African Republic, DR Congo, Egypt, Eritrea, Ethiopia, Ghana, Ivory Coast, Morocco, Niger, Senegal, Sudan, Togo, Uganda (Olivier 1800; Hintz 1920; Fiedler 1938; Gilmour 1956a; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: San (Villiers 1962).

52 . *Ceroplesis buettneri* (Kolbe, 1893)

Distribution in Africa: the Gambia, Ivory Coast, Senegal, Togo (Kolbe 1893a; Lepesme 1953b; Adlbauer 1993; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

Subfamily Lamiinae , Tribe Tragocephalini

53 . *Isochariesthes lesnei sudanica* Breuning, 1962

Distribution in Africa: Central African Republic, Chad, Ivory Coast, Mali, Senegal (Breuning 1962b; Villiers 1962; Teocchi 1997a; Tavakilian 2022).

Collection site in Mali: Boré (Breuning 1962b); Boré, 14.VII.1956 (Villiers 1962); Ouronina, VIII.2020 (original data).

54 . *Graciella pulchella* (Klug, 1835)

Synonymy follows Tavakilian 2022:

= *Chariesthes concinnus* Chevrolat, 1858

= *Chariesthes senegalensis* Chevrolat, 1858

= *Chariesthes elegantulus* Chevrolat, 1858

= *Graciella pulchella* v. *fasciata* Gilmour, 1956

Distribution in Africa: Benin, Cameroun, Central African Republic, Congo, DR Congo, Equatorial Guinea, Gabon, Ghana, Guinea, Guinea-Bissau, Ivory Coast, Liberia, Nigeria, São Tomé and Príncipe, Senegal, Togo, Uganda (Chevrolat 1858; Lepesme 1950; Gilmour 1956; Breuning et Teocchi 1981; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

Subfamily Lamiinae , Tribe Ancytonotini

55 . *Ancytonotus tribulus* (Fabricius, 1775)

Distribution in Africa: Angola, Cameroon, Central African Republic, Congo, DR Congo, Ethiopia, Gabon, the Gambia, Ghana, Ivory Coast, Kenya, Malawi, Mali, Morocco, Mozambique, Namibia, Nigeria, Senegal, Somalia, South Africa, Tanzania, Uganda, Zambia, Zimbabwe (Fabricius 1775; Villiers 1962; Adlbauer 1993; Adlbauer et Mourglia 1999; Delahaye 2009; Adlbauer et Beck 2015; Trócoli 2020; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962); Ouronina, VI.2020 (original data).

56 . *Lasiopezus variegator* (Fabricius, 1781)

Distribution in Africa: Cameroun, Central African Republic, Congo, DR Congo, Gabon, Ghana, Guinee, Ivory Coast, Niger, Senegal, Togo (Breuning et Teocchi 1981; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VI.2020.

57 . *Lasiopezus nigromaculatus* Quedenfeldt, 1882

Synonymy follows Tavakilian 2022:

= *Lasiopezus ambiguus* Kolbe, 1900

Distribution in Africa: Angola, Botswana, Cameroon, DR Congo, Ethiopia, Kenya, Malawi, Mozambique, Namibia, Senegal, Somalia, South Africa, Sudan, Tanzania, Togo, Zimbabwe (Quedenfeldt 1882; Kolbe 1900; Adlbauer 2018; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

58 . *Idactus cristulatus* (Fairmaire, 1886)

Synonymy follows Tavakilian 2022:

= *Lasiopezus exiguus* Quedenfeldt, 1891

= *Idactus matabelicus* Aurivillius, 1916

= *Idactus assimilis* Breuning, 1938

= *Idactus fuscovittatus* Breuning, 1971

= *Idactus paralateralis* Breuning, 1986

Distribution in Africa: Botswana, Cameroon, Central African Republic, Djibouti, Ethiopia, Ivory Coast, Malawi, Mali, Mozambique, Somalia, South Africa, Tanzania, Zimbabwe and Mali (**new record**); also recorded from Oman, Saudi Arabia, Yemen (Fairmaire 1886; Quedenfeldt 1891; Aurivillius 1916; Breuning 1938b; Breuning 1971b; Breuning 1986; Sudre et Teocchi 2002; Teocchi et al. 2010; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Ouronina, VI.2020.

Subfamily Lamiinae , Tribe Pteropliini

59 . *Sthenias cylindrator* (Fabricius, 1801)

Synonymy follows Tavakilian 2022:

= *Stenias Mionii* Guérin-Méneville, 1840

= *Stenias verticalis* Chevrolat, 1857

= *Chalarus leucaspis* Fåhraeus, 1872

Distribution in Africa: Angola, Botswana, Burundi, Cameroon, Central African Republic, Congo, DR Congo, Equatorial Guinea, Eswatini, Ethiopia, Ghana, Guinea, Ivory Coast, Liberia,

Malawi, Mozambique, Namibia, Nigeria, Rwanda, Senegal, Sierra Leone, South Africa, Sudan, Tanzania, Togo, Uganda, Zimbabwe and Mali (**new record**); also recorded from Martinique (Fabricius 1801; Guérin-Méneville 1840; Chevrolat 1857a; Fåhraeus 1872; Adlbauer et Beck 2015; Sudre et al. 2018; Tavakilian 2022).

Collection site in Mali: Kangaba, 3–7.VII.2016.

60 . *Pterotragula leucoloma* (Lapote de Castelnau , 1840)

Synonymy follows Tavakilian 2022:

= *Hatlia leucocincta* Guérin-Méneville, 1844

Distribution in Africa: Benin, Cameroon, Chad, Ivory Coast, Mali, Senegal (Lapote de Castelnau 1840; Guérin-Méneville 1844; Lepesme 1952; Villiers 1962; Teocchi 1985; Teocchi et al. 2014a; Tavakilian 2022).

Collection site in Mali: Dogo, région de Macina (Lepesme 1952); Boré, 26.VII.1956 (Villiers 1962).

61 . *Niphotragulus occidentalis* Breuning, 1977

Distribution in Africa: Endemic in Mali (Breuning 1977; Tavakilian 2022).

Collection site in Mali: Koulikoro (Breuning 1977).

Subfamily Lamiinae , Tribe Apomecyni

62 . *Apomecyna binubila* Pascoe, 1858

Synonymy follows Tavakilian 2022:

= *Apomecyna macularia* Harold, 1879

= *Apomecyna binubila* m. *conjuncta* Breuning, 1953

Distribution in Africa: Angola, Benin, Botswana, DR Congo, Ethiopia, Ghana, Guinea, Ivory Coast, Malawi, Mozambique, Namibia, Niger, Nigeria, Rwanda, Senegal, Sierra Leone, Somalia, South Africa, Sudan, Tanzania, Togo, Zambia, Zimbabwe (Pascoe 1858; Harold 1879; Breuning 1953b; Adlbauer et Beck 2015; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

63 . *Apomecyna lameerei* (Pic, 1895)

Synonymy follows Tavakilian 2022:

= *Apomecyna arabica* Breuning, 1938

= *Apomecyna arabica* ssp. *Mateui* Breuning, 1953

Distribution in Africa: Chad, Egypt, Mali, Mauritania, Morocco, Senegal, Western Sahara; also

recorded from Iran, Iraq, Israel, Pakistan, Oman, Saudi Arabia, United Arab Emirates, Yemen (Pic 1895; Breuning 1938c; Breuning 1953b; Lepesme et Breuning 1955a; Villiers 1962; Sama 2008; Löbl et Smetana 2010; Ambrus et Grosser 2012; Trócoli 2020; Tavakilian 2022).

Collection site in Mali: Boré, 13.VII.1956 (Villiers 1962).

64. *Enaretta varia* (Pascoe, 1886)

Synonymy follows Tavakilian 2022:

= *Enaretta fasciculata* Hintz, 1919

Distribution in Africa: Angola, Botswana, Cameroon, Central African Republic, DR Congo, Ethiopia, Guinea-Bissau, Ivory Coast, Kenya, Mozambique, South Africa, Tanzania, Zambia (Pascoe 1886; Hintz 1919; Adlbauer et Beck 2015; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Kangaba, 3–7.VII.2016.

Subfamily Lamiinae , Tribe Agapanthi i ni

65 . *Hyllisia ochreovittata* Breuning, 1940

Distribution in Africa: Cameroon, Chad, DR Congo, the Gambia, Ivory Coast, Mali, Senegal, Sierra Leone, Tanzania, Uganda (Breuning 1940b; Lepesme et Breuning 1955a; Villiers 1962; Teocchi et al. 2009; Tavakilian 2022).

Collection site in Mali: Diafarabé; Macina (Villiers 1962).

6 6 . *Pseudohippopsis filiformis* (Olivier, 1800)

Distribution in Africa: Angola, Central African Republic, Ethiopia, Guinea, Ivory Coast, Mali, Senegal, Somalia, Sudan, Tanzania, Zimbabwe (Olivier 1800; Guérin-Méneville 1840; Breuning 1940b; Lepesme 1952a; Adlbauer et Beck 2015).

Collection site in Mali: Dogo, région de Macina (Lepesme 1952).

Subfamily Lamiinae , Tribe Saperdini

67 . *Nupserha basalis* (Erichson, 1843)

Synonymy follows Tavakilian 2022:

= *Sphenura occipitalis* Chevrolat, 1857

= *Nupserha basipennis* Fairmaire, 1887

= *Nupserha basalis* var. *Ugandae* Aurivillius, 1914

= *Nupserha angulata* Aurivillius, 1914

= *Nupserha ventralis* Hintz, 1919 (nec Gahan, 1894)

= *Stibara sexmaculata* Pic, 1943

= *Nupserha bidentata* m. *subbasipennis* Breuning, 1950

- = *Nupserha bidentata* m. *immaculicollis* Breuning, 1950
- = *Nupserha apicalis* m. *capensis* Breuning, 1950
- = *Nupserha bidentata* v. *quadripunctata* Lepesme et Breuning, 1952
- = *Nupserha bidentata* var. *joveri* Lepesme et Breuning, 1953
- = *Nupserha bidentata* ssp. *urundiensis* Breuning, 1955
- = *Nupserha bidentata* m. *senegalensis* Breuning, 1956
- = *Nupserha bidentata* m. *nigrosternalis* Breuning, 1956
- = *Nupserha basalis* ssp. *basipennis* var. *laterimacula* Breuning, 1958
- = *Nupserha bidentata* ssp. *urundiensis* var. *kenyana* Breuning, 1958
- = *Nupserha basalis* ssp. *angulata* m. *ituriensis* Breuning, 1971
- = *Nupserha basalis* m. *atrata* Teocchi, 1994
- = *Nupserha basalis* m. *infranigra* Teocchi, 1997

Distribution in Africa: Angola, Burundi, Cameroon, Central African Republic, Chad, Congo, DR Congo, Eritrea, Ethiopia, Gabon, Guinea, Ivory Coast, Kenya, Nigeria, Senegal, Sierra Leone, South Africa, Sudan, Tanzania, Uganda and Mali (**new record**); also recorded from India (Erichson 1843; Chevrolat 1857b; Fairmaire 1887a; Aurivillius 1914; Hintz 1919; Pic 1943; Breuning 1950b; Breuning 1950c; Lepesme et Breuning 1952; Lepesme et Breuning 1953; Breuning 1955; Breuning 1956a; Breuning 1958a; Breuning 1971a; Teocchi 1994; Teocchi 1997b; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Sikasso, VIII.2015.

68. *Phytoecia* (*Pseudoblepisanis*) *cylindricollis* (Kolbe, 1893)

Synonymy follows Tavakilian 2022:

- = *Blepisanis geniculata* Kolbe, 1893
- = *Blepisanis flaviceps* Aurivillius, 1925
- = *Phytoecia* (*Pseudoblepisanis*) *geniculata* m. *subternigra* Breuning, 1950
- = *Phytoecia* (*Pseudoblepisanis*) *geniculata* m. *flaviventris* Breuning, 1950
- = *Phytoecia* (*Pseudoblepisanis*) *geniculata* m. *sassensis* Breuning, 1950
- = *Phytoecia* (*Pseudoblepisanis*) *geniculata* m. *rufolateralis* Breuning, 1950
- = *Phytoecia geniculata* m. *pusilla* Breuning, 1950
- = *Phytoecia geniculata* m. *fuscibasicornis* Breuning, 1950
- = *Phytoecia* (*Pseudoblepisanis*) *Kolbei* m. *togoensis* Breuning, 1951

= *Phytoecia (Pseudoblepisanis) Kolbei* m. *Lamottei* Breuning, 1951

= *Phytoecia (Pseudoblepisanis) Kolbei* m. *rufoantennata* Breuning, 1951

= *Phytoecia (Pseudoblepisanis) kolbei* v. *flavoabdominalis* Lepesme et Breuning, 1952

= *Phytoecia (Pseudoblepisanis) kolbei* v. *nigroscutellata* Lepesme et Breuning, 1952

= *Phytoecia (Pseudoblepisanis) Kolbei* m. *Massarti* Breuning, 1953

= *Phytoecia (Pseudoblepisanis) Kolbei* m. *rufoampliata* Breuning, 1953

= *Phytoecia (Pseudoblepisanis) kolbei* m. *atroampliata* Breuning, 1953

= *Phytoecia (Pseudoblepisanis) kolbei* m. *rubriscapa* Breuning, 1970

Distribution in Africa: Congo, DR Congo, Ghana, Guinea, Ivory Coast, Togo (Kolbe 1893a; Aurivillius 1925a; Breuning 1950b; Breuning 1950c; Breuning 1951b; Lepesme et Breuning 1952; Breuning 1953a; Breuning 1953c; Breuning 1967c; Breuning 1970; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015.

Subfamily Lamiinae , Tribe Astathini

69 . *Hecphora testator* (Fabricius, 1781)

Synonymy follows Tavakilian 2022:

= *Hecphora nitida* Aurivillius, 1920

Distribution in Africa: Cameroon, DR Congo, the Gambia, Ivory Coast, Senegal, Sierra Leone, Togo, Uganda (Fabricius 1781; Aurivillius 1920; Lepesme 1950; Breuning 1956b; Lepesme 1957; Adlbauer 1993; Teocchi et al. 1994; Sudre et Jiroux 2014; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Kenieroba, VI.2016

Subfamily Lamiinae , Tribe Desmiphorini

70 . *Sophronica sudanica* Breuning, 1962

Distribution in Africa: Endemic in Mali (Breuning 1962b; Villiers 1962; Tavakilian 2022).

Collection site in Mali: Farimake, Koubita (Breuning 1962b); Farimaké – Koubita (Villiers 1962).

71 . *Sophronica ventralis* Aurivillius, 1925

Synonymy follows Tavakilian 2022:

= *Sophronica curta* Breuning, 1939

Distribution in Africa: Congo, DR Congo, Ghana, Kenya, Mali, Nigeria, Tanzania, Uganda, Zimbabwe (Aurivillius 1925b; Breuning 1939a; Breuning 1963; Teocchi et al. 2016).

Collection site in Mali: Unknown (Teocchi et al. 2016).

Subfamily Lamiinae , Tribe Mesosini**72 . *Coptops aedificator* (Fabricius, 1793)**

Synonymy follows Tavakilian 2022:

= *Lamia ambulator* Fabricius, 1775

= *Lamia fusca* Olivier, 1797

= *Lamia villica* Olivier, 1797

= *Lachnia parallela* Audinet-Serville, 1835

= *Coptops quadristigma* Fåhraeus, 1872

= *Phymasterna inhambanensis* Bertoloni, 1876

= *Coptops bidens* Wollaston, 1877

Distribution in Africa: Cameroon, Central African Republic, Djibouti, DR Congo, Ethiopia, Gabon, the Gambia, Ivory Coast, Kenya, Liberia, Malawi, Mozambique, Namibia, Nigeria, Senegal, Somalia, South Africa, Tanzania, Uganda, Zambia, Zimbabwe and Mali (**new record**); also recorded from Andaman Islands, Cape Verde, China, Comoros, Hawaii, India, Madagascar, Mauritius, Oman, Pakistan, the Philippines, Réunion, Saudi Arabia, Seychelles, Sri Lanka, St. Helena, Taiwan, Yemen (Fabricius 1775; Fabricius 1793; Olivier 1797; Audinet-Serville 1835; Fåhraeus 1872; Bertoloni 1876; Wollaston 1877; Saha et al. 2013; Adlbauer et Beck 2015; Kariyanna et al. 2017; Kahuthia-Gatu et al. 2019; Rapuzzi et al. 2019; Tavakilian 2022).

Collection site in Mali: Kenieroba, VI.2016.

Subfamily Lamiinae , Tribe Crossotini**73 . *Corus pseudocostiger* Breuning, 1936**

Distribution in Africa: Chad, Guinea-Bissau, Ivory Coast, Sudan, Tanzania, Togo, Uganda (Breuning 1936b; Breuning 1942a; Adlbauer 1998; Adlbauer et Mourglia 1999; Teocchi et al. 2009; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015

74 . *Crossotus albicollis* Guérin-Méneville , 1844

Synonymy follows Tavakilian 2022:

= *Crossotides obtusus* Hintz, 1912

= *Crossotus senegalensis* Breuning, 1950

Distribution in Africa: Burkina Faso, Cameroon, Central African Republic, Chad, Ethiopia, Ghana, Ivory Coast, Mali, Mauretania, Morocco, Niger, Nigeria, Senegal (Guérin-Méneville 1844; Hintz 1912; Breuning 1950a; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Kenieroba, VI.2016

75 . *Crossotus plumicornis* Audinet-Serville, 1835

Synonymy follows Tavakilian 2022:

= *Crossotus Natalensis* White, 1858

= *Crossotus vestiticornis* Fairmaire, 1882

= *Crossotus plumicornis* subsp. *damarensis* Hintz, 1912

= *Crossotus excavatipennis* Breuning, 1961

Distribution in Africa: Botswana, Burkina Faso, Chad, Congo, DR Congo, Eritrea, Eswatini, Ethiopia, Guinea, Ivory Coast, Kenya, Malawi, Mali, Mauretania, Mozambique, Namibia, Niger, Senegal, Somalia, South Africa, Sudan, Tanzania, Uganda, Zambia, Zimbabwe (Audinet-Serville 1835; White 1858; Fairmaire 1882a; Hintz 1912; Breuning 1961; Villiers 1962; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Farimaké – Dioura, 13.VIII.1954 (Villiers 1962).

76 . *Crossotus stigmaticus* (Fåhraeus, 1872)

Synonymy follows Tavakilian 2022:

= *Frea (Crossotofrea) fasciculata* Breuning, 1981

Distribution in Africa: Botswana, Ethiopia, Kenya, Malawi, Mali, Mozambique, South Africa, Tanzania (Fåhraeus 1872; Lepesme 1952a; Breuning 1981a; Sudre et al. 2007; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo, région de Macina (Lepesme, 1952).

77 . *Crossotus sublineatus* Gestro, 1892

Synonymy follows Tavakilian 2022:

= *Crossotus (Crossotides) sennaariensis* Hintz, 1912

= *Crossotus Marshalli* Breuning, 1935

= *Crossotus sahariensis* Breuning, 1938

= *Crossotus somaliensis* Breuning, 1972

Distribution in Africa: Algeria, Chad, Djibouti, Ethiopia, Kenya, Mali, Morocco, Mauritania, Niger, Somalia, Sudan (Gestro 1892; Hintz 1912; Breuning 1935; Breuning 1938b; Breuning 1972; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Unknown (Sudre et al. 2007; Adlbauer et Beck 2015).

78 . *Crossotus subocellatus* (Fairmaire, 1886)

Synonymy follows Tavakilian 2022:

= *Crossotus Phillipsi* Gahan, 1896

= *Crossotus (Crossotides) Heimschi* Peyerimhoff, 1922

Distribution in Africa: Algeria, Chad, Djibouti, Egypt, Eritrea, Ethiopia, Kenya, Libya, Mali, Morocco, Mauritania, Niger, Nigeria, Senegal, Somalia, Sudan, Tanzania; also recorded from Oman, Saudi Arabia (Fairmaire 1886; Gahan 1896; Peyerimhoff 1922; Sudre et al. 2007; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Unknown (Sudre et al. 2007; Adlbauer et Beck 2015).

79. *Crossotus tuberculis* (Fairmaire, 1891)

Synonymy follows Tavakilian 2022:

= *Crossotus robustus* Jordan, 1894

= *Crossotus bimaculatus* Aurivillius, 1903

= *Crossotus albomaculatus* var. *vittatus* Aurivillius, 1914

Distribution in Africa: Benin, Cameroon, Central African Republic, DR Congo, Eritrea, Ethiopia, the Gambia, Ghana, Ivory Coast, Malawi, Mali, Morocco, Mozambique, Namibia, Niger, Nigeria, Senegal, Sudan, Tanzania, Uganda, Zambia (Fairmaire 1891; Jordan 1894; Aurivillius 1903; Aurivillius 1914; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962).

80. *Epidichostates strandi* (Breuning, 1935)

Synonymy follows Tavakilian 2022:

= *Crossotus sassensis* Breuning, 1935

Distribution in Africa: Benin, DR Congo, Gabon, Ghana, Guinea, Ivory Coast, Sierra Leone (Breuning 1935; Sudre et al. 2007; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Sikasso, VIII.2015

Subfamily Lamiinae , Tribe Eunidiini

81. *Eunidia kristenseni* Aurivillius, 1911

Synonymy follows Tavakilian 2022:

= *Eunidia kristenseni* m. *albida* Breuning, 1940

= *Eunidia kristenseni* ab. *flavomaculata* Breuning, 1942

= *Eunidia kristenseni* m. *arabica* Breuning, 1962

Distribution in Africa: Botswana, Chad, Djibouti, Eritrea, Ethiopia, Ivory Coast, Kenya, Mali, Namibia, Niger, Senegal, Somalia, Sudan, Tanzania; also recorded from Oman, Saudi Arabia, Yemen (Aurivillius 1911; Breuning 1940a; Breuning 1942c; Breuning 1962a; Villiers 1962; Ambrus et Grosser 2012; Teocchi et al. 2014b; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 7.VIII.1956 (Villiers 1962).

82 . *Eunidia flavoapicata* Breuning, 1939

Synonymy follows Tavakilian 2022:

= *Eunidia apicefulva* Breuning, 1953

Distribution in Africa: Chad, Eritrea, Ethiopia, Kenya, Mali, Nigeria, Somali, Sudan, Tanzania, Uganda (Breuning 1938a; Breuning 1939b; Breuning 1953b; Villiers 1962; Teocchi 2000a; Adlbauer et Beck 2015; Teocchi et al. 2015; Tavakilian 2022).

Collection site in Mali: Boré, 17.VII.1956 (Villiers 1962).

83 . *Eunidia caffra* Fåhraeus , 1872

Synonymy follows Tavakilian 2022:

= *Eunidia lurida* Fåhraeus, 1872

= *Eunidia modesta* Gahan, 1904

= *Eunidia albisparsa* Breuning, 1938

= *Eunidia collarti* Breuning, 1948

= *Eunidia rufifrons* Breuning, 1954

= *Eunidia paraflavicans* Breuning, 1971

= *Eunidia grisea* Breuning, 1986

Distribution in Africa: Botswana, DR Congo, Ethiopia, Kenya, Malawi, Mali, Mozambique, Somalia, South Africa, Tanzania, Uganda, Zimbabwe (Fåhraeus 1872; Gahan 1904; Distant 1904–1906; Breuning 1938d; Breuning 1948; Breuning 1954; Villiers 1962; Breuning 1971b; Breuning 1986; Adlbauer 1997; Teocchi 2000; Teocchi et al. 2015; Tavakilian 2022).

Collection site in Mali: Dogo (Villiers 1962).

84 . *Eunidia similis* Breuning, 1942

Distribution in Africa: Cameroon, Central African Republic, Ivory Coast, Nigeria, Togo (Breuning 1942b; Teocchi 2000a; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VIII.2020.

Subfamily Lamiinae , Tribe Acanthocinini**85 . *Exocentrus* (*Pseudocentrus*) *girardi* Breuning et Teocchi, 1975**

Distribution in Africa: Ivory Coast (Breuning et Teocchi 1975a; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Kenieroba, VI.2016.

86 . *Exocentrus* (*Ispaterus*) *mirei* Lepesme et Breuning, 1955

Synonymy follows Tavakilian 2022:

= *Exocentrus crassepunctus* Lepesme et Breuning, 1955

= *Exocentrus (Camptomyme) rufotibialis* Breuning, 1957

= *Exocentrus (Pseudocentrus) demangei* Breuning, 1962

= *Exocentrus (s. s.) rufohumeralis* Breuning, 1967 (nec Breuning, 1957)

= *Exocentrus (Pseudocentrus) rufohumeralis* Breuning, 1972 (nec Breuning, 1957)

= *Exocentrus (Camptomyme) rufipennis* Breuning, 1974

= *Exocentrus (Camptomyme) fuscipes* Breuning, 1981

= *Exocentrus (Camptomyme) paravariegatus* Breuning, 1981

= *Exocentrus (Pseudocentrus) neubeckeri* Breuning, 1981

Distribution in Africa: Chad, Central African Republic, Eritrea, Ethiopia, Ivory Coast, Kenya, Mali, Niger, Senegal, South Africa, Tanzania (Lepesme et Breuning 1955a; Breuning 1957b; Breuning 1962b; Breuning 1967a; Breuning 1968b; Breuning 1974a; Breuning 1977; Breuning 1981a; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Boré, 13.VII.1956 (Villiers 1962).

87 . *Exocentrus (Pseudocentrus) chevaugoni* Lepesme et Breuning, 1955

Synonymy follows Tavakilian 2022:

= *Exocentrus (Camptomyme) decorsei* Breuning, 1958

Distribution in Africa: Chad, Ivory Coast, Malawi, Mali, Niger, Senegal, Rwanda, Uganda (Lepesme et Breuning 1955b; Breuning 1958b; Breuning 1963; Sudre et Teocchi 2002; Tavakilian 2022).

Collection site in Mali: Unknown (Breuning, 1963).

Subfamily Lamiinae , Tribe Dorcasomatini

88 . *Olenecamptus macari* Lameere, 1892

Synonymy follows Tavakilian 2022:

= *Olenecamptus macari* ssp. *insularis* Breuning, 1960

Distribution in Africa: Angola, Bioko Island, Cameroon, Central African Republic, Congo, DR Congo, Guinea, Ivory Coast, Uganda (Lameere 1892; Breuning 1960; Breuning et Teocchi 1981; Teocchi et al. 2010; Tavakilian 2022) and Mali (**new record**).

Collection site in Mali: Ouronina, VI.2020.

Subfamily Lamiinae , Tribe Apomecynini

89 . *Enaretta castelnaudii* Thomson, 1864

Synonymy follows Tavakilian 2022:

= *Phoryctus mucoreus* Gerstaecker, 1871

= *Enaretta intermedia* Aurivillius, 1925

= *Fouquetia fasciculata* Pic, 1932

Distribution in Africa: Angola, Benin, Botswana, Chad, Egypt, Ethiopia, Ghana, Ivory Coast, Kenya, Malawi, Mali, Namibia, Niger, Nigeria, South Africa, Sudan, Tanzania, Uganda, Zambia, Zimbabwe (Thomson 1864; Gerstaecker 1871; Aurivillius 1925b; Pic 1932; Breuning et Teocchi 1978; Adlbauer et Beck 2015; Tavakilian 2022).

Collection site in Mali: Bamako (Pic 1932).

Discussion

Presently 89 species of Ceramycidae are known from Mali, 42 of them are new records. Being aware of the patchy collection efforts and the lack of material from the most southern regions of the country we expect a significant number of additional species which are so far not recorded. Among the recorded species 2 are suggested to be endemic to Mali, 8 show a West African distribution, 64 species are widely distributed on the continent, while 15 species are also known from outside of the continent.

All in all, little material from the West African Sahel can be found in international and local collections. Much work needs to be done to understand distribution patterns and ecological needs for almost all species. We would not be surprised if so far much less than 50% of the local species have been recorded. Mali is from a point of biodiversity one of the least researched countries (115th place in the world and 31st place in sub-Saharan Africa; Wolf et al. 2022). Even in the best-researched group of insects, butterflies, there is not a single publication dedicated to the local fauna and only a few dozen of species were according to Larsen (2005) sporadically recorded without any attempt to publish a local fauna. The authors of this study meanwhile compiled a list of more than 150 species of butterflies from Mali with minimal effort, the total number of local species should be again far above this number (original unpublished data). The same is the case for moths; only a few dozens of noctuid species are published while we compiled a preliminary list of more than 250 morphospecies (original unpublished data). It can be expected that this situation is similar to most other invertebrate groups. The lack of data can easily be explained by only a few entomologists having collected in Mali, only over short periods, during the last decades. This resulted in the false assumption that biodiversity is low in this “arid” country which even further discouraged potential research on the local fauna and biodiversity.

The ecosystems in the Sahelian belt and the deciduous scattered forests towards the south are presently threatened by rapid climate change with resulting desertification (FAO UN / UN EP 2020). This is especially the case in the Western Sahel and particularly in Mali with one of the highest population growth rates within Africa (8th place in the world with 2.9% annual population growth; World Bank 2022). There seems to be only a short window in which the present biodiversity can be recorded in these threatened and vanishing ecosystems and valuable data can be preserved for future generations and long-term projects measuring the changes in the local ecosystems over the next decades.

Acknowledgements

We are grateful to the late Prof. Dr. Vasiliy D. Kravchenko (Tel Aviv University, Israel, USTTB Bamako, Mali) for his dedicated entomological work in Mali: operating trapping systems, training local entomologists, and taking care of the collected material in harsh conditions.

We are thankful to the University of Sciences, Techniques, and Technologies of Bamako, which allowed us to use their field station near Ouronina, the numerous village councils that approved our activities, and the citizens of the surrounding villages that kindly tolerated our activities in their land.

We thank Karl Adlbauer for the species identification and valuable comments.

References

Adlbauer K (1993) Cerambyciden aus Senegambien (Coleoptera: Cerambycidae). Koleopterologische Rundschau 63: 279-292.

Adlbauer K (1997) Bockkäfer aus Zimbabwe und Transvaal, Teil III - Lamiinae (Coleoptera Cerambycidae). Lambillionea 97(2): 298-320.

Adlbauer K (1998) Cerambyciden von Kyambura Game Reserve in W-Uganda (Coleoptera, Cerambycidae). Linzer biologische Beiträge 30(2): 533-568.

Adlbauer K (2000) Bockkäfer aus dem Comoé-Nationalpark in der Elfenbeinküste (Coleoptera, Cerambycidae). Mitteilungen aus der Museum für Naturkunde in Berlin 76(1): 101-112.

Adlbauer K (2003) Neue Bockkäfer aus Westafrika, eine neue Gattung aus dem südlichen Afrika und eine neue Synonymie bei einem Taxon aus Ostafrika (Coleoptera, Cerambycidae). Linzer biologische Beiträge 35(2): 1327-1334.

Adlbauer K (2006a) Die Cerambycini der Äthiopischen Region. - Die Gattungen *Ptycholaemus*, *Striatoptycholaemus*, *Odzala*, *Sudreana* und *Djabiria*. (Cerambycidae: Cerambycinae). Coleoptera, Schwanfelder Coleopterologische Mitteilungen 10: 43-60.

Adlbauer K (2006b) Die Cerambycini der Äthiopischen Region. - Die Gattungen *Diorthus*, *Margites*, *Dissaporus*, *Graciliderolus*, *Microderolus* und *Spiniderolus*. (Cerambycidae: Cerambycinae). Coleoptera, Schwanfelder Coleopterologische Mitteilungen 10: 61-82.

Adlbauer K (2018) Zweiter Nachtrag zum Katalog und Fotoatlas der Bockkäfer Namibias (Coleoptera, Cerambycidae). Les Cahiers Magellanes 30: 44-55.

Adlbauer K, Beck R (2015) Katalog und Fotoatlas der Bockkäfer Äthiopiens (Coleoptera, Cerambycidae). Taita Publishers, Czech Republic, 312 pp.

Adlbauer K, Delahaye N (2006) The Plectogasterini Quentin et Villiers, 1969 of Gabun with revalidation of *Plectogaster puncticollis* Burgeon, 1947 and description of his hitherto unknown female. (Coleoptera, Cerambycidae, Cerambycinae). Coleoptera, Schwanfelder Coleopterologische Mitteilungen 10: 35-42.

Adlbauer K, Mourglia R (1999) Zur Bockkäferfauna von Togo II. Lamiinae (Col., Cerambycidae). Bonner Zoologische Beiträge 48(3-4): 329-352.

Allard V (1993) Les Sternotomini. Les Coléoptères du Monde, Vol. 17. Sciences Nat, 112 pp.

Ambrus R, Grosser W (2012) Contribution to the knowledge of longhorn beetles from Dhofar region

in sultanate of Oman (Coleoptera: Cerambycidae). *Humanity space - International Almanac* 1(2): 448–457.

Atkinson DJ (1953) New species of *Plocaederus* (Col., Cerambycidae) from Africa. *The Annals and Magazine of Natural History* 6(12): 223–227.

Audinet-Serville J-G (1832) Nouvelle classification de la famille des longicornes. *Annales de la Société Entomologique de France* 1(1): 118–201.

Audinet-Serville J-G (1834) Nouvelle classification de la famille des longicornes. *Annales de la Société Entomologique de France* 1(3): 5–110.

Audinet-Serville J-G (1835) Nouvelle classification de la famille des Longicornes 3ème Tribu, Lamiaires, Lamiariae. *Annales de la Société Entomologique de France* 1(4): 5–100.

Aurivillius Ch (1916) Neue oder wenig bekannte Coleoptera Longicornia. 16. *Arkiv för Zoologi* 10(19): 335–360.

Aurivillius Ch (1891) Neue Coleoptera Longicornia. III. *Entomologisk Tidskrift* 12: 97–106.

Aurivillius Ch (1903) Neue oder wenig bekannte Coleoptera Longicornia. 8. *Arkiv för Zoologi* 1: 313–328.

Aurivillius Ch (1907) Neue oder wenig bekannte Coleoptera Longicornia. 9. *Arkiv för Zoologi* 3(18): 93–131.

Aurivillius Ch (1911) Neue oder wenig bekannte Coleoptera Longicornia. 12. *Arkiv för Zoologi* 7(19): 187–228.

Aurivillius Ch (1914) Neue oder wenig bekannte Coleoptera Longicornia. 14. *Arkiv för Zoologi* 8(29): 265–318.

Aurivillius Ch (1920) Neue oder wenig bekannte Coleoptera Longicornia. 17. *Arkiv för Zoologi* 13(9): 361–403.

Aurivillius Ch (1922) Cerambycidae: Lamiinae. I. *Coleopterorum Catalogus, pars 73*. W. Junk et S. Schenkling, Berlin, 322 pp.

Aurivillius Ch (1925a) Cérambycides nouveaux du Musée du Congo Belge. *Revue de Zoologie et de Botanique Africaines* 13: 52–56.

Aurivillius Ch (1925b) Neue oder wenig bekannte Coleoptera Longicornia. 20. *Arkiv för Zoologi* 17A(12): 481–501.

Aurivillius Ch (1927) Neue oder wenig bekannte Coleoptera Longicornia. 22. *Arkiv för Zoologi* 19A(17): 525–547.

Bates HW (1879) New Genera and Species of Callichrominæ (Coleoptera Longicornia). *Cistula Entomologica* 21(2): 395–419.

Bertoloni G (1876) Descrizione di quattro specie novelle di Coelotteri Mosambicesi e notizie intorno alla *Acidalia Herbariata* F. *Memorie dell'Accademia di Scienze di Bologna* (3)7: 263–270.

Bouyer Th (2017) Révision des *Plectogaster* Waterhouse, 1881 du “groupe jordanii” (Cerambycidae, Cerambycinae, Plectogasterini). *Entomologia Africana* 22(2): 21–42.

Breuning S (1935) Novae species Cerambycidae. *Folia Zoologica et Hydrobiologica* 7(2): 153-174.

Breuning S (1936a) Études sur les Lamières (Coléop. Cerambycidae). Troisième Tribu: Prosopocerini Thomson. *Novitates Entomologicae* 3ème supplément (21-29): 157-230.

Breuning S (1936b) Novae species Cerambycidae. V. Festschrift zum 60. Geburtstage von Professor Dr. Embrik Strand 1: 274-325.

Breuning S (1937) Études sur les Lamières (Coléop. Cerambycidae). Cinquième Tribu: Phrynetini Thomson. *Novitates Entomologicae*, 3ème supplément 46-49: 271-302.

Breuning S (1938a) Longicorni (Lamiinae) della Somalia raccolti dal Maggiore Cesare Lomi durante la campagna Etiopica. *Atti del Museo Civico di Storia Naturale di Trieste* 14(6): 107-123.

Breuning S (1938b) Novae species Cerambycidae VI. Festschrift zum 60. Geburtstage von Professor Dr. Embrik Strand 4: 180-392.

Breuning S (1938c) Nouveaux Cerambycidae (Col.). *Novitates Entomologicae* 9: 30-63.

Breuning S (1938d) Spedizione Zoologica del Marchese Saverio Patrizi nel Basso Giuba e nell'Oltregiuba. Giugno-Agosto 1934 XII. Quelques nouvelles espèces de Lamiinae (Col. Ceramb.) de l'Afrique orientale. *Annali del Museo Civico di Storia Naturale Giacomo Doria* 58(16): 204-206.

Breuning S (1939a) Neue Lamiinae (Cerambycidae, Coleoptera) aus dem Museo civico di Storia naturale in Genua. *Memorie della Società Entomologica Italiana* 18: 53-79.

Breuning S (1939b) Novae species Cerambycidae VII. Festschrift zum 60. Geburtstage von Professor Dr. Embrik Strand 5: 144-290.

Breuning S (1940a) Lamiini (Coleopt.) dell'Africa orient. Italiana esistenti nel Museo di Milano. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano* 79(19): 157-174.

Breuning S (1940b) Novae species Cerambycidae. IX. *Folia Zoologica et Hydrobiologica* 10(1): 115-214.

Breuning S (1942a) Études sur les Lamières (Coleop. Cerambycidae). Dixième tribu: Crossotini Thoms. *Novitates Entomologicae* (73-84): 8-101.

Breuning S (1942b) Novae species Cerambycidae XI. *Folia Zoologica et Hydrobiologica* 11: 113-175.

Breuning S (1942c) Über einige neue ostafrikanische Cerambyciden (Lamiinae) des Museo civico di Storia naturale, Trieste. *Atti del Museo Civico di Storia Naturale di Trieste* 15(4): 87-94.

Breuning S (1948) Nouvelles formes de Lamières, (deuxième partie). *Bulletin du Musée Royal d'Histoire Naturelle de Belgique* 24(47): 1-47.

Breuning S (1950a) Lamières nouveaux de la collection Lepesme. *Longicornia* I: 511-535.

Breuning S (1950b) Nouvelles formes de Lamières, (troisième partie). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique* 26(12): 1-32.

Breuning S (1950c) Nouvelles formes de Lamières du Congo Belge (Cerambycidae). *Annales du*

Musée Royal du Congo Belge, série in 8°, Sciences Zoologiques 4: 7–46.

Breuning S (1951a) Lamiaires nouveaux du Congo belge (Col. Cerambycidae). *Revue de Zoologie et de Botanique Africaines* 44(2): 148–155.

Breuning S (1951b) Revision du genre *Phytoecia* Muls. (Col. Cerambycidae). *Entomologische Arbeiten aus dem Museum G. Frey* 2(2): 353–460.

Breuning S (1953a) Lamiinae. Parc National Albert I. Mission G. F. de Witte 1933–1935 79(4): 57–82.

Breuning S (1953b) Nouvelles formes de Lamiaires. (Quatrième partie). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique* 29(8): 1–38.

Breuning S (1953c) Verzeichnis der von Dr. J. Bechyné in Französisch Guinea im Jahre 1951 gesammelten Lamiinae (Col. Cerambycidae). *Entomologische Arbeiten aus dem Museum G. Frey* 4: 1–9.

Breuning S (1954) Nouvelles formes de Lamiaires (Sixième partie). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique* 30(28): 1–23.

Breuning S (1955) Contributions à l'étude de la faune entomologique du Ruanda-Urundi (Mission P. Basilewsky, 1953) – XXII. Coleoptera Cerambycidae Lamiinae. *Annales du Musée Royal du Congo Belge* 8 (Sciences Zoologiques), 36: 216–239.

Breuning S (1956a) Lamiaires nouveaux de la collection Lapesme (2e note). *Longicornia* 3: 665–702.

Breuning S (1956b) Révision des “Astathini.” *Longicornia* 3: 417–519.

Breuning S (1957a) Le genre *Oxilus* Buquet. *Bulletin de la Société Entomologique de France* 62: 196–199.

Breuning S (1957b) Nouvelles formes de Lamiaires (neuvième partie). *Bulletin de l'Institut Royal des Sciences Naturelles de Belgique* 33(8): 1–24.

Breuning S (1958a) Révision des espèces africaines du genre *Nupserha* Thoms. (Col. Cerambycidae). *Bulletin de l'Institut Français d'Afrique Noire* 20(2), série A: 369–456.

Breuning S (1958b) Révision du genre *Exocentrus* Mulsant (Col., Cerambycidae). *Bulletin of the British Museum (Natural History)* 7(5): 211–328.

Breuning S (1959) Catalogue des Lamiaires du Monde (Col. Céramb.). 2. Lieferung. Tuntzing bei München, Verlag des Museums G. Frey: 49–107.

Breuning S (1960) Neue Cerambyciden aus den Sammlungen des zoologischen Museums der Humboldt-Universität zu Berlin (Coleoptera, Cerambycidae) Zweiter Teil. *Mitteilungen aus dem Zoologischen Museum in Berlin* 36(2), 2: 463–486.

Breuning S (1961) Neue oder schlecht bekannte Cerambyciden (Col.). *Entomologische Arbeiten aus dem Museum G. Frey* 12(1): 140–160.

Breuning S (1962a) Nouveaux Lamiaires du British Museum (Natural History) (Col. Cerambycidae). *The Annals and Magazine of Natural History (Series 13)* 5: 233–241.

Breuning S (1962b) Quelques nouvelles espèces de Lamiinae de l'Ouest africain (Col. Cerambycidae). Mémoires de l'Institut Français d'Afrique Noire 24 (série A) 3: 794-798.

Breuning S (1962c) Révision du genre *Paroeme* (Col. Cerambycidae). Mémoires de l'Institut Français d'Afrique Noire 24 (série A): 1115-1120.

Breuning S (1963) Catalogue des Lamiaires du Monde (Col. Céramb.). Verlag des Museums G. Frey (7): 463-555.

Breuning S (1964) Coléoptères Cerambycidae nouveaux des collections du Musée Royal de l'Afrique Centrale, à Tervuren. Revue de Zoologie et de Botanique Africaines 69(3/4): 369-376.

Breuning S (1965a) Contribution à la faune du Congo (Brazzaville) (Mission A. Villiers et A. Descarpentries) IX - Coléoptères Cérambycides Lamiinae. Bulletin de l'Institut Français d'Afrique Noire 27 série A (3): 1057-1078.

Breuning S (1965b) Descriptions de nouveaux Lamiaires (Col. Cerambycidae). Bulletin de la Société Entomologique de Mulhouse: 45-53.

Breuning S (1966) Nouveaux Lamiaires du Musée Royal de l'Afrique centrale (Coleoptera Cerambycidae). Revue de Zoologie et de Botanique Africaines 74(1-2): 175-183.

Breuning S (1967a) Descriptions de Lamiaires africains nouveaux (Coleoptera Cerambycidae). Revue de Zoologie et de Botanique Africaines 75(1-2): 82-94.

Breuning S (1967b) Nouveaux Coléoptères Cérambycidae Lamiinae des collections du Muséum de Paris. Bulletin de l'Institut Fondamental d'Afrique Noire 29, série A (4): 1645-1654.

Breuning S (1967c) The Scientific Results of the Hungarian Soil Zoological Expedition to the Brazzaville-Congo, I. Die Arten der Familie Cerambycidae (Coleoptera). Opuscula Zoologica 7(2): 83-88.

Breuning S (1968a) Contributions à la connaissance de la faune entomologique de la Côte-d'Ivoire (J. Decelle, 1961-1964) XXI. - Coleoptera Cerambycidae Lamiinae. Annales du Musée Royal de l'Afrique Centrale série 8, 165: 297-355.

Breuning S (1968b) Quelques rectifications systématiques sur les Lamiaires (Col. Cerambycidae). Bulletin de la Société Entomologique de Mulhouse: 57-58.

Breuning S (1969) Neue Lamiinae (Cerambycidae, Coleoptera) aus Ghana. Annales Historico-Naturales Musei Nationalis Hungarici 61: 267-269.

Breuning S (1970) Descriptions de Lamiinae nouveaux de l'Afrique centrale (Coleoptera Cerambycidae). Revue de Zoologie et de Botanique Africaines 82(1-2): 171-173.

Breuning S (1971a) Descriptions de Lamiaires nouveaux d'Afrique (Coleoptera Cerambycidae). Revue de Zoologie et de Botanique Africaines 83(1-2): 95-99.

Breuning S (1971b) Neue Lamiinae aus Südafrika (Coleoptera: Cerambycidae). Annals of the Transvaal Museum 27(9): 183-192.

Breuning S (1974a) Descriptions de nouvelles formes de Lamiinae d'Ethiopie (Coleoptera Cerambycidae). Revue de Zoologie et de Botanique Africaines 88(2): 413-420.

Breuning S (1974b) Neue Arten und Gattungen von Lamiinen (Coleoptera, Cerambycidae).

Mitteilungen aus dem Zoologischen Museum in Berlin 50(1): 149-165.

Breuning S (1977) Quelques rectifications systématiques sur les Lamiaires (Col. Cérambycidae). Bulletin de la Société Entomologique de Mulhouse: 59-60.

Breuning S (1978a) Coléoptères Cerambycidae nouveaux récoltés par M. Claude Girard à la Station d'Écologie tropicale de Lamto (Côte d'Ivoire). Bulletin de l'Institut Fondamental d'Afrique Noire 40, série A (4): 893-897.

Breuning S (1978b) Descriptions de Lamiaires africains nouveaux (Coleoptera Cerambycidae). Revue de Zoologie et de Botanique Africaines 92(1): 93-116.

Breuning S (1981a) Beschreibungen neuer Lamiinae des Transvaal Museums, Pretoria (Coleoptera: Cerambycidae). Folia Entomologica Hungarica 42(34), 2: 35-54.

Breuning S (1981b) Descriptions supplémentaires de Lamiinae, Sud Africains nouveaux (Coleoptera: Cerambycidae). Folia Entomologica Hungarica 42(34), 2: 55-59.

Breuning S et Teocchi P (1975a) Mises au point relatives au genre *Exocentrus* Mulsant (Coleoptera Cerambycidae Lamiinae) Synonymies, redescriptions, sous-genres nouveaux 5e note. Bulletin de l'Institut Fondamental d'Afrique Noire 37 série A (3): 641-656.

Breuning S et Teocchi P (1981) Notes concernant la bionomie de quelques Lamiaires africains. Bulletin Mensuel de la Société Linnéenne de Lyon 50(8): 254-272.

Breuning S, Teocchi P (1975b) Notes et mises au point concernant quelques *Prosopocerini* (Coleoptera Cerambycidae Lamiinae). Bulletin de l'Institut Fondamental d'Afrique Noire 37 série A (2): 345-370.

Breuning S, Teocchi P (1978) Notes et mises au point concernant quelques *Apomecynini* africains (Coleoptera Cerambycidae Lamiinae). Bulletin de l'Institut Fondamental d'Afrique Noire 40 série A (3): 605-620.

Breuning S, Villiers A (1972) Coléoptères Cerambycidae de l'Afrique orientale. Monitore Zoologico Italiano (N. S. supplemento 4) 11: 247-294.

Brown P (1776) Nouvelles illustrations de zoologie: contenant cinquante planches enlumineés d'oiseaux curieux, et qui non etés jamais décrits, et quelques de quadrupedes, de reptiles et d'insectes, avec de courtes descriptions systematiques. B. White, London, 4, 136 pp.

Buquet JBL (1859) Mémoire sur deux genres nouveaux de coléoptères de la famille des Longicornes (*Oxilus* et *Sthelenus*), suivi de la description de plusieurs espèces appartenant aux genres *Platyarthron*, *Oeme* (*Sclerocerus* Dej.), *Clytus*, *Apriona*, *Cerosterna* et *Acanthoderus*. Annales de la Société Entomologique de France 7(3): 619-636.

Buquet JBL (1860) Description de quelques Longicornes nouveaux. Annales de la Société Entomologique de France 8(3): 617-628.

Burgeon L (1931) Liste des Callichromini du Congo Belge des collections du Musée du Congo. Revue de Zoologie et de Botanique Africaines 20(2): 108-131.

Chevrolat LAA (1855) Description de douze espèces de Longicornes du vieux Calabar, à la côte occidentale d'Afrique. Revue et Magasin de Zoologie (2)7: 513-523.

Chevrolat LAA (1856) Description de longicornes nouveaux du vieux Calabar, côte occidentale

d'Afrique. Revue et Magasin de Zoologie (2)8: 566-572.

Chevrolat LAA (1857a) Description de longicornes nouveaux du vieux Calabar, côte occidentale d'Afrique. Revue et Magasin de Zoologie (2)9: 74-83.

Chevrolat LAA (1857b) Description de longicornes nouveaux du vieux Calabar, côte occidentale d'Afrique. Revue et Magasin de Zoologie (2)9: 166-171.

Chevrolat LAA (1858) Description de longicornes nouveaux du vieux Calabar, côte occidentale d'Afrique. Revue et Magasin de Zoologie (2)10: 348-358.

Cools J (1993) Liste du matériel typique conservé dans les collections entomologiques de l'Institut royal des Sciences naturelles de Belgique. Coleoptera, Cerambycoidea, Cerambycidae. VIII. Sous-famille des Lamiinae. Documents de Travail de L'Institut Royal des Sciences Naturelles de Belgique, 74, 115 pp.

Coulibaly O, Kone AK, Niaré-Doumbo S, Goïta S, Gaudart J, Djimdé AA, Piarroux R, Doumbo OK, Thera MA, Ranque S (2016) Dermatophytosis among Schoolchildren in Three Eco-climatic Zones of Mali. PLoS Neglected Tropical Diseases 10(4): e0004675.
<https://doi.org/10.1371/journal.pntd.0004675>

de Peyerimhoff P (1922) Nouveaux Coléoptères du Nord-Africain. Quarante-troisième note: Récoltes de M. le Dr Heimsch dans l'Erg Iguidi (Sahara occidental). Bulletin de la Société Entomologique de France: 318-321.

de Peyerimhoff P (1936) Mission au Tibesti. Coléoptères (avec la description d'un genre et de deux espèces nouvelles). Mémoires de l'Académie des Sciences (2)62: 74-81.

Delahaye N (2009) Premier inventaire des Cerambycidae collectés en Zambie entre 2006 et 2008 et description d'un nouveau Callichromatini Blanchard, 1845: *Dictator juheli* n. sp. (Coleoptera, Cerambycidae, Cerambycinae). Les Cahiers Magellanes 96: 1-28.

Distant WL (1898) Coleoptera collected in the Transvaal. The Annals and Magazine of Natural History (7)1: 366-378.

Distant WL (1904-1906) Insecta Transvaaliensia. A contribution to a knowledge of the entomology of South Africa. Order Coleoptera. Fam. Cerambycidae. (Longicornia.) 1(5-6): 98-252.

Duffy EAJ (1952) A synopsis of the genus *Cordylomera* Serville (Coleoptera Cerambycidae). The Annals and Magazine of Natural History 5(12), 57: 797-813.

Duffy EAJ (1955) Contribution à l'étude de la faune entomologique du Ruanda-Urundi (Mission P. Basilewsky 1953). XXI. Coleoptera Cerambycidae Cerambycinae. Annales du Musée Royal du Congo Belge 8 (Sciences Zoologiques), 36: 202-215.

Duvivier A (1892) Contributions à la faune entomologique de l'Afrique Centrale. II. Note sur les Coléoptères des vallées de l'Itimbiri-Rubi et de l'Uellé (régions du Haut-Congo) recueillis en 1890 par MM. Joseph Duvivier et Milz. Liste des espèces et descriptions nouvelles. Annales de la Société Entomologique de Belgique 36(7): 257-383.

Duvivier A (1892b) Mélanges Entomologiques. X. - Diagnoses de Coléoptères nouveaux du Congo. Annales de la Société Entomologique de Belgique 36: 56-60.

Erichson WF (1843) Beitrag zur Insecten-Fauna von Angola, in besonderer Beziehung zur geographischen Verbreitung der Insecten in Afrika. Archiv für Naturgeschichte 9: 199-267.

Fabricius JCh (1775) *Systema Entomologiae, sistens insectorum classes, ordines, genera, species, adiectis synonymis, locis, descriptionibus, observationibus*. Officina Libraria Kortii, Flensburgi et Lipsiae, 832 pp.

Fabricius JCh (1781) *Species insectorum exhibentes eorum differentias specificas, synonyma auctorum, loca natalia, metamorphosin adiectis observationibus, descriptionibus*. Hamburgi et Kilonii, Carol Ernest Bohnii 1, 552 pp.

Fabricius JCh (1793) *Entomologia systematica emendata et aucta. Secundum classes ordines, genera, species adiectis synonymis, locis, observationibus, descriptionibus*. Hafniae, C. G. Proft, 1(2), 538 pp.

Fabricius JCh (1801) *Systema eleutheratorum secundum ordines, genera, species: adiectis synonymis, locis, observationibus, descriptionibus*. Vol. 2. Bibliopoli Academici Novi, 687 pp.

Fåhraeus OI (1872) *Coleoptera Caffrariae, annis 1838–1845 a J. A. Wahlberg collecta. Longicornia descripsit. Öfversigt af Kongliga Svenska Vetenskaps-Akademiens Förhandlingar* 29(1): 45–69.

Fairmaire LMH (1882a) *Comptes rendus des séances. Rapports, lectures, communications. Bulletin de la Société Entomologique de Belgique (3ème série)* 16: XLIII–LVII.

Fairmaire LMH (1882b) *Coléoptères recueillis par M. G. Révoil chez les Çomalis. Cerambycidae*. In: Révoil G (Ed.), *Faune et Flore des Pays Çomalis (Afrique Orientale)*: 94–101.

Fairmaire LMH (1882c) *Diagnoses de Coléoptères abyssins*. *Le Naturaliste* 2(6): 48.

Fairmaire LMH (1886) *Note sur les Coléoptères recueillis par M. Laligant à Obock*. *Annales de la Société Entomologique de France* 5(6): 435–462.

Fairmaire LMH (1887a) *Coléoptères des Voyages de M. G. Révoil chez les Somâlis et dans l'intérieur du Zanguebar*. *Annales de la Société Entomologique de France* 7(6): 277–368.

Fairmaire LMH (1887b) *Diagnoses de deux Coléoptères nouveaux de l'Afrique Orientale*. *Le Naturaliste* 9, 19(2): 223.

Fairmaire LMH (1890) *Note supplémentaire sur les Coléoptères d'Obock*. *Annales de la Société Entomologique de France* 10(6): 547–554.

Fairmaire LMH (1891) *Notes sur quelques Coléoptères de l'Afrique intertropicale et descriptions d'espèces nouvelles*. *Annales de la Société Entomologique de France (septième série)* 60(1): 231–274.

Fairmaire LMH (1892) *Coléoptères d'Obock. Troisième partie*. *Revue d'Entomologie* 11: 77–127.

Fairmaire LMH (1897) *Coléoptères nouveaux de l'Afrique intertropicale et australe. 4e note*. *Annales de la Société Entomologique de France* 66: 109–155.

Fiedler H (1938) *Systematik und Zoogeographie von Ceroplessis. (afrikan. Lamiinae, Col.)*. *Deutsche entomologische Zeitschrift* 1938 (3–4): 571–616.

Food and agriculture organization of the UN / UN environment programme (2020) *The State of the World's Forests 2020. Forests, biodiversity and people*. Rome. <https://doi.org/10.4060/ca8642en>

Forchhammer P et Breuning S (1986) *Annotated list of Cerambycidae, subfamily Lamiinae collected in Tanzania, Zambia and Botswana 1969–80 with description of 86 new taxa*. *Contributions to the*

knowledge of Ethiopian Cerambycidae (Coleoptera) II. *Natura Jutlandica* 22(1): 1-72.

Fuchs E (1969) Missioni 1962 e 1965 del Prof. Giuseppe Scortecci nell'Arabia meridionale (Coleoptera, Cerambycidae: Prioninae e Cerambycinae). *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale in Milano* 109(4): 380-382.

Gahan ChJ (1890) On New Longicornia from Africa and Madagascar. *The Transactions of the Entomological Society of London* (4 ser.) 1890: 297-328.

Gahan ChJ (1891) Notes on Longicorn Coleoptera of the Group Cerambycinae, with Descriptions of new Genera and Species. *The Annals and Magazine of Natural History* 7(37): 19-34.

Gahan ChJ (1894) On some new Longicorn Coleoptera obtained by Mr. Th. Greenfield in Somali. *The Annals and Magazine of Natural History* 14(6), 84: 407-411.

Gahan ChJ (1896) On Coleoptera from Aden and Somaliland. *The Annals and Magazine of Natural History* 6(16), 108: 448-461.

Gahan ChJ (1898) Descriptions of new Longicorn Coleoptera from East Africa. *The Annals and Magazine of Natural History* 2(7), 7: 40-59.

Gerstäcker CEA (1871) Beitrag zur Insectenfauna von Zanzibar - III - Coleoptera. *Archiv für Naturgeschichte* 37(1): 42-86.

Gestro R (1892) Di alcuni Coleotteri raccolti nel paese dei Somali dall'Ing. L. Bricchetti Robecchi. *Annali del Museo Civico di Storia Naturale (Serie 2)* 12: 747-790.

Gilmour EF (1956a) New varieties of Lamiinae. *Longicornia* 3: 747-754.

Gilmour EF (1956b) Revision of the "Prioninae" of tropical and South Africa. *Longicornia* 3: 1-252.

Gory, HL (1835) Description de deux Lamia nouvelles. *Annales de la Société Entomologique de France* (1)4: 139-141.

Guérin-Méneville FE (1840) Coléoptères nouveaux, découverts au Sénégal, par M. Mion, capitaine au 3e régiment d'infanterie de la marine, commandant le fort Toubab, près de Sedhiou, sur la Cazaminza (Sénégalie). *Revue Zoologique* 3: 107-110.

Guérin-Méneville FE (1844) Iconographie du Règne Animal de G. Cuvier, ou représentation d'après nature de l'une des espèces les plus remarquables et souvent non encore figurées, de chaque genre d'animaux. Avec un texte descriptif mis au courant de la Science. *Insectes*. 7. Paris, J. B. Baillière, 576 pp.

Harold E (1879) Bericht über die von den Herren A. v. Homeyer und P. Pogge in Angola und im Lunda - Reiche gesammelten Coleopteren. *Coleopterologische Hefte* 16: 1-224.

Heath EA (1905) Descriptions of three new Beetles from the Gold Coast, and Angola, West Africa. *The Entomologist* 38(502): 74-76.

Heller KM (1919) Eine neue, Necydalis ähnliche Callichrominen-Gattung. *Entomologische Blätter* 15: 211-213.

Hintz E (1909) Beiträge zur Kenntnis der Cerambycidenfauna der deutschen Kolonien Afrikas. I. (Col.). *Deutsche entomologische Zeitschrift* 4: 556-562.

Hintz E (1912) Die Cerambycidengruppe der Crossotidae. (Col.). Deutsche entomologische Zeitschrift: 190–202.

Hintz E (1919) Cerambyciden. In: Ergebnisse der Zweiten Deutschen Zentral-Afrika-Expedition 1910–1911 unter Führung Adolf Friedrichs, Herzogs zu Mecklenburg. Band 1. Hamburgische Wissenschaftliche Stiftung, Leipzig, 599–638 pp.

Hintz E (1920) Die Cerambycidengattung *Ceroplesis* Serville (Col.). Archiv für Naturgeschichte 84 A (10): 162–176.

Hope FW (1843) On some new Insects from Western Africa. The Annals and Magazine of Natural History (71)11: 364–369.

<https://doi.org/10.2747/1548-1603.46.4.424>

Illiger JCW (1805) Zusätze, Berichtigungen und Bemerkungen zu Fabricii Systema Eleutheratorum. Magazin für Insektenkunde 4: 69–174.

Joly C, Bernaud D, Pierre J, Murphy R (2008) Second entomological mission to Malawi. Lambillionea 108(4) supp.: 3–47.

Jordan HEK (1894) On African Longicornia. Novitates Zoologicae 1(1): 139–266.

Juhel P (2017) Vingt-quatrième contribution à l'étude des Callichromatini africains: révision du Genre *Oxyprosopus* Thomson, 1864 (Cerambycidae, Cerambycinae). Les Cahiers Magellanes 27: 82–115.

Juhel P (2020) Vingt-huitième contribution à l'étude des Callichromatini africains: révision du genre *Hintziellus* Schmidt et description de quatre espèces nouvelles (Coleoptera, Cerambycidae, Cerambycinae, Callichromatini). Les Cahiers Magellanes 36: 87–100.

Juhel P, Bentanachs Calvo J (2009) Révision du genre *Helymaeus* Thomson, 1864 et des genres voisins (Coleoptera, Cerambycidae, Cerambycinae). Magellanes. Collection systématique, 22, 3–81.

Kahuthia-Gathu R, Abonyo A, Imbayi BM (2019) Composition and diversity of xylophagous and predatory beetles in *Vachellia xanthophloea* (Benth.) P. J. H. Hurter (Fabales: Fabaceae) at Kenyatta University and Mitaboni, Kenya. Journal of Asia-Pacific Biodiversity 12(2): 217–222.

Kariyanna B, Mohan M, Gupta R, Vitali F (2017) The checklist of longhorn beetles (Coleoptera: Cerambycidae) from India. Zootaxa 4345(1): 1–317.

Kiyani P, Andoh J, Lee Y, Lee, DK (2017) Benefits and challenges of agroforestry adoption: a case of Musebeya sector, Nyamagabe District in southern province of Rwanda, Forest Science and Technology 13(4): 174–180, <https://doi.org/10.1080/21580103.2017.1392367>

Kline DL, Hogsette JA, Müller GC (2011) Comparison of various configurations of CDC-type traps for the collection of *Phlebotomus papatasi* Scopoli in southern Israel. Journal of Vector Ecology 36(1): 212–218. <https://doi.org/10.1111/j.1948-7134.2011.00133.x>

Klug JCF (1835) Insecten (Coleoptera). In: Verzeichniss von Thieren und Pflanzen, welche auf einer Reise um die Erde durch Nord-Asien und die beiden Oeane in den Jahren 1828, 1839 und 1830 ausgeführt von Adolf Erman, Naturhistorischer Atlas. Berlin, 64 pp.

Kolbe HJ (1893a) Beiträge zur Kenntniss der Longicornier (Coleoptera). I. Die von Hauptmann Kling und Dr. Büttner im Hinterlande von Togo (Westafrika) gesammelten Arten. Stettiner

Entomologische Zeitung 54(1-3): 59–80.

Kolbe HJ (1893b) Beiträge zur Kenntniss der Longicornier (Coleoptera). II. Ueber die von Herrn Leopold Conradt in Deutsch-Ostafrika auf der Plantage Derema in Usambara und in den benachbarten Küstengegenden gesammelten Arten. Stettiner Entomologische Zeitung 54(7-9): 241–290.

Kolbe HJ (1894) Die Coleopteren-Fauna Central-Afrikas. I. Von Herrn Dr. F. Stuhlmann im Seengebiete (Victoria-, Albert-Edward- und Albert-See) gesammelte Longicornier. Stettiner Entomologische Zeitung 55(1-3): 50–65.

Kolbe HJ (1898) Coleopteren. Die Käfer Deutsch-Ost-Afrikas. Verlag von Dietrich Reimer, Berlin, 367 pp. <https://doi.org/10.5962/bhl.title.53492>

Kolbe HJ (1900) Ueber einige Cerambyciden aus Mhonda in Deutsch-Ost-Afrika. Berliner Entomologische Zeitschrift 45: 297–308.

Kotán A, Sama G (2011) A new species of *Neoplocaederus* Sama, 1991 from Syria (Coleoptera, Cerambycidae: Cerambycinae). *Folia Entomologica Hungarica* 72: 59–63.

Lameere AAL (1892) Nouveaux Lamiides de l’Afrique Tropicale. *Annales de la Société Entomologique de Belgique* 36: 506–510.

Lameere AAL (1902) Révision des Prionides. Troisième mémoire. – Spondylines. *Annales de la Société Entomologique de Belgique* 46(6): 303–334.

Lameere AAL (1912a) Révision des Prionides. Vingtième mémoire. – Prionines (VII). *Annales de la Société Entomologique de Belgique, Bruxelles* 56(6): 185–260.

Lameere AAL (1912b) Révision des Prionides. Vingt-deuxième Mémoire. – Addenda et Corrigenda. *Mémoires de la Société Entomologique de Belgique* 21: 113–188.

Laporte de Castelnau FL (1840) *Histoire Naturelle des Insectes Coléoptères*. P. Duménil, Paris, 2, 563 pp.

Larsen TB (2005) *Butterflies of West Africa*. Apollo Books, Stenstrup, 595 pp.

Le Moult E (1939) Quelques formes nouvelles ou peu connues du genre *Zoographus* (Coleop., Ceramb., Lamiinæ) (1re note). *Novitates Entomologicae* 12: 92–95.

Lepesme P (1950) Longicornes récoltés au Liberia par P. L. Dekeyser et B. Holas. *Bulletin de l’Institut Français d’Afrique Noire* 12(2): 395–400.

Lepesme P (1952a) Coléoptères Cérambycides récoltés au Soudan par G. Remaudière. *Bulletin de la Société Entomologique de France* 57(3): 38–41.

Lepesme P (1952b) Mission A. Villiers au Togo et au Dahomey (1950) XX. Coléoptères Cérambycides. *Bulletin de l’Institut Français d’Afrique Noire* 14(4): 1159–1169.

Lepesme P (1953a) Coleoptera Cerambycidae de l’Angola obtenus par le Musée de Dundo. *Publicações Culturais da Companhia de Diamantes de Angola* 16: 49–60.

Lepesme P (1953b) Coléoptères Cérambycides (Longicornes) de Côte d’Ivoire. *Institut Français d’Afrique Noire. Catalogues XI*, 103 pp.

- Lepesme P (1957) Coleoptera Cerambycidae récoltés par P. Lepesme et P. Cachan dans la région du Bas-Cavayilly (Côte d'Ivoire). Bulletin de l'Institut Français d'Afrique Noire 19 série A (1): 201-208.
- Lepesme P, Breuning S (1952) La Réserve Naturelle intégrale du Mt Nimba IV. - Coléoptères Cerambycidae. Mémoires de l'Institut Français d'Afrique Noire 19: 49-66.
- Lepesme P, Breuning S (1953) Lamiaires nouveaux de Côte d'Ivoire [Col. Cerambycidae]. Bulletin de la Société Entomologique de France 58(7): 98-104.
- Lepesme P, Breuning S (1955a) Coléoptères Cérambycides récoltés par MM. de Miré et Coste dans la région saharo-sahélienne. Bulletin de l'Institut Français d'Afrique Noire 17 série A (3): 847-855.
- Lepesme P, Breuning S (1955b) Lamiaires nouveaux de Côte d'Ivoire [Col. Cerambycidae Lamiinae] (2e Note). Bulletin de la Société Entomologique de France 60 (7-8): 122-128.
- Lepesme P, Breuning S (1955c) Révision des *Hypoeshrus* Thoms. (Cerambycina, Oemini) [Col.] Bulletin de la Société Entomologique de France 60(3-4): 40-45.
- Lepesme P, Breuning S (1956) Les "Sternotomini" de l'Ouest africain. Longicornia 3: 269-347.
- Lepesme P, Breuning S (1958) Les *Derolus* Africains (Coleoptera: Cerambycidae). Proceedings of the Tenth International Congress of Entomology 1: 171-181.
- Löbl I, Smetana A (2010) Chrysomeloidea. In: Löbl I, Smetana A (Eds.), Catalogue of Palaearctic Coleoptera. Vol. 6. Apollo books, Stenstrup, 924 pp.
- Martins UR (1980) Notes, Descriptions and Checklist of African *Xystrocera* (Col., Cerambycidae). Papéis Avulsos de Zoologia 33(5): 99-125.
- Mensah KE, Damnyag L, Kwabena NS (2020) Analysis of charcoal production with recent developments in Sub-Sahara Africa: a review. African Geographical Review 41(1): 35-55. <https://doi.org/10.1080/19376812.2020.1846133>
- Meunier J-Y (2007) Description de la femelle de *Chasmogaster camerunensis* Quentin et Villiers, 1969, et complément à la répartition de *Neoclosterus bernardii* Quentin et Villiers, 1969 (Coleoptera, Cerambycidae). Bulletin de la Société Entomologique de France 112(3): 371-374.
- Morton J (2007) Fuelwood Consumption and Woody Biomass accumulation in Mali, West Africa. Ethnobotany Research et Applications 5: 037-044. <https://doi.org/10.17348/era.5.0.37-44>
- Mourgliar R, Teocci P (1994) Terzo contributo alla conoscenza dei Cerambicidi (Coleoptera Cerambycidae) della Sierra Leone, con note sulle loro piante ospiti, sinonimie e descrizione di nuove specie del genere *Denticerus* Jordan, 1894. Accademia Nazionale dei Lincei 391(267): 399-421.
- Nicholson SE (2000) Land surface processes and Sahel climate. Reviews of Geophysics 38: 117-139. <https://doi.org/10.1029/1999RG900014>
- Nicholson SE (2001) Climatic and environmental change in Africa during the last two centuries. Climate Research 17: 123-144. <https://doi.org/10.3354/cr017123>
- Oberthür R (1911) Un Longicorne nouveau du Soudan français. Insecta 1(9): 183-186.
- Olivier G-A (1790) Encyclopédie Méthodique. Histoire Naturelle. Insectes. Panckoucke Libraire,

Paris 5(1): 1-368. <https://doi.org/10.5962/bhl.title.82248>

Olivier G-A (1795-1800). Entomologie ou Histoire Naturelle des Insectes, avec leurs caractères génériques et spécifiques, leur description, leur synonymie et leur figure enluminée. Coléoptères, 4. Imprimerie de Lanneau, Paris, pp. 1-124 (1795); pp. 125-492 (1800).

Olivier G-A (1797) Encyclopédie Méthodique, ou par ordre des matières; par une société de gens de lettres, de savans et d'artistes; précédée d'un vocabulaire universel, servant de table pour tout l'ouvrage, ornée des portraits de Mm. Diderot et d'Alembert, premiers éditeurs de l'Encyclopédie. Histoire Naturelle. Insectes. Histoire Naturelle, Insectes. Paris, Plomteux, Liège, Panckocke Libr. 7(2): 369-827.

Pascoe FP (1858) XVII. On New Genera and Species of Longicorn Coleoptera. Part III. The Transactions of the Entomological Society of London (2), 4(6-7): 236-266.

Pascoe FP (1862) Notices of new or little-known genera and species of Coleoptera. Part III. The Journal of Entomology 1(5) 26: 319-370.

Pascoe FP (1886) Descriptions of some new Longicornia, chiefly Asiatic and African. The Annals and Magazine of Natural History 5(17), 99: 239-246.

Pic M (1895) Longicornes de la collection H. Tournier. L'Échange, Revue Linnéenne 11(127): 75-78.

Pic M (1898) Coléoptères nouveaux de Tombouctou. Revue de Sciences du Bourbonnais et du Centre de la France 11: 164-165.

Pic M (1932) Longicornes nouveaux [Col.] Bulletin de la Société Entomologique de France 37(10): 150-151.

Pic M (1943) Opuscula martialia IX. L'Échange, Revue Linnéenne, Numéro spécial 9: 1-16.

Quedenfeldt FOG (1882) Kurzer Bericht über die Ergebnisse der Reisen des Herrn Major a. D. v. Mechow in Angola und am Quango-Strom, nebst Aufzählung der hierbei gesammelten Longicornen. Berliner Entomologische Zeitschrift 26(2): 317-362.

Quedenfeldt FOG (1887) Drei neue Cerambyciden von Kamerun. Berliner Entomologische Zeitschrift 31: 141-144.

Quedenfeldt FOG (1891) Neue Käfer von Ost-Afrika. Berliner Entomologische Zeitschrift 36(1): 167-174.

Quentin RM (1956) Contribution à l'étude des Coléoptères Cerambycidae - II Prioninae et Cerambycinae récoltés par Ph. Bruneau de Miré et Coste au Sud du Sahara français, avec description de trois formes nouvelles. Bulletin de la Société Entomologique de France 61(1-2): 40-43.

Quentin RM, Simonetta J (1992) Révision du genre *Cantharocnemis* Serville (Coleoptera: Cerambycidae). Les sous-genres *Cantharocnemis* sensu stricto et *Cantharofaedus* Gilmour. Annales de la Société Entomologique de France 28(1): 83-100.

Quentin RM, Villiers A (1971) *Paraguitelia*, nouveau genre africain de Callichromini [Col. Cerambycidae Cerambycinae], et notes sur les genres *Guitelia* et *Scalenus*. Bulletin de l'Institut Fondamental d'Afrique Noire 33, série A (1): 129-134.

Quentin RM, Villiers A (1972) Révision du genre *Guitelia* Oberthür [Col. Cerambycidae Callichromini]. Bulletin de l'Institut Fondamental d'Afrique Noire 34, série A (2): 343–351.

Quentin RM, Villiers A (1979) Coléoptères Cerambycidae récoltés au Kenya par M. Michel Boulard. Revue Française d'Entomologie 1(2): 79–85.

Rapuzzi P, Kuleshov DA, Fazal TM, Ahmed Z, Hussain A (2019) New or interesting records of longhorn beetles fauna of Pakistan (Coleoptera: Cerambycidae). Munis Entomology et Zoology 14(1): 62–79.

Reitter E (1903) Sechzehnter Beitrag zur Coleopteren-Fauna von Europa und den angrenzenden Ländern. Wiener Entomologische Zeitung 22(2): 43–46.

Rian S, Xue Y, MacDonald GM, Touré MB, Yu Y, De Sales F, Levine PA, Doumbia S, Taylor ChE (2009) Analysis of Climate and Vegetation Characteristics along the Savanna-Desert Ecotone in Mali Using MODIS Data. GIScience et Remote Sensing 46(4): 424–450.

Rungs Ch (1947) A propos de quelques coléoptères Cérambycidae du Maroc. Bulletin de la Société Entomologique de France 52(7): 97–101.

Rungs Ch (1952) Un nouveau Cérambycide marocain. Bulletin de la Société Entomologique de France 57(10): 146–147.

Saha S, Özdikmen H, Biswas MK, Raychaudhuri D (2013) Exploring Flat Faced Longhorn Beetles (Cerambycidae: Lamiinae) from the Reserve Forests of Dooars, West Bengal, India. Psyche 1–8. <https://doi.org/10.1155/2013/737193>

Sama G (2008) Family Cerambycidae. In: van Harten A (Ed.), Arthropod fauna of the United Arab Emirates. Vol. 1. Dar Al Ummah Printing, Abu Dhabi, United Arab Emirates, 754 pp.

Schmidt M (1922) Die afrikanischen Callichrominen (Col. Ceramb.) nach systematischen, phylogenetischen und geographischen Gesichtspunkten. Archiv für Naturgeschichte 88(6): 61–232.

Schmidt M (1923) Bemerkungen über Callichrominen (Col. Ceramb.). Deutsche entomologische Zeitschrift 7(2): 272–276.

Schönherr CJ (1817a) Appendix ad C. J. Schönherr Synonymiam Insectorum. Tom 1, Part 3. Sistens Descriptiones Novarum Specierum. Scaris, Lewerentziana, 266 pp.

Schönherr CJ (1817b) Synonymia insectorum, oder: Versuch einer Synonymie Aller bisher bekannten Insecten; nach Fabricii Systema Eleutheratorum etc. geordnet. Skara, Lewerentzischen Buchdrükerey, 1(3), 506 pp.

Sheikh AH, Thomas M, Bhandari R, Bunkar K (2016). Light trap and insect sampling: an overview. International Journal of Current Research 8(11): 40868–40873.

Sudre J, Jiroux E, Vitali C (2018) Descriptions et synonymies concernant quelques lamiaines africains (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 30: 22–37.

Sudre J, Teocchi P (2002) Contribution à la connaissance des longicornes du Malawi et description de nouveaux taxa (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 14: 1–38.

Sudre J, Teocchi P, Sama G, Rousset F (2007) Les genres *Crossotus*, *Biobessoides* et *Epidichostates* (Coleoptera, Cerambycidae, Lamiinae, Crossotini). Notes préliminaires pour une révision du genre *Crossotus*, avec remarques synonymiques et bionomiques sur les genre *Corus*, *Biobessoides* et

Epidichostates. Magellanes. Collection systématique 15: 1-80.

Sylla MB (2020) Rapport Final. Identification des risques climatiques rencontrés au Mali à court, moyen et long terme dans le cadre du Projet de Gestion des Risques Climatiques et d'Inondations (PGRCI) au Mali. PGRCI, Bamako, Mali, 70 pp.

Tandina F, Doumbo O, Yaro AS, Traoré SF, Parola Ph, Robert V (2018) Mosquitoes (Diptera: Culicidae) and mosquito-borne diseases in Mali, West Africa. Parasites and Vectors 11: 467. <https://doi.org/10.1186/s13071-018-3045-8>

Tavakilian G (2022) Base de données Titan sur les Cerambycides ou Longicornes. Available from <http://titan.gbif.fr/> (Accessed 5.04.2022)

Teocchi P (1985) A propos de *Pterolophia leucoloma* Cast. (Coleoptera Cerambycidae Lamiinae). Bulletin de la Société Sciences Nat 47: 2-3.

Teocchi P (1989) Réhabilitation de *Prosopocera (Alphitopola) nivosa* (Fairmaire) et remarques concernant quelques espèces affines [Coleoptera, Cerambycidae, Lamiinae]. Revue Française d'Entomologie 11(1): 15-16.

Teocchi P (1993) Plantes-hôtes et bionomie de quelques Cerambycidae africains (Coleoptera). Bulletin Mensuel de la Société Linnéenne de Lyon 62(7): 265-272.

Teocchi P (1994) Diagnose de Lamiaires africains nouveaux (Coleoptera Cerambycidae Lamiinae). Lambillionea 94(2), 1: 206-210.

Teocchi P (1997) Diagnoses, synonymies et chorologie de quelques Lamiaires africains (suite) (Coleoptera Cerambycidae). Lambillionea 97(3): 418-428.

Teocchi P (1997) Quelques rectifications systématiques sur les Lamiaires (Coleoptera Cerambycidae). Bulletin de la Société Entomologique de Mulhouse: 59-62.

Teocchi P (2000a) Note sur une petite collection de Lamiaires du Musée d'Helsinki (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 1: 1-32.

Teocchi P (2000b) Synonymies, diagnoses et bionomie de quelques Cérambycides africains (Coleoptera Cerambycidae) (suite 1). Lambillionea 100(4), 2: 639-652.

Teocchi P, Jiroux E, Sudre J (2004) Synonymies, diagnoses et bionomie de quelques Cerambycidae africains (10ème note) 1ère partie (Coleoptera, Cerambycidae). Les Cahiers Magellanes 38: 1-30.

Teocchi P, Jiroux E, Sudre J, Ture A (2009) Note concernant une petite collection de Lamiaires colligés en Côte d'Ivoire, 2e partie (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 98: 1-28.

Teocchi P, Sudre J (2002) Synonymies, diagnoses et bionomie de quelques lamiaires africains (8e note) (suite) (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 12: 1-24.

Teocchi P, Sudre J, Adlbauer K, Jiroux E (2010) Réhabilitation de la tribu des Protonarthrini Thomson, 1864. Diagnose d'un genre africain nouveau la composant, et transfert dans la tribu des Acmoderini de *Protonarthron tuberculicollis* Breuning, 1950 (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 2: 48-61.

Teocchi P, Sudre J, Jiroux E (2010) Synonymies, diagnoses et bionomie de quelques Lamiaires africains (13e note) (Coléoptères Cerambycidae Lamiinae). Les Cahiers Magellanes 107: 1-24.

Teocchi P, Sudre J, Jiroux E (2013) Synonymies, diagnoses et bionomie de quelques Lamiaires africains (18e note) (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 11: 1–31.

Teocchi P, Sudre J, Jiroux E (2014a) Synonymies, diagnoses et bionomie de quelques Lamiaires africains (19e note) (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 14: 16–50.

Teocchi P, Sudre J, Jiroux E (2014b) Synonymies, diagnoses et bionomie de quelques Lamiaires africains (20e note) (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 15: 51–95.

Teocchi P, Sudre J, Jiroux E (2015) Synonymies, diagnoses et bionomie de quelques Lamiaires africains (21e note) (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 17: 35–85.

Teocchi P, Sudre J, Jiroux E (2016) Synonymies, diagnoses et bionomie de quelques Lamiaires africains (23e note) (Coleoptera, Cerambycidae, Lamiinae). Les Cahiers Magellanes 21: 1–25.

Thomas N, Nigam S (2018) Twentieth-century climate change over Africa: Seasonal hydroclimate trends and Sahara desert expansion. *Journal of Climate* 31(9): 3349–3370.
<https://doi.org/10.1175/JCLI-D-17-0187.1>

Thomson J (1858) Voyage au Gabon. Histoire Naturelle des Insectes et des Arachnides recueillis pendant un voyage fait au Gabon en 1856 et en 1857 par M. Henry C. Deyrolle sous les auspices de MM. Le Comte de Mniszech et James Thomson précédé de l'histoire du voyage. *Archives Entomologiques*, 2, 472 pp.

Thomson J (1860–1861) Essai d'une classification de la famille des cérambycides et matériaux pour servir à une monographie de cette famille. Paris, 404 pp.

Thomson J (1864) *Systema Cerambycidarum* ou exposé de tous les genres compris dans la famille des Cérambycides et familles limitrophes. *Mémoires de la Société Royale des Sciences de Liège* 19: 1–540.

Thomson J (1867) IV. Révision du groupe des *Mallodonites* (insectes Coléoptères, Prionites, Cérambycides). *Physis Recueil d'Histoire Naturelle* 1(2): 85–106.
<https://doi.org/10.5962/bhl.title.38189>

Thomson J (1877) *Typi cerambycidarum musei Thomsoniani*. *Revue et Magasin de Zoologie* 5(40): 249–279.

Thomson J (1878) *Typi cerambycidarum musei Thomsoniani*. E. Deyrolle, Paris, 21 pp.

Trócoli S (2019) Actualización del catálogo de Longicornios de Marruecos Actualisation du catalogue des Longicornes du Maroc (Parte III/Partie III: Cerambycidae, Cerambycinae). *Revue de l'Association Roussillonnaise d'Entomologie (R.A.R.E.)* 28(3): 143–185.

Trócoli S (2020) Actualización del catálogo de Longicornios de Marruecos Actualisation du catalogue des Longicornes du Maroc (Parte IV/Partie IV: Cerambycidae: Lamiinae). *Revue de l'Association Roussillonnaise d'Entomologie (R.A.R.E.)* 29(1): 26–65.

United Nations Department of Economic and Social Affairs, Population Division (2022) *World Population Prospects 2022: Summary of Results*. UN DESA/POP/2022/TR/NO. 3.

Villiers A (1959) Coléoptères Cérambycides d'Angola (Prioninae et Cerambycinae). *Publicações Culturais da Companhia de Diamantes de Angola* 41: 25–32.

Villiers A (1961) A propos d'une série de *Polyarthron* (Col. Cerambycidae Prioninae) récoltée au

Sahara par F. Pierre. Bulletin de l'Institut Français d'Afrique Noire 23 série A (1): 63-82.

Villiers A (1962) Coléoptères Cérambycides récoltés au Mali par R. Demange. Bulletin de l'Institut français d'Afrique noire 24 série A (4): 1121-1125.

Villiers A (1968) Coléoptères Cerambycidae d'Arabie. Bulletin du Muséum National d'Histoire Naturelle de Paris 39(5): 846-850.

Wappes JE, Santos-Silva A, Galileo MHM (2018) Description of two new Bolivian Cerambycidae and a Costa Rican record for an African prionine (Coleoptera). Insecta Mundi 625: 1-9.

White A (1853). Catalogue of the coleopterous insects in the collection of the British Museum. Part 7. Longicornia I. London, 174 pp. <https://doi.org/10.5962/bhl.title.60043>

White A (1858) XXVIII. Descriptions of some apparently unrecorded species of Longicorn Beetles, belonging to the genera *Phrissoma*, *Nyphona*, etc. The Annals and Magazine of Natural History 3(2), 10: 264-276.

Wolf MJ, Emerson JW, Esty DC, de Sherbinin A, Wendling ZA, *et al.* (2022). 2022 Environmental Performance Index. Chapter 9: Biodiversity et Habitat. New Haven, CT: Yale Center for Environmental Law et Policy, 206 pp.

The World Bank (2022) Population growth (annual %) - Mali. Available from <https://data.worldbank.org/indicator/SP.POP.GROW?locations=ML> (Accessed 5.04.2022)

World Meteorological Organization (2021) 2020 State of the climate in Africa. No 1275. WMO, Geneva, 44 pp.

Wollaston ThV (1877) Coleoptera Sanctæ-Helenæ. John Van Voorst, London, 256 pp.