

# Thereuoneminae Verhoeff, 1905 (Chilopoda: Scutigermorpha: Scutigeridae), a new subfamily for the Russian fauna

Yurii V. Dyachkov

Altai State University, 61 Lenina Ave., Barnaul, 656049, Russia

*Thereuonema tuberculata* (Wood, 1862), genus *Thereuonema* Verhoeff, 1904, and subfamily Thereuoneminae Verhoeff, 1905 are recorded from Russia for the first time. A distribution map and illustrations are provided.

---

Corresponding author: Yurii V. Dyachkov (dyachkov793@mail.ru)

Academic editor: R. Yakovlev | Received 16 August 2022 | Accepted 8 September 2022 | Published 11 October 2022

<http://zoobank.org/C8404A32-A56F-4651-AF02-00B0CABE6EDA>

**Citation:** Dyachkov YuV (2022) Thereuoneminae Verhoeff, 1905 (Chilopoda: Scutigermorpha: Scutigeridae), a new subfamily for the Russian fauna. *Acta Biologica Sibirica* 8: 469–473. <https://doi.org/10.14258/abs.v8.e28>

## Keywords

biodiversity, fauna, Maritime Territory, new records, Russian Far East

## Introduction

The family Scutigeridae Leach, 1814 contains two subfamilies: Scutigerinae Leach, 1814 and Thereuoneminae Verhoeff, 1905 (Edgecombe 2011). The first one is represented by *Scutigera coleoptrata* (Linnaeus, 1758) in Russia (Sseliwanoff 1884; Muralewicz 1929; Nefediev et al. 2016; Zuev 2016; Zuev & Evsyukov 2016), while the latter subfamily has never been recorded. The present paper provides the first record of a member of Thereuoneminae, *Thereuonema tuberculata* (Wood, 1862), from Russia.

## Material and methods

The material was collected by Yu.V. Dyachkov; specimens were taken by hand, preserved in 70% ethanol, and deposited in ASU (see abbreviations below).

The taxonomy follows Edgecombe (2011). Localities were mapped with SimpleMappr (Shorthouse 2010).

The specimens were examined with an Olympus SZX16 stereo microscope and an Olympus BX51 microscope; photographs were taken using an Olympus DP74 or Olympus XC50 digital cameras. Tergite, gonopods, and legs were mounted in permanent slides using euparal.

Abbreviations: ASU – Altai State University (Barnaul, Russia), juv. – juvenile.

## Results

**Family Scutigeridae Leach, 1814**

**Subfamily Thereuoneminae Verhoeff, 1905**

**Genus *Thereuonema* Verhoeff, 1904**

***Thereuonema tuberculata* (Wood, 1862)**

Figs 1–8.

**Material.** 1♀ (ASU No. 508), 10♂♂, (ASU No. 509), 1♀ (ASU No. 510), 1♂ (ASU No. 512), 5♀♀, 3 juv. ♀♀ (ASU No. 513), 1♀ (ASU No. 514), 1♀ (ASU No. 515), Russia, Maritime Territory, Khasansky District, 8 km NE from Khasan (urban-type settlement), Golubiniy Utes Mt., N42°24'35", E130°44'59", 15–20 m, in grass, under stones, 14–17 July 2022.

**Distribution.** Eastern China (incl. Taiwan), Korean Peninsula, Japan (Würmli 1975; Stoev & Geoffroy 2004; Dunlop et al. 2017), and Russia (new); introduced to Great Britain (Barber 2011) and North America (Reeves & Miller 2022).

**Remarks.** Based on molecular data, Yang et al. (2022) hypothesized that cryptic species could exist in *T. tuberculata*. Therefore, the identity of specimens from Russia should be clarified by molecular analysis.

This species, genus *Thereuonema*, and subfamily Thereuoneminae are recorded from Russia for the first time.

## Conclusions

To date, at least two species from the family Scutigeridae are known to occur in Russia: *Scutigera coleoptrata* (subfamily Scutigerinae) and *Thereuonema tuberculata* (subfamily Thereuoneminae).




**Figure 1.** Distribution of *Thereuonema tuberculata* (Wood, 1862) in Russia.



**Figure 2. Figures 2-5.** *Thereuonema tuberculata* (Wood, 1862), dorsal view (2, 3 - male, ASU No. 512; 4, 5 - female, ASU No. 508): 2 - habitus of male; 3 - head of male, 4 - head of female, 5 - habitus of female. Scale: 2 mm (2, 5), 0.5 mm (3-4).





**Figure 3. Figures 6–8.** *Thereuonema tuberculata* (Wood, 1862) (female, ASU No. 514): **6** – tarsus of leg 10, lateral view; **7** – gonopods, ventral view; **8** – surface of stomatotergite 7, dorsal view. Scale: 0.5 mm (6), 0.2 mm (7), 0.05 mm (8).

## Acknowledgements

I wish to thank M.M. Omelko senior (Gornotaezhnoe, Russia) and M.M. Omelko junior (Vladivostok, Russia), and A.A. Fomichev (Barnaul, Russia) for their help during the expedition. I am grateful to P. Stoev (Sofia, Bulgaria) for his comments on the draft of the manuscript and to Mariia A. Iuzhakova (Tomsk, Russia) who checked the English of an advanced draft. This study was supported in the framework of “Priority-2030” Program by the Altai State University.

## References

- Barber AD (2011) *Thereuonema tuberculata* (Wood, 1863), a scutigermorph centipede from China, found in a warehouse at Swindon. *Bulletin of the British Myriapod and Isopod Group* 25: 49–50.
- Dunlop JA, Friederichs A, Langermann J (2017) A catalogue of the scutigermorph centipedes in the Museum für Naturkunde, Berlin. *Zoosystematics and Evolution* 93 (2): 281–295. <http://dx.doi.org/10.3897/zse.93.12882>
- Edgecombe GD (2011) Scutigermorpha. In: Minelli A (ed.). *Treatise on zoology—anatomy, taxonomy, biology. The Myriapoda. Vol. 1.* Brill, Leiden-Boston, 363–370.
- Muralewicz WS (1929) Scutigeridae and Lithobiidae of the Caucasian fauna. *Memoires de la Section Zoologique de la Societe des Amis des Sciences Naturelles, d’Anthropologie et d’Ethnographie* 4: 1–120. [in Russian]
- Nefediev PS, Tuf IH, Dyachkov YuV, Efimov DA (2016) First record of *Scutigera coleoptrata* (Linnaeus, 1758) in the south of Western Siberia, Russia (Chilopoda: Scutigermorpha: Scutigeridae). *Biological Bulletin of Bogdan Chmelnskiy Melitopol State Pedagogical University* 6 (1): 428–432.
- Reeves WK, Miller MM (2022) *Thereuonema tuberculata* (Wood, 1862) (Chilopoda, Scutigermorpha, Scutigeridae) from forested habitats in North America. *Check List* 18 (3): 431–434. <https://doi.org/10.15560/18.2.431>
- Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps. <http://www.simplemappr.net>
- Sselivanoff AV (1884) Materials towards the study of Russian myriapods. *Trudy russkago entomologicheskago obshchestva, St. Petersburg* 18: 69–121. [in Russian]

Stoev P & Geoffroy J (2004) An annotated catalogue of the scutigeromorph centipedes in the collection of the Muséum National d'Histoire Naturelle, Paris (France) (Chilopoda: Scutigeromorpha). *Zootaxa* 635: 1-12.

Würmli M (1975) Revision der Hundertfüßer-Gattung *Thereuonema* (Chilopoda: Scutigeridae). *Entomologica Germanica* 2 (2): 189-196.

Yang Y, Zhang L, Lin Y, Zheng Y, Jin W, Storey KB, Yu D, Zhang J (2022) The Genetic Diversity in *Thereuonema tuberculata* (Wood, 1862) (Scutigeromorpha: Scutigeridae) and the Phylogenetic Relationship of Scutigeromorpha Using the Mitochondrial Genome. *Insects* 13 (620): 1-15. <https://doi.org/10.3390/insects13070620>

Zuev RV (2016) Centipedes (Chilopoda) from the Stavropol Territory, northern Caucasus, Russia. *Arthropoda Selecta* 25 (1): 23-38. <https://doi.org/10.15298/arthsel.25.1.03>

Zuev RV & Evsyukov AP (2016) Centipedes (Chilopoda) from the Rostov-on-Don Region, southern Russia. *Russian Entomological Journal* 25 (4): 417-426. <https://doi.org/10.15298/rusentj.25.4.12>