



First record of *Dieckmanniellus nitidulus* (Gyllenhal, 1838) (Coleoptera, Brentidae) from Altaiskii Krai, Western Siberia

Andrei A. Legalov

Institute of Systematics and Ecology of Animals, SB RAS,
11 Frunze st., Novosibirsk, 630091, Russia; Altai State
University, 61 Lenina ave., Barnaul, 656049, Russia

The first record of *Dieckmanniellus nitidulus* (Gyllenhal, 1838) (Brentidae: Nanophyinae: Nanophyini) from Altaiskii Krai is given. The distribution map, illustrations and redescription of this species are presented.

doi: 10.5281/zenodo.10078434

Corresponding author: Andrei A. Legalov (fossilweevils@gmail.com)

Academic editor: R. Yakovlev | Received 8 September 2023 | Accepted 31 October 2023 | Published 9 November 2023

<http://zoobank.org/7BE1C787-C4B8-4713-AD2C-19B79F505C23>

Citation: Legalov AA (2023) First record of *Dieckmanniellus nitidulus* (Gyllenhal, 1838) (Coleoptera, Brentidae) from Altaiskii Krai, Western Siberia. *Acta Biologica Sibirica* 9: 937–942. <https://doi.org/10.5281/zenodo.10078434>

Keywords

Curculionoidea, Nanophyinae, Nanophyini, weevil, fauna, new finding, Siberia

Introduction

Representatives of the tribe Nanophyini are distributed almost throughout the world (Alonso-Zarazaga and Lyal 1999). Six species from five genera were recorded in Siberia (Zherikhin 1981; Korotyaev 1984; Legalov and Opanassenko 2000; Galich and Legalov 2012; Legalov 2020). The genus *Dieckmanniellus* Alonso-Zarazaga, 1989 includes four species distributed in the West and Central Palaearctic (Alonso-Zarazaga et al. 2017). It differs from related genera in the incompletely crenulate elytral interstria 8, strongly carinate metorostrum and very long tarsi (Alonso-Zarazaga 1989). *Dieckmanniellus nitidulus* (Gyllenhal, 1838) was recorded in Tyva (Korotyaev 1984) and Novosibirsk region (Tshernyshev and Legalov 2008).

It is the first record of this species from Altaiskii Krai.

Materials and methods

Studied specimens are kept in the ISEA – Institute of Systematics and Ecology of Animals (Russia: Novosibirsk).

Photographs, descriptions and body measuring were performed using a Zeiss Stemi 2000-C dissecting stereomicroscope.

The terminology of weevil body is according to Lawrence et al. (2010) and Alonso-Zarazaga (1989). The systematics of studied taxa are based on Alonso-Zarazaga et al. (2017) and Legalov (2018).

Result

Superfamily Curculionoidea Latreille, 1802

Family Brentidae Billberg, 1820

Subfamily Nanophyinae Gistel, 1848

Tribe Nanophyini Gistel, 1848

Genus *Dieckmanniellus* Alonso-Zarazaga, 1989

***Dieckmanniellus nitidulus* (Gyllenhal, 1838)**

Figs 1–3

Material examined. RUSSIA 1 ex. (ISEA), Altaiskii Krai, Klyuchevsky District, 4.2 km NW of Petukhi, S shore of Petukhovo lake, 52.25002° N, 79.48652° E, 17.VI.2022, A. Legalov; 1 ex. (ISEA), Novosibirsk Oblast, Karasuksky District, 20 km W of Karasuk, env. Troitskoe, 30.V.2007, S. Tshernyshev; 1 ex. (ISEA), Volgograd Oblast, east shore of Bulukhta Lake, 49.351° N, 46.116° E, 20.V.2021, R.Yu. Dudko; 2 ex. (ISEA), Volgograd Oblast, west shore of Elton Lake, 49.165° N, 46.564° E, 19-20.V.2021, R.Yu. Dudko; KAZAKHSTAN 1 ex. (ISEA), East Kazakhstan Region, 10 km SSW of Slavyanka, 8.V.1999, R. Dudko, V. Zichchenko; 5 ex. (ISEA), Jambyl Region, Karatau, 21.VII.1994, A.V. Boiko.

Description. Male. Body brownish-black, covered with narrow piliform white appressed scales. Scape, antennomeres 1–5, base of femora, tibiae, basal half of protarsomere 1, meso- and metatarsomeres 1 yellow-brown. Elytra and apex of rostrum brownish. Rostrum curved, subcylindrical, subglabrous, carinate from antennal insertion to base, slightly shorter than head and pronotum combined. Antennal scrobes distinct. Forehead very narrow. Eyes large, rounded, not protruding from contour of head, quite coarsely faceted. Vertex finely punctate. Temples short. Antennae geniculate, inserted laterally before middle of rostrum, rather long. Funicle 5-segmented. Scape long, curved in apical one third, reaching eye. Antennomere 2 suboval. Antennomere 3 long-conical, shorter and narrower than antennomere 2. Antennomere 4 conical, distinctly shorter and barely narrower than antennomere 3. Antennomeres 5 and 6 wide-conical. Antennomeres 5 shorter and wider than antennomere 4. Antennomere 6 shorter and wider than antennomere 5. Club noncompact, longer than antennomeres 2–6 combined. Pronotum almost campaniform, about 1.2 times as long as wide at apex, about 0.9 times as long as wide in middle and about 0.8 times as long as wide at base. Sides almost straight. Disk finely punctate. Greatest width at base. Scutellum very small, concealed. Elytra suboval, strongly convex, about 1.7 times as long as base width, about 1.4 times as long as wide in middle, about 2.0 times as long as wide in apical quarter, about 2.5 times as long as pronotum. Greatest width in middle. Base of elytra with granulate carina. Humeri flattened. Striae distinct. Interstriae wide, slightly convex, finely punctate. Interstria 8 incompletely crenulate. Elytral apices rounded separately. Pre- and postcoxal portions of prosternum short. Pro- and mesocoxal cavities rounded. Procoxal cavities contiguous. Mesocoxal cavities separated. Metacoxal cavities transversely extended. Andomen convex. Ventrites 1 and 2 fused, equal in length. Ventrite 1 slightly shorter than metacoxal length. Ventrites 3 and 4 equal in length, short. Ventrite 3 about 0.2 times as long as ventrite 2. Ventrite 5 about 5.0 times as long as ventrite 4.



Legs long. Procoxae conical. Meso- and metacoxae convex. Trochanters elongate. Femora slightly clavate, separated from coxae, with two teeth. Tibiae almost straight, mucronate. Tarsi long, with erect setae dorsally. Tarsomere 1 elongate-conical. Tarsomere 2 wide-conical. Tarsomere 3 bilobed. Tarsomere 5 long-conical. Claws relatively large, fused at base, without teeth. Body length: 1.7 mm.

Remarks. This species develops on *Lythrum salicaria* L. (Lythraceae) (Smreczyński 1976) and *Frankenia tuvinica* Lom. (Frankeniaceae) (Korotyaev 1984).

Distribution. North Africa, Europe, Caucasus, Asia Minor, Kazakhstan, Siberia (Alonso-Zarazaga et al. 2017). This species is found in Siberia (Novosibirsk Oblast, Altaiiskii Krai, Tyva Republic) and East Kazakhstan (Fig. 4).

Acknowledgements

The author thanks B.A. Korotyaev (Russia: St. Petersburg) for the opportunity to study of comparative material of this species and R.Yu. Dudko (Novosibirsk) who collected this species in Kazakhstan and Volgograd Oblast.



Figure 1. *Dieckmanniellus nitidulus* from Altaiskii Krai, male, dorsal view. Scale bar 0.2 mm.**Figure 2.** *Dieckmanniellus nitidulus* from Altaiskii Krai, male, lateral view. Scale bar 0.2 mm.




Figure 3. *Dieckmanniellus nitidulus* from Altaiskii Krai, male. a - aedeagus, dorsal view, b - aedeagus, lateral view, c - tegmen, dorsal view. Scale bar 0.1 mm.




Figure 4. Distribution of *Dieckmanniellus nitidulus* in Central Asia.

References

Alonso-Zarazaga MA (1989) Revision of the supraspecific taxa in the Palaearctic Apionidae Schoenherr, 1823. 1. Introduction and subfamily Nanophyinae. *Fragmenta Entomologica* 21: 205-262.

Alonso-Zarazaga MA, Barrios H, Borovec R, Bouchard P, Caldara R, Colonnelli E, Güttekin L, Hlavá P, Korotyaev B, Lyal CHC, Machado A, Meregalli M, Pierotti H, Ren L, Sánchez-Ruiz M, Sforzi A, Silfverberg H, Skuhrovec J, Trýzna M, Velázquez de Castro AJ, Yunakov NN (2017) Cooperative catalogue of Palaearctic Coleoptera Curculionoidea. *Monografías electrónicas* 8: 1-729.

Alonso-Zarazaga MA, Lyal CHC (1999) A world catalogue of families and genera *Curculio noidea* (Insecta: Coleoptera) (excepting Scolytidae and Platypodidae). *Entomopraxis*, Barcelona, 315 pp.

Galich DE, Legalov AA (2012) First record of *Nanomimus hemisphaericus* (Olivier, 1807) (Coleoptera, Brentidae, Nanophyinae) from Siberia. *Euroasian Entomological Journal* 11 (4): 354-355. [In Russian]

Korotyaev BA (1984) To the knowledge of the weevil fauna (Coleoptera, Curculionidae) of Mongolia and adjacent territories. 2. Insects of Mongolia 9: 311-355. [In Russian]

Lawrence JF, Beutel RG, Leschen RAB, Slipinsky SA (2010) Chapter 2. Glossary of Morphological



Terms. Handbook of Zoology. Arthropoda: Insecta. Tb. 40: Coleoptera (Beetles). 2: Morphology and Systematic (Elateroidea, Bostrichiformia, Cucujiformia partim), 9–20.

Legalov AA (2017) Weevils (Coleoptera, Curculionoidea) from plains of Western Siberia, Kazakhstan and Middle Asia. Part 1. Euroasian Entomological Journal 16 (3): 259–282. [In Russian]

Legalov AA (2018) Annotated key to weevils of the world. Part 1. Families Nemonychidae, Anthribidae, Belidae, Ithyceridae, Rhynchitidae, Brachyceridae and Brentidae. Ukrainian Journal of Ecology 8 (1): 780–831. https://doi.org/10.15421/2018_280

Legalov AA (2020) Revised checklist of superfamily Curculionoidea (Coleoptera) from Siberia and the Russian Far East. Acta Biologica Sibirica 6: 437–549. <https://doi.org/10.3897/abs.6.e59314>

Legalov AA, Opanassenko FI (2000) A review of the fauna of the superfamily Curculionoidea (Coleoptera) of Novosibirsk Province. Entomological Review 80 (3): 282–303.

Smreczyński S. 1976. Ryjkowce – Curculionidae. Podrodzina Curculioninae. Podrodzina Curculioninae. Plemiona: Nanophyini, Mecinini, Cionini, Anoplini, Rhynchaenini i uzupełnienia do zeszytów 98 a-e. Klucze do oznaczania owadów Polski. Warszawa, XIX (98f), 9–21.

Tshernyshev SE, Legalov AA (2008) Species composition of chortoantobiont beetles (Coleoptera: Cantharidae, Malachiidae, Dasytidae, Meloidae, Oedemeridae, Bruchidae, Anthribidae, Rhynchitidae, Brentidae, Curculionidae) from the Kulundinskaya forest-steppe of West Siberia. Euroasian Entomological Journal 7 (3): 323–333. [In Russian]

Zherikhin VV (1981) Weevils of subfamily Nanophyinae (Coleoptera, Curculionidae) from Siberia and Far East of the USSR. New data on insects of the Far East. Vladivostok, 55–62. [In Russian]