

# To the fauna of Geophilomorpha (Chilopoda) of Uzbekistan and Turkmenistan

Yurii V. Dyachkov

Altai State University, 61 Lenin Ave., Barnaul, 656049,  
Russia

Genus *Stenotaenia* C.L. Koch, 1847 is new to the fauna of the Middle Asia. The following taxa are new to the fauna of Turkmenistan: family Mecistocephalidae, genus *Krateraspis* Lignau, 1929, *K. meinerti* (Sselivanoff, 1881), and genus *Stenotaenia* C.L. Koch, 1847. Genus *Geophilus* Leach, 1814 and *G. lindbergi* (Loksa, 1971) are new to Uzbekistan. All new records are illustrated. An updated list of the chilopod species dwelling in Uzbekistan and Turkmenistan is provided.

doi: 10.5281/zenodo.10239340

Corresponding author: Yurii V. Dyachkov (dyachkov793@mail.ru)

Academic editor: R. Yakovlev | Received 1 November 2023 | Accepted 16 November 2023 | Published 3 December 2023

<http://zoobank.org/0BFFFAC1-CEFC-4870-B777-811670DC131B>

**Citation:** Dyachkov YuV (2023) To the fauna of Geophilomorpha (Chilopoda) of Uzbekistan and Turkmenistan. *Acta Biologica Sibirica* 9: 1073–1082. <https://doi.org/10.5281/zenodo.10239340>

## Keywords

Biodiversity, fauna, *Krateraspis*, Mecistocephalidae, Middle Asia, new records, Repetek, *Stenotaenia*

## Introduction

The centipede fauna of Turkmenistan and Uzbekistan is still poorly studied (Muralewicz 1926; Zalesskaja & Schileyko 1992; Krivokhatsky 1994; Dyachkov & Nedoev 2021; Dyachkov 2022b; Dyachkov 2022c; Dyachkov & Bonato 2022). This paper provides some new faunistic records from these countries.

## Materials and methods

Material is deposited in the ZISP (abbreviations below).

The standardized terminology follows Bonato et al. (2010). Locality data are given as on the original labels; additional information is provided in square brackets.

The pictures have been taken using an Olympus DP74 digital camera attached to an Olympus SZX16 stereo microscope. The distribution map (Fig. 1) was generated using SimpleMappr software (Shorthouse 2010).

Abbreviations: coll. – collector, fragm. – fragment, ZISP – Zoological Institute of the Russian Academy of Sciences, Saint Petersburg.



**Figure 1.** Distribution of *Geophilus lindbergi* (Loksa, 1971) (oval), *Stenotaenia* sp. (star), and *Krateraspis meinerti* (Sselivanoff, 1881) (square) in Middle Asian countries. Kg – Kyrgyzstan, Pa – Pakistan, Tj – Tajikistan. White color – previous records, green one – new data.

## Result

### Family Geophilidae Leach, 1816

#### Genus *Geophilus* Leach, 1814

##### *Geophilus lindbergi* (Loksa, 1971)

Figures 2–7

*Clinopodes lindbergi* Loksa 1971: 110.

*Clinopodes lindbergi* – Titova 1975: 309.

*Geophilus lindbergi* – Bonato et al. 2011: 197; Dyachkov & Nedoev 2021: 47; Dyachkov 2022a: 11.

**Material.** 1 male (ZISP chilo-64), [Uzbekistan, Bukhara Region], Bukhara [ca. N39°46', E64°26'], behind the Uglanovskie vorota [Uglon Gate], swamp, 7 April 1925, coll. M. Sokolov.

**Distribution.** Afghanistan (Loksa 1971; Dyachkov 2022a), Turkmenistan (“Kushka” [now Serhetabat, ca. N35°16', E62°20']) (Titova 1975), and Uzbekistan (new).

**Remarks.** Specimen has 53 leg-bearing segments.



This species and genus *Geophilus* are new to the fauna of Uzbekistan; present record is the northernmost range limit of the species.

**Genus Stenotaenia C.L. Koch, 1847**

**Remarks.** This genus is new to the fauna of the Middle Asia. Present record is easternmost range limit of the genus (Bonato & Minelli 2008).

**Stenotaenia sp.**

Figures 8-12

**Material.** 1 body fragm. (ZISP chilo-192), Turkmenistan, [Balkan Region], Ai-Dere [river valley, ca. N38°24', E56°44', Sunt-Hasardag Nature Reserve], 20-22 November [19]70, coll. unknown.

**Remarks.** Species identification is not possible due to absence of the rear body part.

**Family Mecistocephalidae Bollman, 1893**

**Genus Krateraspis Lignau, 1929**

**Krateraspis meinerti (Sseliwanoff, 1881)**

Figures 13-16

*Mecistocephalus meinerti* Sseliwanoff 1881a: 9; 1881b: 232; 1884: 73.

*Krateraspis meinerti* - Dyachkov 2019: 368; 2020: 79; 2023: 67; Dyachkov & Nedoev 2021: 44; Dyachkov & Bonato 2022: 149.

*Tygarrup asiaticus* Verhoeff 1930: 260.

*Tygarrup asiaticus* - Dyachkov & Bonato 2022: 149.

**Material.** 1 female (ZISP chilo-176), [Turkmenistan, Lebap Region], Repetek [N38°33'45", E63°10'38"], [date unknown], coll. Y. Bekman.

**Distribution.** Kazakhstan, Kyrgyzstan, Uzbekistan, Tajikistan (Dyachkov & Bonato 2022), and Turkmenistan (new).

**Remarks.** This species, genus *Krateraspis* Lignau, 1929, and family Mecistocephalidae are new to the fauna of Turkmenistan.



**Figure 2. Figures 2-7.** *Geophilus lindbergi* (Loksa, 1971): 2, 3 - front body fragment, ventrally and dorsally; 4 - forcipules and first segment, ventrally; 5, 6 - rear body fragment, ventrally and dorsally; 7 - 7-8th segments, ventrally. Scale: 0.1 mm (7), 0.2 mm (2-6).

## Conclusions

At present, the chilopod fauna of Uzbekistan is represented by at least 28 species in 16 genera, 8 families, and 4 orders; while the fauna of Turkmenistan consists of at least 16 species in 11 genera, 7 families, and 4 orders.

It should be recognized that our knowledge of Middle Asian Geophilomorpha is still far from being complete.

**List of the Chilopoda species occurring in Uzbekistan****Order Geophilomorpha****Family Geophilidae**

1. *Geophilus lindbergi* (Loksa, 1971)
2. *Pachymerium ferrugineum* (C.L. Koch, 1835)
3. *Taschkentia parthorum* (Pocock, 1891)
4. *T. bucharensis* Verhoeff, 1930

**Family Himantariidae**

5. *Bothriogaster signata* (Kessler, 1874)
6. *Polyporogaster porosa* (Sselivanoff, 1881)
7. *P. turkestanica* Verhoeff, 1930

**Family Mecistocephalidae**

8. *Arrup asiaticus* (Titova, 1975)
9. *Krateraspis meinerti* (Sselivanoff, 1881)
10. *K. sselivanovi* Titova, 1975

**Order Lithobiomorpha****Family Henicopidae**

11. *Cermatobius kirgisicus* (Zalesskaja, 1972)

**Family Lithobiidae**

12. *Australobius magnus* (Trotzina, 1894)
13. *Bothropolys desertorum* Lignau, 1929
14. *B. ghilarovi* Zalesskaja, 1975
15. *B. lutulentus* Verhoeff, 1930
16. *Hessebius plumatus* Zalesskaja, 1978
17. *Lithobius crassipes* L. Koch, 1862
18. *L. javanicus* (Zalesskaja, 1978)
19. *L. krali* (Dobroruka, 1979)
20. *L. praeditus* Zalesskaja, 1975

21. *L. turkestanicus* Attems, 1904

**Order Scutigeromorpha**

**Family Scutigeridae**

22. *Scutigera asiatica* Sseliwanoff, 1884


23. *Thereuonema turkestana* Verhoeff, 1905

**Order Scolopendromorpha**


**Family Cryptopidae**

24. *Cryptops doriae* Pocock, 1891

25. *C. hortensis* (Donovan, 1810)



**Figure 3. Figures 8-12.** *Stenotaenia* sp.: **8, 9** - front body fragment, ventrally and dorsally; **10** - forcipules, ventrally; **11** - clypeus and labrum, ventrally; **12** - 6-7th segments, ventrally. Scale: 0.1 mm (**10-12**), 0.2 mm (**8, 9**).



**Figure 4. Figures 13-16.** *Krateraspis meinerti* (Sseliwanoff, 1881): 13, 14 – front body fragment, ventrally and dorsally; 15, 16 – rear body fragment, ventrally and dorsally. Scale bar: 0.2 mm.

### Family Scolopendridae

26. *Scolopendra canidens* Newport, 1844

27. *S. cingulata* Latreille, 1829

28. *S. mirabilis* (Porat, 1876)

### List of the Chilopoda species occurring in Turkmenistan

#### Order Geophilomorpha

##### Family Geophilidae

1. *Geophilus lindbergi* (Loksa, 1971)

2. *Pachymerium ferrugineum* (C.L. Koch, 1835)

3. *Stenotaenia* sp.

##### Family Himantariidae

4. *Bothriogaster signata* (Kessler, 1874)

5. *Polyporogaster geminata* (Silvestri, 1895)

6. *P. porosa* (Sseliwanoff, 1881)

##### Family Mecistocephalidae

7. *Krateraspis meinerti* (Sseliwanoff, 1881)

**Order Lithobiomorpha****Family Lithobiidae**

8. *Hessebius barbipes* (Porat, 1893)

9. *L. icis* Zalesskaja, 1978

10. *L. juniperius* Zalesskaja, 1978

11. *L. vinciguerrae* Silvestri, 1895

**Order Scutigeromorpha****Family Scutigeridae**

12. *Scutigera coleoptrata* (Linnaeus, 1758)

**Order Scolopendromorpha****Family Cryptopidae**

13. *Cryptops hortensis* (Donovan, 1810)

14. *C. caucasius* Verhoeff, 1934

**Family Scolopendridae**

15. *Scolopendra canidens* Newport, 1844

16. *S. mirabilis* (Porat, 1876)

## Acknowledgements

I am grateful to Viktor A. Krivokhatsky (ZISP) who loaned the material. I also express my gratitude to Mariia A. Iuzhakova (Tomsk, Russia) who kindly checked the English of the draft. The reported study was funded by state assignment of the Ministry of Science and Higher Education of the Russian Federation (project FZMW- 2023-0006 "Endemic, local and invasive arthropods (Arthropoda) of the mountains of South Siberia and Central Asia: a unique gene pool of a biodiversity hotspot").

## References

Bonato L, Edgecombe GD, Lewis JG, Minelli A, Pereira LA, Shelley RM & Zapparoli M (2010) A common terminology for the external anatomy of centipedes (Chilopoda). ZooKeys 69: 17–51. <https://doi.org/10.3897/zookeys.69.737>

Bonato L, Iorio E, Minelli A (2011) The centipede genus *Clinopodes* C. L. Koch, 1847 (Chilopoda, Geophilomorpha, Geophilidae): reassessment of species diversity and distribution, with a new species from the Maritime Alps (France). Zoosystema 33: 175–205.

Bonato L, Minelli A (2008) *Stenotaenia* Koch, 1847: a hitherto unrecognized lineage of western Palaearctic centipedes with unusual diversity in body size and segment number (Chilopoda: Geophilidae). Zoological Journal of the Linnean Society 153: 253–286.

Dyachkov YuV (2019) New data on the family Mecistocephalidae Bollman, 1893 (Chilopoda: Geophilomorpha) from Middle Asia. *Arthropoda Selecta* 28 (3): 368–373.  
<https://doi.org/10.15298/arthsel.28.3.02>

Dyachkov YuV (2020) New data on the centipede (Chilopoda) fauna from Tajikistan. *Ecologica Montenegrina* 36: 78–86. <http://dx.doi.org/10.37828/em.2020.36.6>

Dyachkov YuV (2022a) An annotated checklist of Chilopoda from Afghanistan. *Ecologica Montenegrina* 53: 8–24. <https://dx.doi.org/10.37828/em.2022.53.2>

Dyachkov YuV (2022b) New records of lithobiid centipedes (Chilopoda: Lithobiomorpha) from Middle Asia. *Acta Biologica Sibirica* 8: 399–407. <https://doi.org/10.5281/zenodo.7703408>

Dyachkov YuV (2022c) On new records of Geophilomorpha (Chilopoda) from Middle Asia. *Ecologica Montenegrina* 60: 70–79. <https://dx.doi.org/10.37828/em.2022.60.11>

Dyachkov YuV (2023) On the Chilopoda fauna of the Pamir Mts, Tajikistan, with notes on the highest records of chilopod species in Middle Asia. *Ecologica Montenegrina* 65: 67–75.  
<https://dx.doi.org/10.37828/em.2023.65.9>

Dyachkov YuV, Bonato L (2022) Morphology and distribution of the Middle Asian centipede genus *Krateraspis* Lignau, 1929 (Chilopoda, Geophilomorpha, Mecistocephalidae). *ZooKeys* 1095: 143–164. <https://doi.org/10.3897/zookeys.1095.80806>

Dyachkov YuV, Nedoev KhKh (2021) A contribution to the centipede (Chilopoda: Geophilomorpha, Scolopendromorpha) fauna of Uzbekistan and Turkmenistan. *Ecologica Montenegrina* 41: 41–50.  
<http://dx.doi.org/10.37828/em.2021.41.6>

Krivokhatsky VA (1994) Arthropods Inhabiting Rodent Burrows in the Karakum Desert. In: Fet V, Atamuradov KI (Eds) Biogeography and Ecology of Turkmenistan. Kluwer Academic Publ., Netherlands, 389–402. [In Russian]

Loksa I (1971) Die von K. Lindberg in Afghanistan gesammelten Chilopoden. *Senckenbergiana biologica* 52 (1/6): 103–112.

Muralewicz WS (1926) Übersicht über die Chilopodenfauna des Kaukasus. II. *Zoologischer Anzeiger* 69 (1/2): 27–44.

Shorthouse DP (2010) SimpleMappr, an online tool to produce publication-quality point maps.  
<http://www.simplemappr.net>

Sseliwanoff AV (1881a) Geophilidae from the Museum of Imperial Academy of Sciences. *Zapiski Imperatorskoi Akademii Nauk* 40: 1–27. [In Russian]

Sseliwanoff AV (1881b) Turkestanskiya stonozhki (Geophilidae Leach). *Izvestiya Imperatorskogo Obshchestva Lyubitelei Estestvoznanija, Antropologii i Etnografii pri Imperatorskom Moskovskom Universitete* 37: 229–232. [In Russian]

Sseliwanoff AV (1884) Materials towards the study of Russian myriapods. *Trudy Russkago Entomologicheskago Obshchestva* 18 (1–2): 69–121. [In Russian]

Titova LP (1975) The centipede genera *Clinopodes* and *Pleurogeophilus* in the USSR fauna (Chilopoda, Geophilomorpha, Geophilidae). *Problemy pochvennoj zoologii. Materialy V Vsesoyuznogo soveshchaniya*. Vilnius: 308–309. [In Russian]



Zalesskaja NT, Schileyko AA (1992) The distribution of Scolopendromorpha in the USSR (Chilopoda). Advances in Myriapodology. Berichte des Naturwissenschaftlich-Medizinischen Vereins in Innsbruck, Supplement 10: 367-372.

Verhoeff KW (1930) Über Myriapoden aus Turkestan. Zoologischer Anzeiger 91 (9/12): 243-266.