

Coenotic and biomorphological analysis of Lamiaceae in the West Siberian floristic province

Vera A. Cheryomushkina

Central Siberian Botanical Garden SB RAS, 101
Zolotodolinskaya St., Novosibirsk, 630090, Russia

Evgenia B. Talovskaya

Central Siberian Botanical Garden SB RAS, 101
Zolotodolinskaya St., Novosibirsk, 630090, Russia

Anastasia E. Sharabarina

Central Siberian Botanical Garden SB RAS, 101
Zolotodolinskaya St., Novosibirsk, 630090, Russia

We analyzed the distribution, coenotic habitat, ecological preferences, and biomorphology of the 80 Lamiaceae species from 30 genera found in the West Siberian province. The plant species are primarily found in boreal and forest-steppe vegetation communities and are equally distributed in both lighted and shaded areas. Most species have a narrow range of humidity preferences. The study also highlights the diversity of life forms, with long-rhizomatous grasses being the most common, and morphological traits such as monocyclic elongated shoots, orthotropic growth direction, terminal arrangement of inflorescences, sympodial development, appearance of replacement axes due to basiton branching, and a long development period of more than three years. The research reveals that the diversity of life forms and biomorphological features of the shoot system play a significant role in determining the distribution of species in different ecological and coenotic conditions within the West Siberian province.

Acta Biologica Sibirica 9: 1207–1225 (2023) doi: 10.5281/zenodo.10448401

Academic editor: A. Matsyura | Received 4 December 2023 | Accepted 20 December 2023 |
Published 31 December 2023

<http://zoobank.org/DCD68B3B-2D98-4004-8830-4E050D57A238>

Citation: Cheryomushkina VA, Talovskaya EB, Sharabarina AE (2023) Coenotic and biomorphological analysis of Lamiaceae in the West Siberian floristic province. Acta Biologica Sibirica 9: 1207–1225. <https://doi.org/10.5281/zenodo.10448401>

Keywords

West Siberian floristic province, Lamiaceae, ecological and coenotic characterization, biomorphological traits

Introduction

The Lamiaceae Martinov family is a significant component of the global flora, consisting of approximately 236 genera and over 7000 species (Harley et al. 2004; APG IV 2016). These plants are found on all continents, from tropical to Arctic regions, and hold key ecological and economic importance. They thrive in diverse ecological niches, ranging from coastal water areas of plain rivers to cryophytic plant communities of rocky slopes in subalpine and alpine mountain belts. In Eurasia, there are over 500 Lamiaceae taxa organized into systematic groups of varying sizes and

ranks. Researchers from Asian and European countries have shown particular interest in these plants due to their theoretical and practical significance (Jamzad 2013; Lazkov 2016; Massoud, Chonour 2013; 2018; Safhhani et al. 2018; Zhao et al. 2020; 2021) and Europe (Paton 1990; Hedge 1992; Mathiesen et al. 2011; Carovic-Stanko et al. 2016; Salimov et al. 2018). On the territory of Russia and neighboring countries, Lamiaceae species have been studied for the most part only in terms of taxonomic diversity (Klokov 1954; Shishkin 1954; Shishkin Yuzepchuk 1954; Adylov 1987; Budantsev 1990; Kamelin, Makhmedov 1990) and chemical composition of their natural compounds (Budantsev 2011; Mamadaliyeva et al. 2017). Often individual taxa are used by researchers as model objects when studying the mechanisms of plant adaptation to different habitat conditions (Astashenkov 2015; Talovskaya (Kolegova) 2015; Cheryomushkina et al. 2020, 2022; Talovskaya, Cheryomushkina 2022). The study of the distribution of large taxa is of particular importance, as these data will further make it possible to assess the current distribution of species of the family in a certain area, predict the response of vegetation to changes in habitat conditions, and characterize the peculiarities of the morphological structure of species in the spectrum of habitats (Pyšek, Liška 1991; Körner 1992; Anthelme et al. 2014; Talovskaya, Cheryomushkina 2022).

In Russia and neighboring countries, Lamiaceae species have primarily been studied for taxonomic diversity and the chemical composition of their natural compounds. Some taxa are also utilized as model organisms for studying plant adaptation mechanisms to different habitats. Understanding the distribution of these taxa is crucial for assessing their current presence in specific areas, predicting vegetation responses to habitat changes, and characterizing species' morphological structures across different habitats. Siberia, the largest region in North Asia, offers a wide range of ecological conditions and is home to 148 Lamiaceae species from 30 genera (Doronkin 2012). These species exhibit various life forms, including perennial and annual herbs, semi-shrubs, and shrubs.

The goal of this study was to investigate the distribution, coenotic habitat, ecological preferences, and biomorphology of Lamiaceae species in the West Siberian floristic province.

Materials and methods

The study was carried out in Western Siberia within the West Siberian province, encompassing Tobolsk, Kurgan, Omsk, Tomsk, Novosibirsk, Kemerovo, and Barnaul floristic regions (Takhtajyan 1978; Malyshev et al. 2000). Floristic zoning was performed with contrast zoning (Malyshev et al. 2000) by flora species composition (Fig. 1). The province is situated in the West Siberian Plain and features a predominantly uniform landscape, with occasional minor elevations and depressions (0-300 m above sea level). The distinctiveness of the province's vegetation cover lies in its extensive swampy areas. The taiga and forest-steppe zones dominate the province, while the steppe zone is only sporadically present.



Figure 1. Map of the West Siberian floristic province according to L.I. Malyshev et al. (2000). Bold dashed line – border of

Western Siberia, blue line - border of the floristic province, dotted line - borders of the province districts.

Species distribution

The material for the analysis consisted of herbarium collections gathered independently during field studies in Western Siberia, as well as from the NS (CSBS SB RAS, Novosibirsk) and ALTB (Altai State University, Barnaul) herbarium collections. Floristic works by L.I. Malyshev (1997), V.M. Doronkin (2012), and M.M. Silantieva (2013), along with data from the Global Biodiversity Information Facility (GBIF) at <https://www.gbif.org/ru/> and Plants of the World Online (POWO) at <https://powo.science.kew.org/>, were utilized to establish the general distribution pattern of species. Relying on R.V. Kamelin's approach (2017), the species of the Lamiaceae family were classified into 8 chorological (geographical) groups: multiregional, Holarctic, Palaeartic, Volga-South Ural-Southwest Siberian, South Siberian-Central Asian, South Siberian, Altai-Central Asian, and endemic (subendemic).

The phytocoenotic assignment of Lamiaceae species was determined based on A.V. Kuminova (1960), L.I. Malyshev (1997), N.B. Ermakov (2003), D.M. Doronkin (2005), N.I. Makunina (2016), and data from herbarium collections. The species names were referenced from the international database World Checklist of Vascular Plants at <https://powo.science.kew.org/>. Eight coenotic complexes were identified, namely: sandy shores and dunes, shrub thickets, mountain-tundra, anthropophytic, petrophytic, meadow, steppe, and forest.

Ecological traits

To determine the ecological preferences of Lamiaceae species, we conducted an analysis of their tolerance to moisture, light, and substrate type. Regarding moisture, we identified eight groups: mesoxerophytes, euxerophytes, xeromesophytes, eumesophytes, mesohygrophytes, hygrophytes, hydrophytes, and psychrophytes. In terms of illumination, we distinguished three groups: heliophytes, sciophytes, sciogeliophytes, or facultative heliophytes. Additionally, we categorized all Lamiaceae species into three groups based on substrate type: psammophytes, petrophytes, and non-petrophytes. We then used Ward's method, as outlined by Siegel and Castellan in 1989, to associate Lamiaceae species distributed in the West Siberian province according to their ecological and coenotic preferences. This analysis was conducted using the Statistics 10 software package.

Morphological traits

Biomorphological characters were analyzed on at least five herbarium specimens of each species. A total of 400 Lamiaceae specimens were analyzed. Eight traits were selected as initial data: stem elongation cycle (mono-, di- and polycyclic), stem structure (semi-rosette, rosette, and elongated), auxiliary shoot position/branching (acrotonic, basitonic, mesotonic, monopodial or sympodial), inflorescence position (terminal or lateral), flowering types (polycarpic, monocarpic), life cycle (annual, biennial, and perennial), axes differentiation (orthotropic, plagiotropic, and anisotropic) (Serebryakov 1959; Barthélémy, Caraglio 2007). When characterizing branching, the positioning of the second-order shoots along the parent axis was determined (acrotonic, basitonic, and mesotonic).

Life form and biormorph types

In this study, we examine two approaches to classifying life forms. The first is the C. Raunkiaer (1934) classification, which is based on the position of the renewal buds. The second approach is the ecological-morphological classification of E. Warming (1909) and I.G. Serebryakov (1962, 1964), which analyzes a set of morphological features of plants based on specific habitat conditions.

Result

Taxonomic composition

The study showed that 80 species of Lamiaceae belonging to 30 genera grow on the territory of the West Siberian province. The leading among them are *Thymus* L. (10 species), *Dracocephalum* L. (9 species), *Scutellaria* L. (6 species) and *Mentha* L. (6 species) (Table 1). The character of distribution of genera and species over the territory of the province shows that their greatest number is concentrated in the Barnaul district (95% of the total number of genera and 89% of the total number of species). A nucleus consisting of 8 genera is distinguished, which are found in all districts of the province: *Dracocephalum*, *Galeopsis* L., *Leonurus* L., *Mentha*, *Origanum* L., *Prunella* L., *Scutellaria*, and *Stachys* L.

Distribution and ecology of Lamiaceae

As a result of chorological analysis, all studied species of Lamiaceae were divided into 8 chorological groups (Table 2). The largest number of species (43 species) constitutes the Palaearctic realm. Among them, most species have Palaearctic origin (18 species, like *Lamium album* L., *Scutellaria supina* L., and *Thymus marschallianus* Willd.) and West Palaearctic (17 species, among them *Betonica officinalis* L., *Chaiturus marrubiastrum* (L.) Ehrh. ex Rchb., *Dracocephalum thymiflorum* L.) origin. The Altai-Central Asian group is represented by 14 species, among them *Dracocephalum integrifolium* Bunge, *Lagopsis marrubiastrum* (Stephan) Ikonn.-Gal., and *Phlomoides agraria* (Bunge) Adylov, Kamelin & Makhm. as well as species not entering Mongolian territories, like *Hyssopus ambiguus* (Trautv.) Iljin ex Prochorov. & Ledel, *Thymus petraeus* Serg., *T. proximus* Serg., and *T. roseus* Schipcz. The same number of species form South Siberian-Central Asian (7 species, among them *Dracocephalum discolor* Bunge, *D. grandiflorum* L., and *Leonurus deminutus* V.I.Krecz.), Holarctic (6 species, like *Glechoma hederacea* L., *Lamium purpureum* L., and *Lycopus europaeus* L.) and pluriregional or cosmopolitan groups (5 species, among them *Clinopodium acinos* (L.) Kuntze, *Leonurus japonicus* Houtt., and *Mentha aquatica* L.). For some species the cosmopolitan range is secondary. For example, *Clinopodium acinos* is a weed in neglected fields; *Mentha aquatica*, *M. arvensis* L., and *M. longifolia* (L.) L. are the species widely used by humans in culture as medicinal and aromatic plants; *Leonurus japonicus* - formerly cultivated as a medicinal plant, but now occurring as a weed. There are three endemic and subendemic species of Lamiaceae in the study area: *Scutellaria altaica* Ledeb. ex Sweet, *Scutellaria mongolica* Sobolevsk. and *Thymus elegans* Serg. *Thymus jenisseensis* Iljin belongs to South Siberian areal and *Thymus punctulosus* Klokov belongs to Trans-Volga-South Ural-South-West Siberian range.

Genera	Species number	Species number per administrative district						
		1	2	3	4	5	6	7
<i>Acinos</i>	1	0	1	0	1	1	0	1
<i>Amethystea</i>	1	0	0	0	0	1	0	1
<i>Antonina</i>	1	0	0	0	0	0	0	1
<i>Betonica</i>	1	1	0	0	1	0	1	1
<i>Chaiturus</i>	1	0	0	1	0	1	0	1
<i>Clinopodium</i>	1	1	0	0	0	0	0	0
<i>Dracocephalum</i>	9	2	3	3	3	3	6	8
<i>Elsholtzia</i>	1	0	0	0	1	1	1	1
<i>Galeopsis</i>	3	3	3	3	3	3	3	3
<i>Glechoma</i>	1	1	0	0	1	1	1	1
<i>Hyssopus</i>	1	0	0	0	0	0	1	1
<i>Lagopsis</i>	1	0	0	0	0	0	0	1
<i>Lamium</i>	3	3	2	1	3	3	3	3
<i>Leonurus</i>	5	2	2	2	3	3	4	4

<i>Lycopus</i>	2	1	2	2	2	2	2	2
<i>Melissa</i>	1	0	0	0	0	0	0	1
<i>Mentha</i>	6	2	1	1	1	2	2	6
<i>Nepeta</i>	4	0	2	2	3	4	1	4
<i>Origanum</i>	1	1	1	1	1	1	1	1
<i>Panzerina</i>	1	0	0	0	0	0	0	1
<i>Phlomis</i>	4	1	1	1	1	2	1	4
<i>Prunella</i>	1	1	1	1	1	1	1	1
<i>Salvia</i>	4	0	2	1	0	2	2	4
<i>Satureja</i>	1	0	0	0	0	0	0	1
<i>Schizonepeta</i>	2	1	0	1	0	1	1	1
<i>Scutellaria</i>	6	2	1	2	3	4	3	4
<i>Stachys</i>	4	2	2	2	2	3	3	4
<i>Teucrium</i>	1	0	0	0	0	0	0	1
<i>Thymus</i>	10	0	2	3	3	1	6	8
<i>Ziziphora</i>	1	0	0	0	0	1	1	1
Total	80	24	26	27	33	41	44	71

Table 1. Distribution of Lamiaceae in the West Siberian Province

Notes: 1 - Tobolsk, 2 - Kurgan, 3 - Omsk, 4 - Tomsk, 5 - Novosibirsk, 6 - Kemerovo, 7 - Barnaul.

	Areal	Species number	Species number (% from total)
1	Multiregional	5	6.3
2	Holarctic	6	7.5
3.1	Palaeartic realm	18	22.5
3.2	Eastern Palaeartic	5	6.3
3.3	Western palaeartic	17	21.3
3.4	Central palaeartic	3	3.8
4	Trans-Volga-South Ural-Western Siberia	1	1.3
5	South Siberian- Central Asian	7	8.8
6	South-Siberian	1	1.3
7	Altai-Central Asian	14	17.5
8.1	Altai-Sayan-Tuva	2	2.5
8.2	Altai	1	1.3

Table 2. Chorological groups of Lamiaceae

The analysis of ecological tolerance has shown that according to water content adaptation, most species belong to mesophytes (*Amethystea caerulea* L., *Dracocephalum nutans* L., and *Glechoma hederacea* L.), slightly less are mesoxerophytes (*Nepeta nuda* L., *Phlomis tuberosa* (L.) Moench, and *Salvia verticillata* L.) and euxerophytes (*Hyssopus ambiguus* and *Ziziphora clinopodioides* Lam. – see Fig. 2. According to light adaptation, most of the studied species belong to sciogeliophytes (*Clinopodium vulgare* L., *Origanum vulgare* L., and *Satureja hortensis* L.). In relation to the substrate, the absolute majority of species are non-petrophytic. The obligate petrophytes include *Thymus altaicus* Klokov & Des.-Shost. and *T. punctulosus*, while the psammophytes include *Satureja hortensis*.




Figure 2. Number of *Lamiaceae* species (%) in different ecological groups (a - humidity, b - light, c - nutrition).

Lamiaceae in the West Siberian province plant communities

The greatest number of *Lamiaceae* species is found in communities of four coenotic complexes: petrophytic (61 species), meadow (56), anthropophytic (52 species), and steppe (35 species, see Table 3). Only two *Lamiaceae* species were recorded in the communities of the mountain-tundra complex. The petrophytic complex unites *Lamiaceae* growing in rock crevices (*Thymus elegans* and *T. petraeus*), on stonyrubble slopes (*Lagopsis marrubiastrum*, *Hyssopus ambiguus*, and *Ziziphora clinopodioides*) and scree slopes (*Thymus mongolicus* (Ronniger) Ronniger and *T. roseus*). The meadow complex is one of the most diverse in composition of coenotic groups. Among them, the steppe-meadow group is the richest in the number of species – 26 species (*Chaiturus marrubiastrum*, *Lamium purpureum* and *Leonurus tataricus* L.); in the forest-meadow group there are 16 species (*Dracocephalum thymiflorum*, *Galeopsis speciosa* Mill., and *Glechoma hederacea*); in the floodplain-meadow group – 11 species (*Scutellaria hastifolia* L., *Stachys palustris* L., and *Teucrium scordioides* Schreb.); in mountain-forest and subalpine – 3 species (*Phlomoidea alpina* (Pall.) Adylov, Kamelin & Makhm., *P. oreophila* (Kar. & Kir.) Adylov, Kamelin & Makhm.); in alpine-meadow group only *Dracocephalum grandiflorum*. In the steppe complex, the steppe coenotic group proper is the richest in the number of *Lamiaceae* species, which unites 24 species (*Amethystea caerulea*, *Leonurus glaucescens* Bunge, and *Nepeta ucranica* L.). Meadow-steppe coenotic group includes 9 species (*Nepeta nuda* and *Schizonepeta multifida* (L.) Briq.), shrub-steppe – 2 species (*Salvia deserta* Schangin and *Scutellaria supina*). Anthropophytic complex combines ruderal and segetal species, like *Elsholtzia ciliata* (Thunb.) Hyl., *Mentha spicata* L., and *Stachys byzantina* K. Koch.

Most of the forest complex species are associated with light coniferous (*Clinopodium acinos*, *Dracocephalum nutans*, and *Origanum vulgare*) and small-leaved (*Dracocephalum ruyschiana* L., *Phlomoidea tuberosa*, and *Prunella vulgaris* L.) forests. In dark coniferous forests there is one *Lamiaceae* species *Stachys sylvatica* L. In addition, the flora of alpine tundra is not rich in *Lamiaceae* species, in this habitat species of the genus *Dracocephalum* can occur: *D. grandiflorum* and *D. imberbe* Bunge.

Coenotic complex	Coenotic group	Species number
Forest	1.1 Light coniferous	8
	1.2 Dark coniferous	2
	1.3 Small-leaved	7
Steppe	2.1 Steppe-meadow	9
	2.2 Shrub-steppe	2
	2.3 Steppe	24
Meadow	3.1 Wooded meadow	16
	3.2 Steppe meadow	25
	3.3 Floodplain meadow	11
	3.4 Alpine-forest and subalpine	3
	3.5 Grassland meadows	1

Alpine tundra	Alpine tundra	2
Petrophytic	5.1 Rocks	14
	5.2 Rocky slopes	26
	5.3 Rubble slopes	13
	5.4 Taluses	8
Anthropophytic	6.1 Ruderal	31
	6.2 Segetal	21
Shrub thickets		10
	Sandy riverbanks and dunes	14

Table 3. Phytocoenotic assignment of species of the Lamiaceae family

Shrub thickets with 10 species of Lamiaceae (*Clinopodium delibe* (Bunge) Kuntze, *Galeopsis speciosa*, *Lycopus europaeus*, and *Mentha longifolia*) and sandy river banks and dunes with 14 species (*Leonurus japonicus*, *Lycopus exaltatus* L.f., and *Scutellaria scordiifolia* Fisch. ex Schrank) were identified as special coenotic complexes.

A significant part of species (74 species) of the family participates in several coenotic complexes. For example, *Amethystea caerulea* is found in steppe, petrophytic and anthropophytic complexes; *Dracocephalum ruyschiana* – in forest, meadow and petrophytic complexes; *Origanum vulgare* – in forest, steppe, meadow and petrophytic complexes. Single species are confined to one coenotic complex. *Dracocephalum integrifolium* is found only in communities of the petrophytic complex; *Elsholtzia ciliata* and *Galeopsis bifida* Boenn. – in communities of the anthropophytic complex; *Phlomoides agraria* – in communities of the steppe complex.

Diversity of morphological characters in Lamiaceae

Most plants of the Lamiaceae growing in the West Siberian Province are characterized by monocyclic (55.7%) elongated (57.0%) shoots, orthotropic growth direction (83.5%), terminal inflorescence arrangement (82.3%), sympodial development (73.4%), appearance of substitute axes as a result of basiton branching (52.6%) and long development period (from three years) (80.0%, see Table 4). Plants with polycyclic shoots are less common. Species with at least one of the following traits are found only rarely: rosette shoots, plagiotropic growth direction, acrotonic branching, and biannual growth duration.

Stem elongation cycle			
Monocyclic (55.7%)	Dicyclic (29.1%)		Polycyclic (15.2%)
<i>Amethystea caerulea</i> , <i>Antonina debilis</i> , <i>Galeopsis bifida</i>	<i>Lagopsis marrubastrum</i> , <i>Leonurus japonicus</i> , <i>Panzerina lanata</i> (L.) Soják		<i>Dracocephalum imberbe</i> , <i>Phlomoides oreophila</i> , <i>Thymus jensseensis</i>
Shoot structure			
Elongated (57.0%)	Rosette (5.1%)	Semi-rosette (25.3%)	Mixed (12.7%)
<i>Clinopodium acinos</i> , <i>Clinopodium vulgare</i> , <i>Ziziphora clinopodioides</i>	<i>Dracocephalum grandiflorum</i> , <i>D. imberbe</i> , <i>Phlomoides alpina</i>	<i>Betonica officinalis</i> L., <i>Chaiturus marrubiastrum</i> , <i>Leonurus deminutus</i>	<i>Thymus</i>
Axes differentiation			
Orthotropic (83.5%)	Anisotropic (15.1%)		Plagiotropic (1.3%)
<i>Elsholtzia ciliata</i> , <i>Hyssopus ambiguus</i> , <i>Lamium album</i>	<i>Clinopodium vulgare</i> , <i>Dracocephalum discolor</i> , <i>Thymus</i>		<i>Glechoma hederacea</i>
Inflorescence position			
Terminal (82.3%)		Lateral (17.7%)	
<i>Lycopus exaltatus</i> , <i>Melissa officinalis</i> L., <i>Nepeta cataria</i> L.		<i>Dracocephalum grandiflorum</i> , <i>D. imberbe</i> , <i>Phlomoides oreophila</i> , <i>Thymus</i>	
Flowering type			
Monocarpic (21.5%)		Polycarpic (78.5%)	

<i>Amethystea caerulea</i> , <i>Chaiturus arrubiastrum</i> , <i>Galeopsis ladanum</i> L.		<i>Hyssopus ambiguus</i> , <i>Lamium album</i> , <i>Lycopus exaltatus</i>	
Auxiliary shoot position			
Monopodial (26.6%)		Sympodial (73.4%)	
<i>Galeopsis bifida</i> , <i>Amethystea caerulea</i> , <i>Chaiturus marrubiastrum</i>		<i>Nepeta sibirica</i> L., <i>Lagopsis marrubiastrum</i> , <i>Glechoma hederacea</i>	
Positioning of the second-order shoots along the parent axis			
Acrotonic (10.1%)		Mesotonic (37.3%)	Basitonic (52.6%)
<i>Thymus altaicus</i> , <i>T. proximus</i> , <i>T. petraeus</i>		<i>Scutellaria supina</i> , <i>S. hastifolia</i> , <i>Stachys palustris</i>	<i>Phlomoïdes tuberosa</i> , <i>Thymus marschallianus</i> , <i>T. roseus</i>
Life cycle			
Annual (15.0%)	Biennial (5.0%)		Perennial (80.0%)
<i>Galeopsis ladanum</i> L., <i>Dracocephalum thymiflorum</i> , <i>Elsholtzia ciliata</i>	<i>Leonurus deminutus</i> , <i>L. japonicus</i> , <i>Schizonepeta annua</i> (Pall.) Schischk.		<i>Betonica officinalis</i> , <i>Clinopodium vulgare</i> , <i>Mentha aquatica</i>

Table 4. Diversity of morphological characters in Lamiaceae

The analysis of biomorphological characters showed that plants of 21 Lamiaceae genera distributed in the territory of the West Siberian province have no variability of morphological characters. In the species of the remaining 9 genera, which constitute 60% of the total number of all the Lamiaceae species in the study area, the polyvariance of at least one of the biomorphological characters was revealed (Table 5). Such genera include *Dracocephalum*, *Lamium* L., *Leonurus*, *Phlomoïdes* Moench, *Salvia* L., *Schizonepeta* (Benth.) Briq., *Scutellaria*, *Stachys* and *Thymus*. The characters, diversity of which is noted in representatives of these genera, are shoot development cycle and type of branching. Polyvariance of shoot structure, duration of development of an individual, frequency of flowering and development is also often noted.

Genera	Morphological traits							
	1	2	3	4	5	6	7	8
<i>Dracocephalum</i>	+	+	+	+	+	+	+	+
<i>Lamium</i>	+	+	+				+	+
<i>Leonurus</i>	+	+	+	+			+	+
<i>Phlomoïdes</i>			+	+		+	+	+
<i>Salvia</i>			+	+				
<i>Schizonepeta</i>	+	+					+	+
<i>Scutellaria</i>				+				
<i>Stachys</i>	+	+	+	+				+
<i>Thymus</i>				+				+

Table 5. Matrix of morphological traits variability

Note: 1 - Life cycle, 2 - Flowering type, 3 - Shoot structure, 4 - Shoot elongation structure, 5 - Axes differentiation, 6 - inflorescence position, 7 - auxiliary shoot position, 8 - positioning of the second-order shoots, "+" - polyvariance of the trait.

Life forms and phenorhythmotypes

The analysis of Lamiaceae species using Raunkiaer's (1934) approach revealed that hemicryptophytes (50.6%) are the most common life form among species such as *Betonica*

officinalis, *Dracocephalum peregrinum* L., and *Mentha aquatica*. Therophytes account for 22.8% (including *Chaiturus marrubiastrum*, *Dracocephalum moldavica* L., and *Satureja hortensis*), while chamaephytes make up 20.3% (including *Dracocephalum integrifolium* and *Hyssopus ambiguus*). Additionally, 6.3% of the species (such as *Lycopus europaeus*, *Stachys palustris*, and *Teucrium scordioides*) are cryptophytes.

Furthermore, applying the ecological-morphological approach of C. Warming (1909) and I.G. Serebryakov (1962) to Lamiaceae species in the West Siberian province, it was found that 82.3% are herbs, 11.4% are dwarf semi-shrubs, and 6.3% are dwarf shrubs (see Fig. 3). Herbaceous plants are further categorized based on their lifespan, with perennial and one- or two-year herbaceous plants identified. Perennial grasses exhibit a high diversity in their underground organs, with five types identified: Taproot caudex, short-rhizomatous, long-rhizomatous, ground creeping, and subterranean-stoloniferous (refer to Table 6). Notably, long-rhizomatous perennial grasses (21 species) are the most prevalent among the diverse range of grasses.

Life form	% from total species	Species
Taproot ephemeral grass	16.5	<i>Clinopodium delibe</i> , <i>Elsholtzia ciliata</i> , <i>Lamium purpureum</i>
Taproot biannual grass	6.3	<i>Chaiturus marrubiastrum</i> , <i>Leonurus deminutus</i> , <i>Schizonepeta annua</i>
Perennial grasses		
Taproot caudex	15.2	<i>Panzerina lanata</i> (L.) Sojak., <i>Phlomis alpina</i> , <i>Salvia stepposa</i> Des.-Shost.
Short-rhizomatous	11.4	<i>Dracocephalum nutans</i> , <i>Nepeta ucranica</i> , <i>Prunella vulgaris</i>
Long-rhizomatous	26.6	<i>Clinopodium vulgare</i> , <i>Mentha arvensis</i> , <i>Nepeta sibirica</i>
Creeping	1.3	<i>Glechoma hederacea</i>
Stoloniferous	25	<i>Lycopus exaltatus</i> , <i>Scutellaria scordiifolia</i>

Table 6. Grasses diversity in Lamiaceae

In multi-species genera like *Dracocephalum*, *Scutellaria*, and *Thymus*, a range of life forms is observed (see Fig. 3). For example, *Dracocephalum* species can exhibit dwarf semi-shrub or herbaceous biomorphs. The herbaceous type further encompasses various forms, such as short-rhizomatous, long-rhizomatous, perennial tap-root caudex, and annual taproot. The majority of *Dracocephalum* species develop as rhizomatous (short- or long-rooted) plants. In contrast, *Scutellaria* species are known for forming dwarf semi-shrub and herbaceous (long-rhizomatous and subterranean-stoloniferous) life forms. *Thymus* species, on the other hand, predominantly exhibit shrubby or semi-shrubby life forms.

A research on the phenological development of Lamiaceae species in the West Siberian province found that all these plants exhibit a similar phenorhythmotype.

They are summer-green plants that become dormant in winter. These plants bloom from spring to fall and lose their leaves during the winter months.




Figure 3. Life forms in three multispecies taxa (*Dracocephalum*, *Scutellaria* and *Thymus*).

Discussion

In the West Siberian province, there are 80 species of the Lamiaceae family, which account for 54% of their total number in Siberia. Their distribution across the province is uneven, with a concentration in the Barnaul district, known for its boreal and forest-steppe vegetation. The Barnaulsky District boasts 71 species from 29 genera, rivaling and sometimes surpassing neighboring regions in species diversity. For instance, the Altai Republic has 67 species from 24 genera (Doronkin, Ebel 2012), Tuva has 60 species from 22 genera (Timokhina, Zykova 2007), and Khakassia has 61 species from 20 genera (Myadele 2008). Genera such as *Dracocephalum*, *Thymus*, and *Scutellaria* are known for their multiple species. Most Lamiaceae species in the West Siberian province have wide ranges, with only 4% being endemics or subendemics. The family's representatives are found in various coenotic complexes, and 93% of species thrive in a wide range of ecological and coenotic habitats. In terms of ecological and phytocoenotic features, all species were grouped using clustering (Ward's method) into two clusters (Fig. 4).

Group 1 includes three species (*Dracocephalum grandiflorum*, *D. imberbe*, and *Phlomis oreophila*) primarily found in high mountain cold conditions, although they are also present in meadow and forest communities. These species have different herbaceous life forms but share similar morphological features in their shoot systems.

The second group encompasses the remaining guboflower species, which display a diverse range of features. This group can be further divided into four subgroups. Notably, subgroup 2c consists of 37 species mainly categorized as eumesophytes or euxerophytes, with varying light regime preferences and the ability to grow on different soil types. These species are commonly found in meadow, petrophytic, and anthropophytic communities. They exhibit a high diversity of life forms and morphological characters. Subgroup 2d comprises 24 species associated with forest, steppe,

and petrophytic communities, with a prevalence of annual grasses. Subgroup 2b includes four species with semi-shrubby/shrubby life forms, while subgroup 2a consists of ten species with perennial herbaceous life forms. Some species within the second group share a high degree of similarity in ecological and phytocoenotic features, even across different genera. For example, *Stachys sylvatica* L., *Scutellaria hastifolia* and *Prunella vulgaris* are closely related.




Figure 4. Similarity of ecological-phytocoenotic preferences in Lamiaceae species of the West Siberian province.

The comparison of our results with existing literature on the biomorphology and ecological-phytocoenotic preferences of hibiscus species in Siberia (Myadel 2008; Vodolazova et al. 2010; Kolegova, Cheryomushkina 2013; Cheryomushkina, Guseva 2015; Denisova et al. 2018; Talovskaya et al. 2018; Cheryomushkina et al. 2020; Komarevtseva 2020) revealed that the diverse life forms, prevalence of herbaceous perennials, and variation in morphological characteristics are key factors that have enabled the family's representatives to thrive in this region. Consequently, Lamiaceae species in the West Siberian province exhibit a broad range of ecological and coenotic preferences, spanning from flat steppes to mountain tundra, and adapting to various substrate types and light conditions. The distribution of these species is determined by the diversity of life forms, with perennial grasses being the most common, and biomorphological traits.

Conclusion

A comprehensive approach was applied to study the Lamiaceae family in the West Siberian floristic province. This approach helped determine the taxonomic composition, distribution, coenotic

distribution, ecological preferences, and biomorphology of the species. Among the 30 genera in the study area, the leading genera in terms of species number are *Thymus* L., *Dracocephalum* L., *Scutellaria* L., and *Mentha* L. The species exhibit a wide range of ecological and coenotic preferences. They are mainly found in boreal and forest-steppe vegetation communities, equally distributed in both lighted and shaded areas, and prefer a narrow range of humidity changes. Most Lamiaceae species are nonpetrophytes growing on various types of soils. There are 7 different life forms observed among the species, with herbaceous perennial long-rhizomatous being the most common. The study also revealed common morphological features such as monocyclic elongated shoots, orthotropic growth direction, terminal arrangement of inflorescences, sympodial development, appearance of replacement axes due to basiton branching, and a long development period of more than three years. The diversity of life forms and biomorphological traits in the shoot system is influenced by the spectrum of ecological and coenotic conditions in the West Siberian province, reflecting species adaptation to these conditions.

Acknowledgements

The study was funded by the Russian Science Foundation grant No. 23-24-00254 (<https://rscf.ru/project/23-24-00254>).

References

- Adylov TA (1987) Field guide on Central Asia plants. Critical outline of the flora. Vol. 8. Fan Publishing House of the Uzbek SSR, Tashkent, 402 pp. [In Russian]
- Anthelme F, Cavieres LA, Dangles O (2014) Facilitation among plants in alpine environments in the face of climate change. *Frontiers in Plant Sciences* 5: 387. <https://doi.org/10.3389/fpls.2014.00387>
- APG IV (2016) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG IV. *Botanical Journal of Linnean Society* 181: 1–20. <https://doi.org/10.1111/j.1095-8339.2009.00996.x>
- Astashenkov AYu (2015) Morphological adaptation of *Nepeta pamirensis* Franch. (*Lamiaceae*) to the conditions of the Pamir Mountains. *Contemporary Problems of Ecology* 6 (5): 636–646. <https://doi.org/10.15372/SEJ20150510>[In Russian]
- Barthélémy D, Caraglio Y (2007) Plant architecture: a dynamic, multilevel and comprehensive approach to plant form, structure and ontogeny. *Annals of Botany* 99 (3): 375–407. <https://doi.org/10.1093/aob/mcl260>
- Budantsev AL (1990) Systematics of the genus *Nepeta* (Lamiaceae). Species of the section *Spicata*. *Botanical Journal* 75(7): 1004–1013. [In Russian]
- Budantsev AL (Ed.) (2011) Plant resources of Russia. Wild flowering plants, their component composition and biological activity. Part 4. Family Caprifoliaceae-Lobeliaceae. Partnership of Scientific Publications KMK, St. Petersburg-Moscow, 630 pp. [In Russian]
- Carović-Stanko K, Petek M, Grdiša M, Pintar J, Bedeković D, Čustić MH, Satovic Z (2016) Medicinal plants of the family Lamiaceae as functional foods – a review. *Czech Journal of Food Sciences* 34: 377–390. <https://doi.org/10.17221/504/2015-CJFS>
- Cheryomushkina VA, Guseva AA, Makunina NI, Astashenkov AY, Denisova GR (2020) Phytocenotic characterization, ontogenetic structure and assessment of the state of *Scutellaria scordiifolia* (Lamiaceae) cenopopulations in Siberia. *Plant Resources* 2: 138–150. <https://doi.org/10.7868/S0033994618040031>

Cheryomushkina V, Guseva A, Talovskaya E, Astashenkov A (2022) Dwarf subshrub morphological structure variety in species of the genus *Scutellaria* (Lamiaceae) under different growing conditions. *Taiwania* 67(1): 146–154. <https://doi.org/10.6165/tai.2022.67.146>

Cheryomushkina VA Guseva AA (2015) Life forms of *Scutellaria supina* L. (Lamiaceae). *Contemporary Problems of Ecology* 8(5): 624–635. <https://doi.org/10.1134/S1995425515050054>

Cheryomushkina VA, Astashenkov AYu, Saidov DS (2020) Parallelism in the development of life forms of species of the genus *Kudryaschevia* (Lamiaceae): ontogenes and architectural analysis. *Contemporary Problems of Ecology* 13 (3): 257–265. <https://doi.org/10.1134/S1995425520030038>

Denisova GR, Cheremushkina VA, Astashenkov AYu, Talovskaya EB (2018) Ontomorphogenesis and assessment of the state of *Dracocephalum argunense* Fisch. et Link. cenopopulations at the border of the areal. *Botanical Journal* 103(4): 427–440. <https://doi.org/10.1134/S000681361618040014>

Doronkin VM (2005) The family Lamiaceae Martinov, or Labiatae Juss. In: Baikov KS (Ed.) *Conspectus of the flora of Siberia: Vascular plants*. Siberian Branch Russian Academy of Sciences Publishing House, Novosibirsk, 413–429. [In Russian]

Doronkin VM (2012) The family Lamiaceae Martinov, or Labiatae Juss. In: Baikov KS (Ed.) *Conspectus of the flora of Asian Russia: Vascular plants*. Publishing House of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk, 413–428. [In Russian]

Doronkin VM, Ebel AL (2012) The family Lamiaceae. In: Krasnoborov IM, Artemov IA (Eds) *Field guide on Altai Republic plants*. Siberian Branch Russian Academy of Sciences Publishing House, Novosibirsk, 383–401. [In Russian]

Ermakov AB (2003) Diversity of boreal vegetation of North Asia. Hemoboreal forests. Classification and ordination. Siberian Branch Russian Academy of Sciences Publishing House, Novosibirsk, 232 pp. [In Russian]

GBIF. Available from: <https://www.gbif.org/ru/>.

Harley RM, Atkins S, Budantsev AL, Cantino PD, Conn BJ, Grayer R, Harley MM, de Kok R, Krestovskaja T, Morales R, Paton AJ, Ryding O, Upson T (2004) Labiatae. In: Kadereit JW (Ed.) *The Families and Genera of Vascular Plants. VII Flowering Plants-Dicotyledons, Lamiales, except Acanthaceae including Avicenniaceae, Labiatae*. Springer-Verlag. Berlin, 167–275.

Hedge IC (1992) A global Survey of the biogeography of the Labiatae. In: Harvay RM, Reynolds T (Eds) *Advances in Labiatae Science*. Royal Botanical Garden. Kew, 7–17.

Jamzad Z (2013) A survey of Lamiaceae in the flora of Iran. *Rostaniha* 14(1): 59–67.

Kamelin RV (2017) *Flora of the Earth: floristic zoning of the land*. Five Plus LLC, Barnaul, 129 pp. [In Russian]

Kamelin RW, Makhmedov AM (1990) The system of the genus *Phlomooides* (Lamiaceae). *Botanical Journal* 75(2): 241–250. [In Russian]

Klokov MV (1954). The genus *Thymus* L. In *Flora of the USSR*. Part 21. USSR Academy of Sciences Publishing House, Moscow-Leningrad, 470–557. [In Russian]

Kolegova EB, Cheremushkina VA (2013) Ontomorphogenesis of vegetatively motile shrubs of the genus *Thymus* L. (Lamiaceae) in Khakassia. *MOIP Bulletin. Biological Series* 118(2): 70–76. [In Russian]

Russian]

Komarevtseva EK (2020) Cyclicity and morphogenesis of shoots of *Phlomis tuberosa* (Lamiaceae) in different growing conditions in southern Siberia. Journal of Botany 105(8): 81-93. <https://doi.org/10.31857/S0006813620080098>

Körner C (1992) Response of alpine vegetation to global climate change. Catena. Supp (Giessen) 22: 85-96.

Kuminova AV (1960) Vegetation cover of Altai. Publishing House of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk, 450 pp. [In Russian]

Lavrenko EM, Korchagin AA (1964) Field Geobotany. Vol. 3. USSR Academy of Sciences Publishing. Moscow-Leningrad, 146-208. [In Russian]

Laz'kov GA (2016) Labiatae (Family Labiatae Juss.) in flora of Kyrgyzstan. Vol. 1. In: KH Botanical Monograph & Revision Series. Korea National Arboretum, 375 pp.

Li X-W, Hedge IC (1994) Lamiaceae (Labiatae). In: Flora of China. Part 17. Science Press-Missouri Botanical Garden, Beijing-St. Louis, 50-299.

Makunina NI (2016) Vegetation of the forest-steppe of the West Siberian Plain and the Altai-Sayan mountain region: classification, structure and botanical and geographical patterns. Academic Publishing House "Geo", Novosibirsk, 183 pp. [In Russian]

Malyshev LI (1997) Flora of Siberia: Pyrolaceae - Lamiaceae (Labiatae). Part 11. Publishing house of the Siberian Branch of the Russian Academy of Sciences, Novosibirsk, 296 pp. [In Russian]

Malyshev LI, Baikov KS, Doronkin VM (2000) Floristic division of Asian Russia on the basis of quantitative characters. Krylovia 2(1): 3-16. [In Russian]

Mamadaliyeva NZ, Akramov DKh, Ovidi E, Tiezzi A, Nahar L, Azimova SS, Sarker SD (2017) Aromatic medicinal plants of the Lamiaceae family from Uzbekistan: ethnopharmacology, essential oils composition and biological activities. Medicines 4(8): 2-12. <https://doi.org/10.3390/medicines4010008>[In Russian]

Mathiesen C, Scheen AC, Lindqvist C (2011) Phylogeny and biogeography of the lamioid genus *Phlomis* (Lamiaceae). Kew Bulletin 66: 83-99.

Myadel MA (2008) Guboflowers of Khakassia: species composition, ecology and prospects of use. Thesis of Doctoral Dissertation. Khakassia State University. Abakan. [In Russian]

Paton A (1990) A global taxonomic investigation of *Scutellaria* L. Kew Bulletin 45: 399-450. <https://doi.org/10.2307/4110512>

Plants of the World. Available from: <https://powo.science.kew.org/>

Pyšek P, Liška J (1991) Colonization of *Sibbaldia tetrandra* cushions on alpine scree in the Pamiro-Alai Mountains. Central Asia. Arctic, Antarctic, and Alpine Research 23(3): 263-272. <https://doi.org/10.2307/1551603>

Ranjbar M, Mahmoudi C (2013) Chromosome numbers and biogeography of the genus *Scutellaria* L. (Lamiaceae). Caryologia: International Journal of Cytology, Cytosystematics and Cytogenetics 66(3): 205-214. <http://doi.org/10.1080/00087114.2013.821840>

Ranjbar M, Mahmoudi C (2018) A taxonomic revision of *Scutellaria* sect. *Lupulinaria* subsect. *Lupulinaria* (Lamiaceae) in Iran. Feddes Repertorium 128: 63–101.
<http://dx.doi.org/10.1002/fedr.201600027>

Raunkiaer CC (1934) The life-forms of plants and statistical plant geography: Being the collected papers of C. Raunkiaer. Clarendon Press, Oxford.

Safhhani K, Jamzad Z, Saeidi H (2018) Phylogenetic relationships in Iranian *Scutellaria* (Lamiaceae) based on nuclear ribosomal ITS and chloroplast trnLF DNA data. Plant Systematics and Evolution 304: 1077–1089. <https://doi.org/10.1007/s00606-018-1533-0>

Salimov RS, Karimov V, Babayeva Z, Gasimova G, Aliyeva Z, Akbarova A, Fatdayeva A, Nabiyeva P, Isgandarova L, Hajiyeva S (2018) Vascular plants of Azerbaijan: a nomenclatural update and survey of Lamiaceae L. Plant & Fungal Research 1(1): 69–85. <http://dx.doi.org/10.29228/plantfungalres.37>

Serebryakov IG (1959) Types of shoot development in herbaceous perennials and factors of their formation. Questions of plant biology: Scientific Notes of the Moscow City Pedagogical Institute named after V.P. Potemkin 100 (5): 3–38. [In Russian]

Serebryakov IG (1962) Ecological morphology of plants. Vysshaya Shkola Publishing House. Moscow, 378 pp. [In Russian]

Serebryakov IG (1964) Life forms of higher plants and their study. Field Geobotany. USSR Academy of Sciences Publishing House. Moscow-Leningrad, 146–205. [In Russian]

Shishkin BK (Ed.) (1954) Flora of the USSR. Vol. 21. Labiatae Juss. of the genera *Eremostachys-Plectrancthus*. USSR Academy of Sciences Publishing House. Moscow-Leningrad [In Russian]

Shishkin BK, Yuzepchuk SV (1954) Flora of the USSR. Vol. 20 Labiatae Juss. of the genus *Ajuga-Neustruevia*. USSR Academy of Sciences Publishing House. Moscow-Leningrad, 555 pp. [In Russian]

Siegel S, Castellan NJ (1989) Nonparametric statistics for the behavioral sciences. McGraw-Hill, International Editions. New York, USA.

Silantieva MM (2013) Conspectus of the flora of the Altai region. Altai University Publishing House. Barnaul, 392 pp. [In Russian]

Takhtajyan AL (1978) Floristic regions of the Earth. Nauka Publishing House. Leningrad, 247 pp. [In Russian]

Talovskaya (Kolegova) EB (2015) Morphological Transformation of *Thymus baicalensis* (Lamiaceae). Contemporary Problems of Ecology 8(5): 607–613.
<https://doi.org/10.1134/S1995425515050133>[In Russian]

Talovskaya E, Cheryomushkina V (2022) Morphological variations of *Thymus* L. in the vegetation belts of the Tien Shan mountains (Central Asia). Botany 100(6): 499–508.
<https://doi.org/10.1139/cjb-2021-0101>

Talovskaya EB, Cheryomushkina VA, Denisova GR (2018) Morphological adaptation of *Thymus* (Lamiaceae) species in Yakutia. Contemporary Problems of Ecology 11(6): 624–634.
<https://doi.org/10.15372/SEJ20180608>

Timokhina SA, Zykova EY (2007) Family Lamiaceae. In: Shaulo DN (Ed.) Field guide on the Tyva

Republic plants. Publishing House of the Siberian Branch of the Russian Academy of Sciences. Novosibirsk, 428-446. [In Russian]

Vodolazova SV, Cheremushkina VA, Kolegova EB, Myadele MA (2010) Ontogeny, structure of cenopopulations and ecological and cenotic characterization of *Nepeta sibirica* L. in Khakassia. Plant Resources 46(1): 3-15. [In Russian]

Warming E (1909) Oecology of plants. Clarendon Press, Oxford.
<https://doi.org/10.5962/bhl.title.23133>

Zhao F, Chen YP, Salmaki Y, Drew BT, Wilson TC, Scheen AC, Celep F, Bräuchler C, Bendiksby M, Wang Q, Min DZ, Peng H, Olmstead RG, Li B, Xiang CL (2021) An updated tribal classification of Lamiaceae based on plastome phylogenomics. BMC Biology 19 (2):1-27.
<https://doi.org/10.1186/s12915-020-00931-z>

Zhao F, Li BI, Drew BT, ChenY-P, Wang Q et al (2020) Leveraging plastomes for comparative analysis and phylogenomic inference within Scutellarioideae (Lamiaceae). PLoS ONE 15 (5): e0232602. <https://doi.org/10.1371/journal.pone.0232602>