

Complexes of terrestrial molluscs of various biotopes in the Gissar Range, Uzbekistan

Jasur Kudratov

Department of Biology, Gulistan State University, Gulistan City 4-microdistrict, 120101, Gulistan, Uzbekistan;
Department of Zoology, Samarkand State University named after Sharof Rashidov, 15 University Boulevard, Samarkand, 140104, Uzbekistan

Abduvaeit Pazilov

Department of Biology, Gulistan State University, Gulistan City 4-microdistrict, 120101, Gulistan, Uzbekistan

Dilnoza Zokirova

Department of Zoology, Samarkand State University named after Sharof Rashidov, 15 University Boulevard, Samarkand, 140104, Uzbekistan

Bekzod Otakulov

Department of Ecology and Life Safety, Samarkand State University, named after Sharof Rashidov, 15 University Boulevard, Samarkand, 140104, Uzbekistan

Jamshed Jalilov

Termiz Branch of Tashkent Medical Academy, Termiz, Uzbekistan

Xulkar Urinova

Department of Life Safety, Navoi State Mining and Technology University, Navoi, 210100, Uzbekistan

Fazlitdin Khalimov

Department of Biology, Gulistan State University, Gulistan City 4-microdistrict, 120101, Gulistan, Uzbekistan;
Department of Zoology, Samarkand State University named after Sharof Rashidov, 15 University Boulevard, Samarkand, 140104, Uzbekistan; Kattakurgan branch of Samarkand State University, Kattakurgan, Uzbekistan

Zuvaidullo Izzatullaev

Department of Ecology and Life Safety, Samarkand State University, named after Sharof Rashidov, 15 University Boulevard, Samarkand, 140104, Uzbekistan

The article presents the results of studying the complexes of terrestrial molluscs of various biotopes in the Gissar Range, Uzbekistan. Taxonomic identification of terrestrial molluscs collected from 13 biotopes in 4 altitude regions, determination of individual species' density, and assessment of the similarity of species content among biotopes was carried out. Forty-six species of molluscs are found in the studied biotopes. Biotopes no. 5 (along ditches among thickets of grass, under stones), no. 10 (near springs among grasses), and no. 11 (on the banks of small streams among thickets of grasses) were found to maintain the highest species richness. The most common species in the studied biotopes are *Cochlicopa lubrica*, *Vallonia costata*, and *Pupilla muscorum*.

Acta Biologica Sibirica 10: 349–359 (2024) doi: 10.5281/zenodo.11160044

Corresponding author: Jasur Kudratov (kudratovzasur484@gmail.com)

Academic editor: R. Yakovlev | Received 22 March 2024 | Accepted 12 April 2024 | Published 11 May 2024

<http://zoobank.org/577BCC6E-76A8-4C68-81EC-54D06717777C>

Citation: Kudratov J, Pazilov A, Zokirova D, Otakulov B, Jalilov J, Urinova X, Khalimov F,

Izzatullaev Z (2024) Complexes of terrestrial molluscs of various biotopes in the Gissar Range, Uzbekistan. *Acta Biologica Sibirica* 10: 349–359. <https://doi.org/10.5281/zenodo.11160044>

Keywords

Biotopes, density, Gissar Range, species composition, terrestrial molluscs

Introduction

Mollusca is the second most diverse animal phylum occurring in all major environments except the aerospace (Pazilov et al. 2022). They have significant functions in food webs and ecosystem equilibrium, such as nutrient cycling, biofiltration, and storage (Fatemeh et al. 2019; Kudratov et al. 2023). Molluscs are animals that have the potential to be able to monitor species stability and the environment; they can also be bioindicators of ecosystem changes (Isoni et al., 2023). The most species-rich class of Mollusca is Gastropoda, comprising 80% of this phylum (Bouchet, Rocroi, 2005; Bouchet et al., 2017; Islamy et al., 2020; Abdurasulova and Pazilov, 2021). It is known that the biological diversity of plants and animals in mountain regions uniquely differs from other biotopes, and the number and variety of species are different (Schilkeyko et al. 2017; Akramov et al. 2023; Khalimov 2023). One of the areas with a high probability of changes due to anthropogenic influence is the mountain ranges and their adjacent areas (Khalimov 2020).

Complexes of terrestrial molluscs of various biotopes of the southern slope of the Gissar ridge (Uzbekistan) were analyzed in detail by Z. Izzatullaev (Izzatullaev 1970; Izzatullaev 1975). However, over the past 30 years, the vegetation cover has changed dramatically under the influence of human activity: development for agriculture, logging, and grazing. The exploitation of woody vegetation has led to a reduction in its area and noticeably disrupted the structure of plantings. The herbaceous vegetation of the ridge has changed significantly as a result of grazing and, probably, the excessive overload of many pastures for a long time. Similar changes, although to a lesser extent, occurred in high-mountain heath meadows and groups of alpine short-grass and other heaths located even in the most remote, inaccessible areas.

It should be noted that these changes directly affect the ecology and distribution of terrestrial molluscs. Geobotanically, the slopes of the ridge are heterogeneous. According to the characteristics of Zakirov, high-altitude zones are divided into Desert, Hill, Mountain, and Pasture (Zakirov 1955).

The climate of the mountain region is dry continental, which caused the formation of unique flora and fauna in this region. It has been proven in several studies that the biological diversity of animals (Schilkeyko et al., 2021; Pazilov and Umarov, 2021; Baymuradov et al., 2021; Zokirova and Khalimov, 2022; Khalimov et al. 2023;) and plants (Bazarov et al. 2023; Ruziev et al. 2023; Alikulov et al. 2022) in the mountainous regions of Uzbekistan is high. However, information on the current species content and distribution of the malacofauna of the Gissar mountain range in different biotopes is practically absent from scientific literature and other sources, which indicates that no in-depth research has been conducted in this regard.

Therefore, our research aimed to determine the species composition of terrestrial malacofauna in different biotopes of the Gissar mountain range.

Materials and methods

The research area comprised 13 biotopes: 1 – in gardens and vegetable gardens, in grass, in fallen leaves and rotting plant debris; 2 – on the banks of rivers and ditches among thickets of grass; 3 – on undeveloped lands in semi-shrubs and the stems of grass thickets; 4 – at the foot of the slopes,

among semi-shrubs; 5 - along ditches among thickets of grass, under stones; 6 - close to streams. Among bushes and under rocks; 7 - in tree and shrub vegetation on gravelly slopes; 8 - in bushes with screes; 9 - on rocks and screes with plants; 10 - near springs among grasses; 11 - on the banks of small streams among thickets of grasses; 12 - on rocks and screes; 13 - in subalpine meadows in the Gissar Range of Uzbekistan. A complete description of the sampling station is shown in Fig. 1.

The sampling and processing of the malacological material were carried out using the methods of Izzatullayev (2019) and Pazilov and Umarov (2021). The collected snails were fixed with 70-75% alcohol. If it was impossible to fix them with alcohol, they were fixed in 2% formalin, and soda (NaHCO_3) was added to the mixture to prevent the shells from dissolving. All shells are kept dry in the malacological collection of Samarkand State University.

Likharev's (1952) and Schileyko (1978) keys were used to identify molluscs. After being photographed and identified, the animals were stored in the collection bottle. The method used to identify mollusc species consisted of morphological and anatomical analysis. Molluscs were identified based on the morphology of their shells and the structure of genital organs. At least five specimens from each species was dissected to determine differences in genitalia, including the penis.

The similarity of the species composition of the investigated biotopes was analyzed by the Neighbor-joining method (Yan et al. 2021) and described using the cluster prepared in the PAST 4.0 program (Hammer 2001).

Result

During the study, 46 terrestrial molluscs were collected (Table 1).

Desert. Located at altitudes of 650–900 m above sea level. The malacofauna of this belt has been studied in the following biotopes: gardens and orchards, banks of rivers, and ditches.

In gardens and orchards, on grass, in fallen leaves, and in rotting plant remains were found the following species: *Deroceras laeve* (5 individuals per m^2), *D agreste* (6), *Candacharica levanderi* (3).

Along the banks of rivers and ditches, among thickets of grasses, we identified *Cochlicopa nitens* (10 individuals per m^2), *C. lubrica* (6), *Vallonia costata* (11), *Pupilla muscorum* (14), *Angomphalia regeliana* (4), *Candacharica levanderi* (3), *Zonitoides nitidus* (9).

On undeveloped lands in subshrubs and the stems of grass thickets, *Xeropicta candaharica* forms dense populations (25 individuals per m^2).




Figure 1. A map of the studied area.

Thus, in the desert belt we found 10 species of terrestrial mollusks. In terms of population density, *Xeropicta candaharica* dominates. But *Candacharica levanderi* became more widespread since this species was recorded in two biotopes of the desert belt.

Hill covers the entire hilly-undulating foothill space with a loess cover and occupies most of the territory of the northwestern part of the Gissar ridge. Hill, according to the scheme of Zakirov (1955), is divided into two subzones: 1) lower Hill, where the relief is relatively flat, 500–900–1200 m; 2) upper, where the relief is relatively sharp with frequent exposures of subsoil rocks; its height is 1000–1500 m. The vegetation of the lower Hill consists of the following ranks: *Philomis thapsoides*, *Prosalea drupacea*, etc. The vegetation of the upper Hill is represented by almond trees, which survived only in places more or less remote from populated areas.

In the Hill belt we studied the following biotopes: at the foot of the slopes, among subshrubs: *Pseudonapaeus Sogdiana* (17 individuals per m²), *Sphyradium doliolum* (13), *Chondrulopsina intumescens* (9), *Xeropicta candaharica* (15), *Gibbulnopsis signata* (21).

Along the ditches among thickets of grasses, under stones, the following were found: *Cochlicopa nitens* (11 individuals per m²), *C. lubrica* (9), *Pupilla muscorum* (18), *Vertigo pygmaea* (5), *Phenacolimax annularis* (13), *Angiomphalia regeliana* (4), *D. laeve* (3), *D. agreste* (5), *Zonitoides nitidus* (5), *Vallonia costata* (15).

Not far from streams, among bushes, under stones, *C. levanderi* (5), *C. roseni* (7), and *C. kaznakovi* (3) are living.

In total, 17 species have been recorded in the Hill belt.

No	Species name	Desert			Adyr			Mountain				Pasture		
		1	2	3	4	5	6	7	8	9	10	11	12	13
1	<i>Cochlic</i>	-	+	-	-	+	-	-	-	-	+	+	-	-

	<i>opa nitens</i>													
2	<i>C. lubrica</i>	-	+	-	-	+	-	-	-	-	+	+	-	+
3	<i>C. lubricella</i>	-	-	-	-	-	-	-	-	-	-	+	-	-
4	<i>Sphyradium doliolum</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
5	<i>Vallonia costata</i>	-	+	-	-	+	-	-	-	-	+	+	-	+
6	<i>V. pulchella</i>	-	-	-	-	-	-	-	-	-	+	+	-	-
7	<i>V. ladaensis</i>	-	-	-	-	-	-	-	-	+	-	-	-	-
8	<i>Gibbulinopsis signata</i>	-	-	-	+	-	-	+	-	-	-	-	-	-
9	<i>Pupilla triplicata</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
10	<i>P. muscorum</i>	-	+	-	-	+	-	-	-	-	+	+	-	+
11	<i>P. sterrii</i>	-	-	-	-	-	-	-	-	-	-	-	+	-
12	<i>P. anzobica</i>	-	-	-	-	-	-	-	-	+	-	-	+	-
13	<i>Vertigo antvertigo</i>	-	-	-	-	-	-	+	-	-	+	-	-	-
14	<i>V. pygmaea</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
15	<i>Columella columella</i>	-	-	-	-	-	-	-	-	-	+	-	-	-
16	<i>Pyramidula rustris</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
17	<i>Pseudonapaes albiplicata</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
18	<i>Ps. sovgiana</i>	-	-	-	+	-	-	-	-	-	-	-	-	-
19	<i>Ps. miser</i>	-	-	-	-	-	-	-	-	-	-	-	+	-
20	<i>Ps. kasnakovi</i>	-	-	-	-	-	-	-	+	-	-	-	-	-
21	<i>Ps. ostomus</i>	-	-	-	-	-	-	-	+	-	-	-	-	-
22		-	-	-	-	-	-	-	-	-	-	-	+	-

	<i>Turane na mar tensian a</i>													
23	<i>T. con icula</i>	-	-	-	-	-	-	-	-	-	-	-	+	-
24	<i>Phenac olimax annula ris</i>	-	-	-	-	+	-	-	-	-	-	-	-	-
25	<i>Chondr ulopsin a intu mescen s</i>	-	-	-	+	-	-	-	-	-	-	-	-	-
26	<i>Ch. fe dtsche nko i</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
27	<i>Leucoz onella rubens</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
28	<i>L. me soleuca</i>	-	-	-	-	-	-	-	+	-	-	-	-	-
29	<i>L. rufi spira</i>	-	-	-	-	-	-	-	-	+	-	-	-	-
30	<i>L. retteri</i>	-	-	-	-	-	-	-	-	+	-	-	-	-
31	<i>L. caria</i>	-	-	-	-	-	-	-	-	+	-	-	-	-
32	<i>Xeropi cta can daharic a</i>	-	-	+	+	-	-	-	-	-	-	-	-	-
33	<i>Angio mphali a regel iana</i>	-	+	-	-	+	-	-	-	-	-	-	-	-
34	<i>Deroce ras laeve</i>	+	-	-	-	+	-	-	-	-	+	-	-	-
35	<i>D. agreste</i>	+	-	-	-	+	-	-	-	-	-	-	-	-
36	<i>Lytopel te mac ulata</i>	-	-	-	-	-	-	+	-	-	-	-	-	-
37	<i>Canda haria r utellu m</i>	-	-	-	-	-	-	-	-	-	+	-	-	-
38	<i>C. lev anderi</i>	+	+	-	-	-	+	-	-	-	-	-	-	-
39	<i>C. roseni</i>	-	-	-	-	-	+	-	-	-	-	-	-	-
40	<i>C. kas nakovi</i>	-	-	-	-	-	+	-	-	-	-	-	-	-

41	<i>Macrochlamys turanica</i>	-	-	-	-	-	-	-	-	+	-	-	-	-
42	<i>M. sogdiana</i>	-	-	-	-	-	-	-	+	-	-	-	-	-
43	<i>Zonitoides nitidus</i>	-	+	-	-	+	-	-	-	-	+	-	-	-
44	<i>Novisuccinea evoluta</i>	-	-	-	-	-	-	-	-	-	-	+	-	-
45	<i>Pamirsuccinea eximia</i>	-	-	-	-	-	-	-	-	-	-	+	-	-
46	<i>Oxyloma elegans</i>	-	-	-	-	-	-	-	-	-	-	+	-	-

Table 1. Distribution of terrestrial molluscs across vertical belts and biotopes of the Gissar Range

Mountain. According to natural-historical conditions, this belt can be divided into two subzones.

1. The lower Mountain subzone – a transitional strip from the underlying adyr zone – is characterized by ephemeral plants with more or less xerophilic long-growing steppe-type dominants. Tree and shrub forms are displaced for various reasons, mainly due to human activity. The height of the strip is 1400–1800–2100 m above sea level.

In this subzone, the same complexes of terrestrial molluscs were found as in the Adyr belt.

2. There are almost no ephemeral plants in the upper subzone. Trees and shrubs are better developed here. Under similar conditions, the number of species of more or less mesophilic plants gradually increases.

In this subzone, molluscs live in various biotopes: on tree and shrub vegetation, on rocks and screes, and on the banks of streams and springs.

We found eight species in tree and shrub vegetation on gravelly slopes: *S. doliolum* (10 individuals per m²), *G. signata* (15), *Pupilla triplicata* (12), *Vertigo antivertigo* (12), *Pyramidula rupestris* (7), *Pseudonapaeus albiplicata* (4), *Chondrulopsina intumescens* (8), *L. maculata* (5).

In the biotope of shrubs with screes, six species are found: *Ps. kasnakovi* (5 individuals per m²), *Ps. otostomus* (4), *Ch. fedtschenkoi* (10), *Leucozonella rubens* (6), *L. mesoleuca* (7), *Macrochlamys sogdiana* (2).

Rocks and screes are poorer in mollusks than other biotopes. Here, among the stones and plant remains, six species are observed: *Leucozonella rufispira* (4 individuals per m²), *L. retteri* (3), *L. caria* (1), *Vallonia ladacensis* (10), *Macrochlamys turanica* (5), *P. anzobica* (10).

Along ditches and springs, as well as near streams among thickets of plants, 11 species live: *Cochlicopa nitens* (5 individuals per m²), *C. lubrica* (10), *C. lubricella* (9), *Vallonia costata* (11), *V. pulchella* (8), *Pupilla muscorum* (10), *Vertigo antivertigo* (7), *Columella columella* (6), *Deroceras*

laeve (3), *Candaharia rutellum* (2), *Zonitoides nitidus* (10).

In total, 32 species of terrestrial molluscs were found in the upper subzone of the Mountain belt.

Pasture. This belt consists of two sub-belts: the lower, usually called subalpine, and the upper, known in the literature as “alpine”. There are no conditions for the development of trees and shrubs in the zone.

The lower pasture belt is characterized by the development and distribution of mesophilic formations of mixed-grass meadows, while the upper belt is characterized by short-grass carpet lawns.

In the lower pasture belt, in contrast to the upper one, some woody plants are common, such as *Juniperus turkestanica*, *Sorbus tianschanica*, *Rosa betula*, etc. The malacological complexes on both subbelts of the Pasture are very similar. Therefore, the distribution of terrestrial molluscs in the two sub-belts is considered jointly. Terrestrial molluscs of the Mountain-Pasture belt inhabit the following biotopes: the banks of small streams, rocks, and screes, subalpine meadows:

In the biotope - banks of small streams among thickets of grasses the following were found: *Cochlicopa nitens* (10 individuals per m²), *C. lubrica* (7), *C. lubricella* (13), *Vallonia costata* (11), *V. pulchella* (15), *Pupilla muscorum* (10), *Novisuccinea evoluta* (3), *Pamirsuccinea eximia* (6), *Oxyloma elegans* (5).

On rocks and screes there are: *Pupilla sterrii* (8 individuals per m²), *P. anzobica* (2), *Pseudonapaeus miser* (5), *Turanena martensiana* (3), *T. conicula* (1).

The subalpine meadows contain mainly mesophilic interzonal species: *Cochlicopa lubrica* (5 individuals per m²), *Pupilla muscorum* (10), *Vallonia costata* (11).

In total, 14 species of terrestrial molluscs were found in the Pasture belt.

The density of terrestrial molluscs in biotopes varies. Data on the total density of terrestrial mollusks of the Gissar Range are shown in the figure 2. For example, in biotopes 4, 5, 7, 10, 11, the density of terrestrial molluscs varies from 75 to 81 specimens per m². The lowest density of molluscs is observed in biotopes 1, 6, 12 from 15 to 19 specimens per m².

As for species diversity, this indicator reaches its maximum value in biotopes 5, 10, 11 (from 9 to 11 species). The lowest species diversity is observed in biotopes 1, 3, 6, 13 (from 1 to 3 species).

When comparing the malacocomplexes using the coefficients of species similarity of the Gissar Range, it turned out that in more than half of the cases, the similarity coefficient is equal to 0 (Table 2 and Figure 3). In only two biotopes: on the banks of rivers and ditches among thickets of grass (2), along ditches among thickets of grasses, under stones (5), and also near springs among grasses (10), species similarity is only 54%. Also similar in the studied biotopes 5- and 13- (50%), 2- and 13- (42.8%), 10- and 11- (35.7%), 5- and 10- (33.3%) types are distributed.

We explain the large number of non-repeatable cases in the composition of species of the biotopes studied by the small area of the biotopes being compared and the fact that their structural components are different.




Figure 2. Density (histogram) and number of species (line) of terrestrial molluscs in various biotopes of the Gissar Range.




Figure 3. Dendrogram of faunistic similarity of terrestrial molluscs of various biotopes of the Gissar Range, Uzbekistan (based on Jaccard coefficient).

K	2	3	4	5	6	7	8	9	10	11	12	13
1	11	0	0	18	0	0	0	0	8.3	0	0	0
2		0	0	54	11.1	0	0	0	54	23.7	0	42.8
3			25	0	0	0	0	0	0	0	0	0
4				0	0	0	0	0	0	0	0	0
5					0	0	0	0	33.3	18.7	0	50
6						0	0	0	0	0	0	0
7							0	0	0	0	0	0

8								0	0	0	0	0
9									0	0	0	0
10										35.7	0	0
11											0	33.3
12												0

Table 2. Coefficient of species similarity of malacomplexes of different biotopes of the Gissar Range (Jaccard index)

Conclusion

For the first time, land molluscs of the Gissar mountain range were studied in 13 biotope sections in different altitudinal regions. A list of land mollusks including 46 species was compiled. Species-rich biotopes were identified, and species with high distribution potential (*Cochlicopa lubrica*, *Vallonia costata*, and *Pupilla muscorum*) were revealed. A comparative analysis of the biotopes of terrestrial molluscs is provided. The data obtained by the study of land mollusks in the section of biotopes differing in their altitude position, besides providing important facts about the distribution of these animals, explain the biological diversity of the region's fauna.

Acknowledgements

The author expresses his sincere gratitude to M. Rakhimov, associate professor of the Department of Zoology of Samarkand State University, for his advice in preparing the article.

References

- Abdurasulova S, Pazilov A (2021) Nature of variability of *Candaharia levanderi* (Simroth, 1902) in the Ferghana and Surkhan – Sherabad Valleys, Uzbekistan. Bulletin of the Iraq Natural History Museum 16 (4): 547–555. <http://dx.doi.org/10.26842/binhm.7.2021.16.4.0547>
- Alikulov BS, Shuryhin VV, Davranov KD, Ismailov ZF (2022) The halophytic plant *Halostachys belangeriana* (Moq.) Botsch as a source of plant growth-promoting endophytic bacteria. Microbiological Journal 4: 31–39. <http://dx.doi.org/10.15407/microbiolj84.04.030>
- Akramov I, Axanbayev Sh, Alikulov B, Mukhtorova S, Ergashev A, Ismailov Z (2023) Plant growth-promoting properties of endophytic bacteria isolated from some xerophytic plants distributed in arid regions (Uzbekistan). Plant Science Today 10(4): 228–237. <https://doi.org/10.14719/pst.2725>
- Baymuradov Kh, Zhabborova T, Tuinazarova I, Otakulov B, Egamkulov A (2021) Aquatic ecosystems of the lower reaches of the Zarafshan River. Diversity and ecological groups of molluscs. E3S Web of Conferences 262: 04009. <http://dx.doi.org/10.1051/e3sconf/202126204009>
- Bazarov B, Rajamuradov Z, Safin M, Rajabov A, Khayitov D, Kuziev M, Aminjonov S, Ismayilova M, Kudratov J, Khujabekov M, Khaydarov D (2023) The productivity, chemical composition and nutritional value of pastures dominated by *Artemisia diffusa* and *Cousinia resinosa* in arid lands of southwestern Uzbekistan. Biodiversitas 24: 3916– 3923. <https://doi.org/10.13057/biodiv/d240730>
- Bouchet P, Rocroi J-P (Eds) (2005) Classification and nomenclator of gastropod families. Malacologia: International Journal of Malacology 47(1–2): 1–397.
- Bouchet Ph, Rocroi J-P, Hausdorf B, Kaim A, Kano Y, Nützel A, Parkhaev P, Schrödl M, Strong EE (2017) Revised Classification, Nomenclator and Typification of Gastropod and Monoplacophoran Families. Malacologia 61(1–2): 1–526. <https://doi.org/10.4002/040.061.0201>

Fatemeh A, Sareh Y, Peter G, Mehrdad Z (2019) Ecology and species composition of Molluscs upstream of the Kor River System, with two new records for the Fars Province, Iran. *Journal of Wildlife and Biodiversity* 3 (2): 29–39. <http://dx.doi.org/10.22120/jwb.2019.105850.1062>

Hammer O (2001) PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica* 4 (1): 1–9.

Islamy RA, Hasan V (2020) Checklist of mangrove snails (Mollusca: Gastropoda) in South Coast of Pamekasan, Madura Island, East Java, Indonesia. *Biodiversitas* 21 (7): 3127–3134. <https://orcid.org/0000-0002-2541-4432>

Isoni W, Sari PDW, Sari LA, Daniel K, South J, Islamy RA, Wirabuana PYAP, Hasan V (2023) Checklist of mangrove snails (Gastropoda: Mollusca) on the coast of Lamongan District, East Java, Indonesia. *Biodiversitas* 24: 1676–1685. <http://dx.doi.org/10.13057/biodiv/d240341>

Izzatullaev Z (1970) Materials for studying terrestrial molluscs of the Gissar Range and adjacent regions of Tajikistan. *Proceedings of the Academy of Sciences of the Tajik SSR. Biology* 239: 79 – 86. [In Russian]

Izzatullaev Z (1975) Features of the distribution of terrestrial molluscs of the Gissar Range and adjacent regions of Tajikistan. Dushanbe, 212 pp. [In Russian]

Izzatullayev ZI (2019) Fauna of The Molluscs of Water Ecosystems of Central Asia and The Contiguous Country Territories. Lesson-Press Pub., Tashkent, 430 pp. [In Russian]

Khalimov F (2020) Seasonal dynamics of dominant species of soil predators (Coleoptera: Carabidae, Staphylinidae) in agro landscapes and their potential gluttony. *Travaux du Muséum National d’Histoire Naturelle “Grigore Antipa”* 63 (2): 175–187. <https://doi.org/10.3897/travaux.63.e54116>

Khalimov F, Rakhimov M, Khamzaev R, Abdullaev E, Usanov U (2023) Composition and structure of the entomofauna of *Ferula (Ferula kuhistanica)* in different sections of the Zarafshan Ridge. *Journal of the Entomological Research Society* 25(2): 275–286. <https://doi.org/10.51963/jers.2023.86>

Khalimov F (2023) Composition and structure of the fauna of ground beetles (Coleoptera, Carabidae) of the Zerafshan Range. *Acta Biologica Sibirica* 9: 113–125. <https://doi.org/10.5281/zenodo.7725474>

Kudratov J, Pazilov A, Maxammadiyev Z, Urazova R, Otakulov B, Bazarov B, Keldiyarov K, Soatova Z, Urinova X (2023) Diversity and ecology of molluscs (Gastropods) in mountain streams, Nurota mountain range, Uzbekistan. *Biodiversitas* 24(4): 2402–2408. <http://dx.doi.org/10.13057/biodiv/d240455>

Likharev IM, Rammelmeyer ES (1952) Terrestrial molluscs of the fauna of the USSR. Key to the fauna of the USSR 43. Nauka, Moscow-Leningrad, 511 pp.

Pazilov A, Umarov F (2021) On the ecology and species diversity of the freshwater gastropods of springs in Andijan Region, Uzbekistan. *Bulletin of the Iraq Natural History Museum* 16(3): 325–340. <https://doi.org/10.26842/binhm.7.2021.16.3.0325>

Pazilov A, Umarov F (2022) Conchological variability of terrestrial mollusc *Chondrulopsina fedtschenkoi* (Ancey, 1886) (Gastropoda, Pulmonata, Enidae) from the Zarafshan Range, Uzbekistan. *Bulletin of the Iraq Natural History Museum* 17 (1): 103–113. <http://dx.doi.org/10.26842/binhm.7.2022.17.1.0103>

Ruziyev F, Djabbarov I, Olimjonova S, Niyozov U, Urokov S, Ishankulova D, Bakhadirov U (2023)

Grain quality indicators and their phenotypic variability of ancient varieties of *Triticum aestivum* in the mountains of Uzbekistan. *Biodiversitas* 24: 5995–6001.

Schileyko AA (1978) Shellfish. Terrestrial molluscs of the superfamily Helicoidea. Fauna of the USSR 3 (6). Science, Leningrad branch, Leningrad, 384 pp. [In Russian]

Schileyko A, Pazilov A, Abdulazizova Sh (2017) A new genus of the Bradybaenidae family (Gastropoda, Pulmonata) from central Asia. *Ruthenica* 27(1): 31–37.
[https://doi.org/10.35885/ruthenica.2017.27\(1\).3](https://doi.org/10.35885/ruthenica.2017.27(1).3)

Schileyko A, Pazilov A, Abdulazizova Sh (2021) One more surprise from Bradybaenidae (Gastropoda, Pulmonata). *Ruthenica* 30(4): 217–221.
[https://doi.org/10.35885/ruthenica.2021.30\(4\).6](https://doi.org/10.35885/ruthenica.2021.30(4).6)

Yan H, Maozu G, Juan W (2021) ENJ algorithm can construct triple phylogenetic trees. *Mol Ther: Nucleic Acids* 23: 286–293. <https://doi.org/10.1016/j.omtn.2020.11.004>

Zakirov KZ (1955) Flora and vegetation of the Zarafshan river basin. Publishing House of the Academy of Sciences of Uzbekistan SSR, Tashkent, 212 pp. [In Russian]

Zokirova D, Khalimov F (2022) Morphometric features of the beetle *Acinopus (Acinopus) laevigatus* Menetries, 1982 (Coleoptera, Carabidae) in the mountain ecosystems of Uzbekistan. *Bulletin of the Iraq Natural History Museum* 17(2): 141–153.
<https://doi.org/10.26842/binhm.7.2022.17.2.0141>