

Lepidoptera of South Ossetia (Northern Transcaucasia). Part V. Superfamily Papilionoidea Latreille, 1809

Aleksandr N. Streltsov

Herzen State Pedagogical University of Russia, 48 Moika Emb., Saint Petersburg 191186, Russia

Petr Ya. Ustjuzhanin

Altai State University, 61 Lenina Ave., Barnaul 656049, Russia; Biological Institute, Tomsk State University, 36 Lenina Ave., Tomsk 634050, Russia

Kirill A. Kolesnichenko

Lomonosov Moscow State University, Leninskie Gory I/12, Moscow 119991, Russia

Roman V. Yakovlev

Altai State University, 61 Lenina Ave., Barnaul 656049, Russia; Biological Institute, Tomsk State University, 36 Lenina Ave., Tomsk 634050, Russia; Western Caspian University, 31, Istiglaliyyat Street, Baku, Azerbaijan

Seventy seven Papilionoidea species have been indicated for South Ossetia (72 – for the first time for this territory). For three species listed in the Red Book of South Ossetia (*P. apollo*, *I. podalirius* and *P. machaon*) new localities are given.

Acta Biologica Sibirica 10: 375–381 (2024)

doi: 10.5281/zenodo.11161306

Corresponding author: Aleksandr N. Streltsov (streltsov@mail.ru)

Academic editor: A. Matsyura | Received 23 April 2024 | Accepted 6 May 2024 | Published 12 May 2024

<http://zoobank.org/B226E122-0BA1-4394-A96B-7CF475001371>

Citation: Streltsov AN, Ustjuzhanin PYa, Kolesnichenko K, Yakovlev RV (2024) Lepidoptera of South Ossetia (Northern Transcaucasia). Part V. Superfamily Papilionoidea Latreille, 1809. Acta Biologica Sibirica 10: 375–381. <https://doi.org/10.5281/zenodo.11161306>

Keywords

Biodiversity, Caucasus, species richness, fauna, HesperIIDae, Papilionidae, Pieridae, Lycaenidae, Nymphalidae

Introduction

This article continues the series of publications devoted to the lepidopterofauna of South Ossetia and covers families of Papilionoidea (HesperIIDae, Papilionidae, Pieridae, Lycaenidae, and Nymphalidae). Families Adelidae, Incurvariidae, Psychidae, Tineidae, Roeslerstammiidae, Gracillariidae, Yponomeutidae, Argyresthiidae, Plutellidae, Ypsolophidae, Ethmiidae, Depressariidae, Elachistidae, Scythrididae, Cryptolechiidae, Oecophoridae, Lecithoceridae, Stathmopodidae, Coleophoridae, Momphidae, Blastobasidae, Autostichidae, Lypusidae,

Cosmopterigidae, Gelechiidae, Epermeniidae, Choreutidae, Pyraloidea, Cossidae, Limacodidae, part of Erebidae, Notodontidae, Lasiocampidae, Lemoniidae, Saturniidae, Sphingidae, Drepanidae, Cimeliidae, Tortricidae, Pterophoridae and Alucitidae were previously treated (Streltsov et al. 2022 a, b, c; Nedoshivina et al. 2023; Sinev et al. 2023). In the fifth part we publish data on Superfamily Papilionoidea; information about which in South Ossetia was not previously available.

A detailed description of the natural conditions and history of studying the lepidopterofauna of the region is contained in the first article in the series, dedicated to Lepidoptera (Streltsov et al. 2022b) and is not discussed here.

The butterflies (Papilionoidea) of South Ossetia are very poorly studied. The information on their distribution in the region is provided only for five species listed in the Red Data Book of South Ossetia (Bazaev et al. 2017). *Parnassius nordmanni* (Menetries, 1850) was reported from north of South Ossetia in the highlands of the Great Caucasus. *P. apollo* (Linnaeus, 1758) was given for the valleys of the Bolshaya and Malaya Liakhva rivers and other locations, mainly in the south of the territory at altitudes of 1200–2000 m. *Iphiclides podalirius* (Linnaeus, 1758) was indicated for Khetagurovo, Belot and Tsadykau villages. *Papilio machaon* Linnaeus, 1758 was regularly found in the Inner Kartli Valley. *Colias thisoa* Menetries, 1832 was spotted in the upper reaches of the Bolshaya and Malaya Liakhva rivers and tributaries of the Rioni river.

Materials and methods

The specimens were collected in South Ossetia in eight localities (Fig. 1) by A. N. Streltsov, P. Y. Ustjuzhanin and R. V. Yakovlev in June–July 2021 and by V.V. Rudoi, P. Y. Ustjuzhanin and R. V. Yakovlev in July 2022.

Methods are described in detail in the first part of the article (Streltsov et al. 2022a). The determination of Material was carried out according to Tshikolovets and Nekrutenko (2012).

The examined material is kept in the collections:

ASSP collection of Alexandr Streltsov (Saint-Petersburg, Russia);

CUK collection of Petr Ustjuzhanin and Vasiliy Kovtunovich (Novosibirsk, Moscow, Russia);

KKM collection of Kirill Kolesnichenko (Moscow, Russia);

RYB collection of Roman Yakovlev (Barnaul, Russia).

List of collecting localities

1. South Ossetia, Tskhinval Distr., 2 km NW Grom, 42°10'6" N / 44°11'53" E, 930 m, 22-25.06.2021, A. Streltsov, P. Ustjuzhanin & R. Yakovlev leg.
2. South Ossetia, Leningor Distr., 4 km E Leningor, 42°08'45" N, 44°30'55" E / 1200 m, 26-27.06.2021, A. Streltsov, P. Ustjuzhanin & R. Yakovlev leg.; 13- 14.07.2022, P. Ustjuzhanin & R. Yakovlev leg.
3. South Ossetia, Dzaus Distr., 4 km NNE Kvaisa, Koz lake, 42°33'32" N / 43°37'59" E, 1580 m, 28-30.06.2021, A. Streltsov, P. Ustjuzhanin & R. Yakovlev leg.
4. South Ossetia, Dzaus Distr., Rachinsky Range, near Dodtota, 42°27'25" N / 43°43'18" E, 1750 m, 1-2.07.2021, A. Streltsov, P. Ustjuzhanin & R. Yakovlev leg.; 19-22.07.2022, V. Rudoi, P. Ustjuzhanin & R. Yakovlev leg.

5. South Ossetia, Dzaus Distr., Dvalet Range, near Kherusel't, 42°32'37" N / 43°47'32" E, 1760 m, 3-5.07.2021, A. Streltsov, P. Ustjuzhanin & R. Yakovlev leg.

6. South Ossetia, Dzaus Distr., Mtiulet Range, near Erman, 42°31'2" N / 44°14'10" E, 2140 m, 7-9.07.2021, P. Ustjuzhanin & R. Yakovlev leg.; 26-27.07.2022, V. Rudoi, P. Ustjuzhanin & R. Yakovlev leg.

7. South Ossetia, Znaur Distr., 2 km W Dzagina, 42°14'34" N / 43°43'11" E, 1100 m, 11-12.07.2021, P. Ustjuzhanin & R. Yakovlev leg.; 15-16.07.2022, P. Ustjuzhanin & R. Yakovlev leg.

8. South Ossetia, Dzaus Distr., Tli, 42°29'31" N / 43°51'22" E, 1860 m, 23- 25.07.2022, V. Rudoi, P. Ustjuzhanin & R. Yakovlev leg.

Result

No	Taxa	Localities							
		1	2	3	4	5	6	7	8
	Hesperidae								
1	<i>Erynnis tages</i> (Linnaeus, 1758)	-	-	-	+	-	-	-	+
2	<i>Spialia orbi fer</i> (Hübner, 1823)	-	-	-	+	-	-	-	-
3	<i>Pyrgus sidae</i> (Esper, 1784)	+	-	-	-	+	-	-	-
4	<i>Pyrgus</i> (Rambur, 1839)	-	-	+	-	+	-	-	-
5	<i>Carterocephalus palaemon</i> (Pallas, 1771)	-	-	-	-	+	-	-	+
6	<i>Carcharodus alceae</i> (Esper, 1780)	+	-	-	-	-	-	-	-
7	<i>Carcharodus</i> (Zeller, 1847)	-	-	-	+	+	-	-	-
8	<i>Thymelicus lineola</i> (Ochsenhei)	+	-	+	+	+	-	-	+

	mer, 1808)								
9	<i>Thymelicus sylvestris</i> (Poda, 1761)	+	-	-	-	-	-	-	-
10	<i>Ochlodes sylvanus</i> (Esper, 1777)	+	-	+	+	-	+	-	-
	Papilionidae								
11	<i>Papilio machaon</i> Linnaeus, 1758	+	-	-	-	+	-	-	+
12	<i>Iphiclides</i> Linnaeus, 1758	+	-	-	-	-	-	-	-
13	<i>Parnassius apollo</i> Arnold, 1909 (Fig. 2)	-	-	+	+	+	+	-	+
	Pieridae								
14	<i>Leptidea sinapis</i> (Linnaeus, 1758)	+	-	-	-	+	-	-	-
15	<i>Colias hyale</i> (Linnaeus, 1758)	+	-	-	-	-	-	-	-
16	<i>Colias croceus</i> (Geoffroy, 1785)	-	-	+	+	+	-	-	-
17	<i>Colias thiso</i> Ménériés, 1832	-	-	-	-	-	+	-	-
18	<i>Gonepteryx rhamni</i> (Linnaeus, 1758)	-	-	+	-	-	-	-	-
19	<i>Pieris brassicae</i> (Linnaeus, 1758)	+	-	-	-	+	-	-	-

20	<i>Pieris napi</i> (Linnaeus, 1758)	-	-	-	+	-	-	-	+
21	<i>Pieris rapae</i> (Linnaeus, 1758)	+	-	-	-	-	-	-	-
22	<i>Pieris bryoniae casica</i> (Lorković, 1968)	-	-	-	-	-	+	-	-
23	<i>Aporia crataegi</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	+
24	<i>Pontia edusa</i> (Fabricius, 1777)	+	-	-	+	+	-	-	+
	Lycaenidae								
25	<i>Nordmannia ilicis</i> (Esper, 1779)	+	-	-	-	-	-	-	-
26	<i>Callophrys chalybeitincta</i> Sovinsky, 1905	+	-	-	-	-	-	-	-
27	<i>Lycaena phlaeas</i> (Linnaeus, 1761)	-	-	-	-	+	-	-	+
28	<i>Lycaena</i> Linnaeus, 1758	-	-	-	+	+	+	-	+
29	<i>Lycaena alciphron</i> (Rottemburg, 1775)	-	-	-	-	-	+	-	+
30	<i>Celastrina argolus</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	-
31	<i>Everes</i>	-	-	+	-	-	-	-	-

	<i>argiades</i> (Pallas, 1771)								
32	<i>Pseudophylotes vicrama</i> (Moore, 1865)	+	-	-	-	-	-	-	-
33	<i>Cyaniris</i> (Rottemburg, 1775)	+	-	-	+	-	+	-	-
34	<i>Plebejus argus</i> (Linnaeus, 1758)	+	-	-	-	-	-	-	-
35	<i>Plebejus argyrognomon</i> (Bergsträsser, 1779)	-	-	+	-	+	-	-	+
36	<i>Plebejus idas</i> (Linnaeus, 1758)	+	-	-	-	-	+	-	-
37	<i>Aricia agestis</i> ([Denis & Schiffermüller], 1775)	+	-	+	+	-	-	-	-
38	<i>Aricia</i> (Fabricius, 1793)	+	-	-	-	-	-	-	+
39	<i>Glaucopsyche alexis</i> (Poda, 1761)	+	-	-	-	-	-	-	-
40	<i>Phengaris teleius</i> (Bergsträsser, 1779)	+	-	-	-	-	-	-	-
41	<i>Polyommatus amandus</i> (Schneider, 1792)	+	-	+	+	+	+	-	+
42	<i>Polyommatus</i>	+	-	-	-	-	-	-	-

	(Herrich-Schäffer, [1852])								
43	<i>Polyommatus icarus</i> (Rottemburg, 1775)	-	-	-	+	+	+	-	+
	Nymphalidae								
44	<i>Limenitis reducta</i> Staudinger, 1901	+	-	-	-	-	-	-	-
45	<i>Limenitis camilla</i> (Linnaeus, 1764)	-	-	+	-	-	-	-	-
46	<i>Aglaia urticae</i> (Linnaeus, 1758)	-	-	+	+	+	+	-	+
47	<i>Polygonia c-album</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	+
48	<i>Vanessa atalanta</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	-
49	<i>Vanessa cardui</i> (Linnaeus, 1758)	+	-	-	+	-	-	-	+
50	<i>Euphydryas aurinia</i> (Rottemburg, 1775)	-	-	-	+	-	-	-	-
51	<i>Melitaea cinxia</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	+
52	<i>Melitaea phoebe</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	-	-
53	<i>Melitaea interrupta</i> Kolenati, 1846	+	-	+	+	+	+	-	-

54	<i>Melitaea caucasogenita</i> Verity, 1930	-	-	-	+	+	-	-	-
55	<i>Melitaea athalia</i> (Rottemburg, 1775)	-	-	+	-	-	-	-	-
56	<i>Clossiana dia</i> (Linnaeus, 1767)	+	-	-	+	-	-	-	-
57	<i>Clossiana euphrosyne</i> (Linnaeus, 1758)	-	-	+	-	-	-	-	+
58	<i>Brenthis ino</i> (Rottemburg, 1775)	-	-	-	-	+	+	-	+
59	<i>Brenthis daphne</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	-	-
60	<i>Issoria latihonia</i> (Linnaeus, 1758)	+	-	+	-	+	-	-	-
61	<i>Argynnis pandora</i> ([Denis & Schiffermüller], 1775)	+	-	-	-	-	-	+	-
62	<i>Argynnis papiria</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	-
63	<i>Argynnis adippe</i> ([Denis & Schiffermüller], 1775)	+	-	+	+	-	-	-	-
64	<i>Argynnis aglaja</i> (Linnaeus, 1758)	-	-	-	+	+	+	-	+
65	<i>Argynnis niobe taura</i> Röber, 1896	+	-	-	+	-	-	-	-
66	<i>Pararge aegeria</i> (Linnaeus, 1758)	+	-	-	-	-	-	-	-
67	<i>Lasiommata maera</i> (Linnaeus,	-	-	-	-	-	+	-	-

	1758)								
68	<i>Lasiommat a petropolit ana</i> (Fabricius, 1787)	-	-	-	-	+	-	-	-
69	<i>Hipparchia p ell ucida</i> Stauder, 1924	+	-	-	-	-	-	-	-
70	<i>Melanargia galathea</i> (Linnaeus, 1758)	+	-	+	-	-	-	+	-
71	<i>Maniola jurtina</i> (Linnaeus, 1758)	+	-	+	-	-	-	-	-
72	<i>Coenonym pha arc ania</i> (Linnaeus, 1761)	+	-	+	-	-	-	-	-
73	<i>Coenonym pha g lyc erion</i> (Borkhause n, 1788)	-	-	-	-	+	+	-	+
74	<i>Coenonym pha</i> (Linnaeus, 1758)	+	-	+	-	-	-	+	-
75	<i>Erebia medusa</i> ([Denis & S chifferrmüll er], 1775)	-	-	-	+	+	+	-	-
76	<i>Erebia aethiops m elu sina</i> Herrich- Schäffer, 1847 (Fig. 3)	-	-	+	+	+	+	-	+

Table 1.




Figure 1. Map of South Ossetia with collecting localities.



Figure 2. Butterflies of South Ossetia in Nature: **2.** *Parnassius apollo suaneticus* Arnold, 1909, male, Dzaus Distr., Rachinsky Range, near Dодtota, 42°27'25" N / 43°43'18" E, 1750 m (photo by Aleksandr Streltsov)



Figure 3. Butterflies of South Ossetia in Nature: **3.** *Erebia aethiops melusina* Herrich-Schäffer, 1847, male, Dzaus Distr., Tli, 42°29'31" N / 43°51'22" E, 1860 m (photo by Roman Yakovlev).

Conclusions

Thus, on the territory of South Ossetia 77 species of butterflies were reliably indicated. One of them (*Parnassius nordmanni* (Menetries, 1850) is known only from published data. For three species listed in the Red data Book (*P. apollo*, *I. podalirius* and *P. machaon*) we give the new localities. It is necessary to notice that *P. apollo* in the mid-mountain belt of the Rachinsky ridge is the dominant species. The relevance of including species with huge ranges, *Iphiclides podalirius* and *Papilio machaon*, in the Red Book of the region is also not obvious.

References

Bazaev FB, Bestaev AZ, Busarova NV, Butaeva FG, Vaniev AG, Gabaev VN, Dzhioeva TsG, Doroshina GYa, Kabulov AZ, Kabulov ZE, Kabulov ID, Kachmazov DG, Kokoev TI, Komarov YuE, Lavrinenko YuV, Lotiev KYu, Nikolaev IA, Pukhaev RV, Pukhaeva ZA, Timuthin IN, Tuniev BS, Tuniev SB, Khetagurov KhA, Tskhovrebova SS, Tskhovrebova NI, Chibirova AKh, Shikov EV, Yusupov ZM (2017) Red data book of South Ossetia. Poligraphservis & T., Nalchik, 304 pp. [In Russian]

Nedoshivina SV, Ustjuzhanin PY, Kovtunovich VN, Streltsov AN, Yakovlev RV (2023) Lepidoptera of South Ossetia (Northern Transcaucasia). Part III. Tortricidae, Pterophoridae and Alucitidae



(Insecta: Lepidoptera). SHILAP Revista de lepidopterología 51 (203): 437–445.
<https://doi.org/10.57065/shilap.529>

Sinev SYu, Anikin VV, Piskunov VI, Streltsov AN, Ustjuzhanin PY, Yakovlev RV (2023) Lepidoptera of South Ossetia (Northern Transcaucasia). Part IV. Microlepidoptera: Adelidae to Choreutidae. Acta Biologica Sibirica 9: 1061–1072. <https://doi.org/10.5281/zenodo.10213217>

Streltsov AN, Ustjuzhanin PYa, Yakovlev RV (2022a) A new species of the genus *Scoparia* Haworth, 1811 (Lepidoptera: Pyraloidea, Crambidae) from the Transcaucasia. Far Eastern Entomologist 457: 1–6. <https://doi.org/10.25221/fee.457.1>

Streltsov AN, Ustjuzhanin PYa, Yakovlev RV (2022b) Lepidoptera of South Ossetia (Northern Transcaucasia). Part I. Introduction and Superfamily Pyraloidea Latreille, 1809. Acta Biologica Sibirica 8: 281–296. <https://doi.org/10.5281/zenodo.7686863>

Streltsov AN, Ustjuzhanin PYa, Morozov PS, Naydenov AE, Spitsyn VM, Yakovlev RV (2022c) Lepidoptera of South Ossetia (Northern Transcaucasia). Part II. Cossidae, Limacodidae, Erebidae (Lymantriinae, Arctiinae, Syntominae, Notodontinae), Lasiocampidae, Lemoniidae, Saturniidae, Sphingidae, Drepanidae and Cimeliidae. Acta Biologica Sibirica 8: 647–654.

Tshikolovets V, Nekrutenko Y (2012) The Butterflies of Caucasus and Transcaucasia (Armenia, Azerbaijan, Georgia and Russian Federation). Tshikolovets Publications, Kyiv–Pardubice, 423 pp.