

Wasps of the subfamily Bembicinae (Hymenoptera, Crabronidae) in the collection of the Institute of Zoology, Ministry of Science and Education of the Republic of Azerbaijan

Aydin A. Mammadov

Institute of Zoology, Ministry of Science and Education, passage 1128, district 504, Baku, AZ 1004, Azerbaijan; Western Caspian University, 31 Istiglaliyyat st., Baku, AZ1001, Azerbaijan

Mahir M. Maharramov

Nakhchivan State University, Nakhchivan, AZ7012, Azerbaijan

The collection materials of Crabronidae wasps in Institute of Zoology (Baku) presented at the following 16 species from seven Bembicinae genera. Sixteen species are recorded from Azerbaijan, two – from Russia, and six species from Iran. *Gorytes laticinctus* Lepeletier, 1832 is newly recorded from Caucasus and *Gorytes procrustes* Handlirsch, 1888 from Azerbaijan.

Acta Biologica Sibirica 10: 1379–1387 (2024)

doi: 10.5281/zenodo.14258294

Corresponding author: Aydin A. Mammadov (aydin.aydin.2021@mail.ru)

Academic editor: R. Yakovlev | Received 19 September 2024 | Accepted 29 October 2024 | Published 4 December 2024

<http://zoobank.org/6BCBE17C-8D2E-46B0-8DA3-2A512CD3B3B8>

Citation: Mammadov AA, Maharramov MM (2024) Wasps of the subfamily Bembicinae (Hymenoptera, Crabronidae) in the collection of the Institute of Zoology, Ministry of Science and Education of the Republic of Azerbaijan. Acta Biologica Sibirica 10: 1379–1387. <https://doi.org/10.5281/zenodo.14258294>

Keywords

Wasp, Crabronidae, Bembicinae, fauna, Azerbaijan, collection

Introduction

Crabronidae is distributed worldwide, except Antarctic and Arctic zones, mainly in tropical and subtropical regions. Crabronids, including 90 percentages of apoid wasps, is the largest group of insects among all families of wasps (Antropov et al. 2017).

There are 250 genera and 8900 species of crabronids in the world, in Palaeartic – 84 genera and 2000 species, in Azerbaijan 42 genera and 232 species (Antropov et al. 2017, Mokrousov et al.

2019, Mokrousov et al. 2020).

The collection materials of Crabronidae wasps presented at the following 16 species from seven Bembicinae genera. *Gorytes laticinctus* Lepeletier, 1832 are newly recorded from Caucasus and *Gorytes procrustes* Handlirsch, 1888 – from Azerbaijan.

Results

Genus *Hoplisoides* Gribodo, 1884

***H. latifrons* (Spinola, 1808)**

Hoplisoides latifrons: Mokrousov et al., 2019: 12.

Material: Azerbaijan: 1♂, Nakhchivan ASSR [=Nakhchivan MR], 20.05.1933, A. Bogachev.

Remarks: This species was previously recorded from Nakhchivan (Mokrousov et al. 2019).

Distribution: Spain, Portugal, France, Belgium, Switzerland, Italy, Germany, Austria, Czech republic, Croatia, Greece, Slovakia, Hungary, Romania, Ukraine, Russia (Crimea, Dagestan, Volgograd prov., Tatarstan, Orenburg prov.), Algeria, Libya, Azerbaijan, Turkey, Uzbekistan, Tajikistan, Kazakhstan (Mokrousov et al. 2020).

Genus *Argogorytes* Ashmead, 1899

***A. mystaceus* (Linnaeus, 1761)**

Argogorytes mystaceus: Mokrousov et al., 2019: 6.

Material: Azerbaijan: 1♀, Eli-su [=Ilisu, Gakh distr.], 21.07.1934, A. Bogachev.

Remarks: This species was previously recorded from Nakhchivan (Mokrousov et al. 2019).

Distribution: Russia, Europe, Turkey, Kazakhstan, Azerbaijan (Nemkov 2017, Mokrousov et al. 2019).

Genus *Gorytes* Latreille, 1805

***G. quinquecinctus* (Fabricius, 1793)**

Gorytes proximus: Handlirsch, 1893: 281; 1895: 924.

Gorytes quinquecinctus quinquecinctus: Nemkov, 2017: 223.

Gorytes quinquecinctus: Mokrousov et al., 2019: 12.

Material: Azerbaijan: 1♀, Karagashly, Samukh distr., 2.06.1935, on the *Euphorbia* sp. (Euphorbiaceae), M. Vinovsky.

Remarks: This species was previously recorded from Helenendorf [=Goygol] (Handlirsch 1893, 1895) and Nakhchivan (Mokrousov et al. 2019).

Distribution: Russia, Europe, N Africa, Azerbaijan, Turkey, Syria, Iran, Uzbekistan, Kyrgyzstan, Kazakhstan, Azerbaijan (Nemkov 2017, Handlirsch 1893, 1895, Mokrousov et al. 2019).

G. *procrustes* Handlirsch, 1888

Material: Russia: 1♂, Uralsk, 21.07.1927, V. Koloskov; **Azerbaijan:** 1♂, Nakhchivan ASSR, Nakhchivan, 20.05.1933, A. Bogachev; 1♀, Eli-su [=Ilisu, Gakh distr.], 21.07.1934, A. Bogachev; 1♂, Karagashly, Samukh distr., 3.06.1935, on the *Euphorbia* sp. (Euphorbiaceae), M. Vinovsky; 1♀, Eldar, Poily, 21.05.1935, A. Bogachev.

Remarks: newly recorded from Azerbaijan.

Distribution: Russia, Europe, Georgia, Turkey, Kazakhstan (Nemkov 2017).

G. *laticinctus* (Lepeletier de Saint Fargeau, 1832)

Material: Azerbaijan: 1♂, Karagashly, Samukh distr., 3.06.1935, on the *Euphorbia* sp. (Euphorbiaceae), M. Vinovsky.

Remarks: newly recorded from Caucasus and Azerbaijan.

Distribution: Russia, Europe, N Africa, Turkey, Uzbekistan, Kazakhstan (Nemkov 2017).

Genus *Sphecius* Dahlbom, 1843**S. *antennatus* (Klug, 1845)**

Sphecius antennatus: Handlirsch, 1889: 452; Nemkov, 1995: 181; 2017: 224; Mokrousov et al., 2019: 17.

Material: Azerbaijan: 2♀♀, Ganja, 2, 5.07.1933, on the onion's flowers, M. Vinovsky; 2♂♂, Station Ganja railway, 19.07.1933, on the flowers of *Anethum*, M. Vinovsky; 1♀, Ganja, 21.06.1932, in the garden, M. Vinovsky; 2♀♀, Ganja, 11, 29.06.1933, on the onion's flowers, M. Vinovsky; 1♀, Malbinasi, distr. Barda, 7.08.1931, A. Bogachev; 1♂, Yevlakh, 15.06.1940, A. Bogachev; 1♂, Bekhmanly, valley Araz, 18.04.1926, Bocharnikov; 4♂♂, Mil's steppe, 9, 12, 14, 15.06.1934, A. Bogachev; 1♂, Gazakh, 11.07.1941, A. Bogachev; 1♀, Shemakha, 25.06.1937, N. Tertishnikov; 1♂, Kurd-Makhmudly distr., Karyagino (now Fizuli), 14.06.1924, Bocharnikov; 1♂, Mingachaur railway st., 23.07.1933, in the gallery, B. Lototosky. **Iran:** 8♂♂, 1♀, Gaur-arkh, Persia, Ungut-Mugan, 10, 19, 21.06.1927, Bocharnikov.

Remarks: This species was previously recorded from Helenendorf [=Goygol], Yevlax (Handlirsch, 1889), Tash-Bulag near Nukha [=Sheki], Kirovobad [=Ganja], Geokchai, Shamkhor, Kuduly, Julfa, Nakhchivan (Nemkov 1995) and Nakhchivan (Mokrousov et al. 2019).

Distribution: Albania, Cyprus, Greece, Iran, Israel, Italy, Kazakhstan, Mongolia, Montenegro, Russia, Syria, Tajikistan, Turkey, Turkmenistan, Ukraine, Uzbekistan, Azerbaijan (Handlirsch, 1889, Nemkov 1995, Jahantigh et al. 2017, Mokrousov et al. 2019).

Genus *Stizoides* Guérin-Méneville, 1844**S. *tridentatus* (Fabricius, 1775)**

Stizus tridentatus: Handlirsch, 1892: 101.

Stizoides tridentatus: Ohl, 1999: 139; Nemkov, 2012: 10; Mokrousov et al., 2019: 17.

Stizoides tridentatus tridentatus: Nemkov, 2017: 225.

Material: Azerbaijan: 3♀♀, Mil's steppe, 9.06.1934, A. Bogachev; 1♀, Ganja, 29.06.1933, on the onion's flowers, M. Vinovsky.

Remarks: This species was previously recorded from Helenendorf [=Goygol], Murut [=Ucbulaq], Kilasi (Handlirsch 1892), Goradiz, Ganja, Helenendorf [=Goygol], Karadag [=Baku] (Ohl 1999), Altan (Nemkov 2012) and Nakhchivan (Mokrousov et al. 2019).

Distribution: Algeria, Bulgaria, Croatia, Cyprus, Egypt, France, Greece, Hungary, Iran, Israel, Italy, Kazakhstan, Libya, Macedonia, Madagascar, Mongolia, Morocco, Portugal, Romania, Russia, Spain, Switzerland, Tajikistan, Tunisia, Turkey, Turkmenistan, Ukraine, Uzbekistan, Azerbaijan (Handlirsch 1892, Ohl 1999, Nemkov 2012, Jahantigh et al. 2017, Mokrousov et al. 2019).

S. *crassicornis* (Fabricius, 1787)

Stizoides crassicornis: Nemkov, 2017: 225.

Material: Azerbaijan: 2♀♀, Mil's steppe, 11, 14.06.1934, A. Bogachev.

Distribution: Algeria, France, Iran, Israel, Italy, Kazakhstan, Palestine, Russia, Spain, Turkey, Turkmenistan, Ukraine, Azerbaijan (Jahantigh et al. 2017, Nemkov 2017).

Genus *Stizus* Latreille, 1802

S. *annulatus* (Klug, 1845)

Stizus annulatus: Mokrousov et al., 2019: 17.

Material: Azerbaijan: 1♀, Ganja, 5.08.1933, on the *Eryngium* sp. (Apiaceae), M. Vinovsky; 1♀, Ganja, 29.06.1933, on the onion's flowers, M. Vinovsky; 4♀♀, Lerik, Kalakhan, 24.07.1976, H. Aliev; 1♀, Ganja, Kara-yeri, 2.06.1932, on the flowers, M. Vinovsky; 1♀, Lerik, Gosmalyan, 24.07.1976, H. Aliev; 1♂, Baku, Surakhany, 7.07.1936, on the *Eryngium* sp. (Apiaceae), N. Tertysnikov; 1♀, Zakataly, 23.06.1924, on the flying, P. Maximovich; 3♂♂, Baku, Surakhany, 6, 7.07.1936, on the *Eryngium* sp. (Apiaceae), N. Tertysnikov; 1♀, Ganja, 21.07.1932, on the garden, M. Vinovsky; 1♀, Ganja, 28.07.1933, on the *Eryngium* sp. (Apiaceae), M. Vinovsky; 1♀, Ganja, 10.06.1932, on the flying in the garden, M. Vinovsky; 1♀, Ganja, 22.07.1932, on the flying in the garden, M. Vinovsky.

Remarks: This species was previously recorded from Nakhchivan (Mokrousov et al. 2019).

Distribution: SE Europe, N Africa, Armenia, Turkey, Syria, Israel, Iran, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, Azerbaijan (Roche 2007, Nemkov 2017, Mokrousov et al. 2019).

S. *rufiventris* Radoszkowski, 1877

Stizus rufiventris: Mokrousov et al., 2019: 17.

Material: Azerbaijan: 5♂♂, 3♀♀, Baku, Botanical garden, 10.07.1937, M. Vinovsky.

Remarks: This species was previously recorded from Nakhchivan (Mokrousov et al. 2019).

Distribution: Turkey, Turkmenistan, Uzbekistan, Tajikistan, Kazakhstan, Mongolia, Iran, Azerbaijan. (Yildirim et al. 2016, Mokrousov et al. 2019).

Genus *Bembix* Fabricius, 1775

B. *eburnea* Radoszkowski, 1877

Bembix eburnea: Mokrousov et al., 2019: 8.

Material: Azerbaijan: 2♀♀, Ganja, 11.07.1933, on the flying, M. Vinovsky; Iran: 1♂, Mugan Persia, Gaur-Ark, 12.06.1927, Bocharnikov; 1♂, Persia, Mugan, Kuruchay, 21.06.1927, Bocharnikov.

Remarks: This species was previously recorded from Nakhchivan (Mokrousov et al. 2019).

Distribution: Turkey, Russia (Crimea, Volgogradskaya Oblast), Kazakhstan, Uzbekistan, Turkmenistan, Tajikistan, Iran, Afghanistan, Mongolia, China (Xingiang, Inner Mongolia, Hebei, Beijing, Tianjin), Azerbaijan (Nemkov 2016, Mokrousov et al. 2019).

***B. rostrata* (Linnaeus, 1758)**

Bembix rostrata: Nemkov, 2016: 22; Mokrousov et al., 2019: 8.

Bembix rostrata rostrata: Nemkov, 2017: 222.

Material: Azerbaijan: 1♂, 1♀, Zakataly, Katekhy, 22.07.1933, A. Bogachev; 2♀♀, Zakataly, 29.06.1938, A. Bogachev; 1♀, Lerik, Gosmalyan, 24.07.1976, H. Aliev; 1♂, Ganja, 26.07.1932, in the garden, M. Vinovsky; **Iran:** 1♀, Mugan, Persia, Gaur-Ark, 12.06.1927, Bocharnikov.

Remarks: This species was previously recorded from Belokany (Nemkov, 2016) and Nakhchivan (Mokrousov et al. 2019).

Distribution: Russia, Europe, Azerbaijan, Turkey, Iran, Afghanistan, Turkmenistan, Tajikistan, Uzbekistan, Kazakhstan, Mongolia, China, Azerbaijan (Nemkov 2016, 2017, Mokrousov et al. 2019).

***B. oculata* Panzer, 1801**

Bembix oculata: Nemkov, 2016: 16; Mokrousov et al., 2019: 8.

Bembix oculata oculata: Nemkov, 2017: 222.

Material: Azerbaijan: 3♀♀, Kalinovka Vilyajchay river, 9, 15.07.1936, M. Vinovsky; 1♀, Baku, 07, Akhundova; 1♂, 1♀, Port-Ilyich, distr. Lenkoran, 3.07.1936, M. Vinovsky; 1♀, Talysh, Alekseyevka, 7.08.1936, A. Bogachev; 2♂♂, Ganja, 31.07.1932, in the garden, M. Vinovsky; 2♂♂, 1♀, Ganja, 5, 13.08.1932, in the steppe, M. Vinovsky; 1♂, 1♀, Ganja, 5, 17.08.1933, on the *Eryngium* sp. (Apiaceae), M. Vinovsky; 1♀, Ganja, 5.07.1932, in the garden, M. Vinovsky; 1♀, Prishib. distr., Lenkoran, 08.1923, Bocharnikov; 1♀, Krym, Dorankoy, 9.07.1926, V. N. Kuznetsov; **Iran:** 1♀, Mugan Pers., Gaur-arkh, 12.06.1927, Bocharnikov; 1♂, Mugan Pers., Tazakend, 17.08.1927, Bocharnikov; 1♂, Mugan Pers., Tazakend, 17.08.1927, Bocharnikov.

Remarks: This species was previously recorded from Balaken, Gumbashi (Nemkov 2016) and Nakhchivan (Mokrousov et al. 2019).

Distribution: Russia, Europe, N Africa, Georgia, Azerbaijan, Syria, Jordan, Lebanon, Israel, Saudi Arabia, United Arab Emirates, Oman, Iran, Afghanistan, Pakistan, Kazakhstan, Uzbekistan, Turkmenistan, Tajikistan, Kazakhstan, Mongolia, China, Azerbaijan (Nemkov 2016, 2017, Mokrousov et al. 2019).

***B. olivacea* Fabricius, 1787**

Bembix olivacea olivacea: Nemkov, 2017: 222.

Material: Azerbaijan: 11♀♀, ost. fl. Samur caucas. or., 23.08.1940, E. Hauser; 1♀, Nakhchivan, 25.07.1928; 1♀, Sabirabad, 10.08.1939, A. Bogachev; **Iran:** 1♀, Mugan Pers., Gaur-arkh, 30.05.1927, Bocharnikov; 1♀, Mugan Pers., Tazakend, 17.08.1927, Bocharnikov.

Distribution: Algeria, Azerbaijan, Bulgaria, Canary Islands, Chad, Cyprus, Egypt, Ethiopia, France, Greece, Hungary, Iran, Italy, Israel, Kazakhstan, Libya, Madagascar, Mauritania, Morocco, Portugal, Romania, Russia, Saudi Arabia, Serbia, Slovakia, Somalia, Spain, Sudan, Syria, Tunisia, Turkey, Turkmenistan, Ukraine, Western Sahara, Yemen (Jahantigh et al. 2017).

B. *bidentata* Vander Linden, 1829

Bembex bidentata: Handlirsch, 1893: 775.

Bembex bidentata: Nemkov, 2016: 9; 2017: 221; Mokrousov et al., 2019: 7.

Material: Azerbaijan: 6♂♂, 1♀, Kirovabad (now Ganja), Bolchaly, 17, 20.07.1933, M. Vinovsky; 1♂, Samukh, Karagashly, 31.05.1935, M. Vinovsky; 1♀, Kirovabad (now Ganja), Karayery, 2.06.1932, M. Vinovsky; 6♀♀, ost. flav. Samur caucas. or., 23, 26.08.1940, E. Hauser; 1♂, 1♀, Samukh, Karagashly, 31.05.1935, A. Bogachev; 2♀♀, Zakataly, Ali-Bayramly, 2, 26.06.1938, A. Bogachev; 2♂♂, Mugan bor. Djafarkhan, 26.06.1925, Alexandrov; 5♀♀, Lerik, Kosmalyan, 24.07.1976, H. Aliev; 1♀, Kalinovka, Vilyaj-chay river, 15.07.1936, M. Vinovsky; 1♀, Ordubad, 2.06.1939, E. Hauser; 1♂, Lerik, Kosmalyan, 24.07.1976, H. Aliev; 5♀♀, Lerik, Galabyn, 21, 22, 23, 27.07.1976, H. Aliev; 2♂♂, Mingachevir, 6.07.1946, on the Salsoletum, A. Bogachev; 1♂, steppe, on the road to Bolchaly village, 17.07.1933, on the fly, Asia, Kirovabad (now Ganja), M. Vinovsky; 1♂, Samukh, Karagashly, 31.05.1935, M. Vinovsky; 1♀, Zakataly, Ali-Bayramly, 2.07.1938, A. Bogachev; 1♀, station Ganja railway, 19.07.1933, on the Anethum's flowers, M. Vinovsky; 2♀♀, Lerik, Gilidara, 17, 27.07.1976, H. Aliev; **Iran:** 1♂, Mugan Pers., Kuruchay, 2.06.1927, Bocharnikov;

Remarks: This species was previously recorded from Murut [=Ucbulaq] (Handlirsch 1893), Agdere; Kudula (Nemkov 2016) and Nakhchivan (Mokrousov et al. 2019).

Distribution: South Europe, Turkey, Kazakhstan, Southern Siberia, Iran, Azerbaijan (Handlirsch 1893, Yildirim et al. 2016, Nemkov 2016, Mokrousov et al. 2019).

B. *bicolor* Radoszkowski, 1877

Bembex bicolor: Mokrousov et al., 2019: 7.

Material: Azerbaijan: 1♂, Ganja, on the fly, 11.07.1933, M. Vinovsky.

Remarks: This species was previously recorded from Nakhchivan (Mokrousov et al. 2019).

Distribution: Afghanistan, Bulgaria, China, Cyprus, Greece, Iran, Iraq, Italy, Israel, Kazakhstan, Kyrgyzstan, Mongolia, Russia (European), Tajikistan, Turkey, Turkmenistan, Uzbekistan, Azerbaijan (Mokrousov et al. 2019, Gadallah, 2020).

References

Antropov AV, Astafurova YV, Belokobylskij SA, Byvaltsev AM, Danilov YuN, Dubovikoff DA, Fadeev KI, Fateryga AV, Kurzenko NV, Lelej AS, Levchenko TV, Loktionov VM, Mokrousov MV, Nemkov PG, Proshchalykin MYu, Rosa P, Sidorov DA, Sundukov YuN, Yusupov ZM, Zaytseva LA (2017) Annotated catalogue of the Hymenoptera of Russia. Volume I. Symphyta and Apocrita: Aculeata. Proceedings of the Zoological Institute RAS. Supplement 6, 475 pp.

Gadallah NS (2020) Biodiversity of the aculeate wasps (Hymenoptera: Aculeata) of the Arabian Peninsula: Apoidea (Spheciformes), Crabronidae. In: Gadallah NS, Brothers DJ, Williams KA (Eds) Biodiversity of the aculeate wasps (Hymenoptera: Aculeata) of the Arabian Peninsula. Zootaxa 4754: 20–73.

Handlirsch A (1889) Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. IV. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I, 98: 440–517.

Handlirsch A (1892) Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. VI. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I, 101: 25–205.

Handlirsch A (1893) Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. VII. (Schluss). Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I, 102: 657–942.

Handlirsch A (1895) Nachträge und Schlusswort zur Monographie der mit *Nysson* und *Bembex* verwandten Grabwespen. Sitzungsberichte der Kaiserlichen Akademie der Wissenschaften. Mathematisch-Naturwissenschaftliche Classe. Abtheilung I, 104: 801–1079.

Jahantigh F, Rakhshari E, Mokhtari A, Ramroodi S (2017) Catalogue of Ampulicidae, Crabronidae and Sphecidae of Iran (Hymenoptera, Apoidea). Zootaxa 4307 (1): 1–96.
<https://doi.org/10.11646/zootaxa.4307.1.1>

Mokrousov MV, Proshchalykin MYu, Aliyev KhA, Maharramov MM (2019) To the knowledge of digger wasps (Hymenoptera: Crabronidae) of Nakhchivan Autonomous Republic of Azerbaijan. Far Eastern Entomologist 394: 1–24.

Mokrousov MV, Proshchalykin MY, Maharramov MM, (2020) Digger wasps of the genus *Hoplisoides* Gribodo (Hymenoptera, Crabronidae, Bembicinae) from the Palaearctic region, with description of two new species. Journal of Hymenoptera Research 79: 213–233.
<https://doi.org/10.3897/jhr.79.56839>

Nemkov PG (1995) Sphecoid wasps of the tribe Gorytini (Hymenoptera, Sphecidae) of the fauna of the CIS. The genera *Argogorytes* Ashmead, *Hoplisoides* Gribodo, *Psammaecius* Lepeletier. Trudy Zoologicheskogo Instituta Rossiyskoy Akademii Nauk 258: 128–137. [In Russian]

Nemkov PG (2012) Digger wasps of the genus *Stizus* Latreille, 1802 (Hymenoptera, Crabronidae, Bembicinae) of the fauna of Russia and neighbouring countries. Euroasian Entomological Journal 11(1): 55–62.

Nemkov PG (2016) Digger wasps of the genus *Bembix* Fabricius, 1775 (Hymenoptera: Crabronidae, Bembicinae) of Russia and adjacent territories. Far Eastern Entomologist 313: 1–34.

Nemkov PG (2017) Subfamily Bembicinae. In: Belokobylskij SA, Lelej AS (Eds) Annotated Catalogue of the Hymenoptera of Russia. Volume I. Symphyta and Apocrita: Aculeata. Proceedings of the Zoological Institute of the Russian Academy of Sciences 6: 219–225.

Ohl M (1999) A revision of *Stizoides* Guérin-Méneville, 1844: taxonomy, phylogenetic relationships, biogeography, and evolution (Hymenoptera: Apoidea: “Sphecidae”). Mitteilungen aus dem Museum für Naturkunde in Berlin, Zoologische Reihe 75: 63–169.

Roche CG (2007) Conspectus of the Sphecoid wasps of Egypt (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae). Egyptian Journal of Natural History 4: 12–149.



Yildirim E, Ljubomirov T, Özbek H, Yüksel M (2016) New data on Spheciformes fauna (Hymenoptera: Ampulicidae, Sphecidae, Crabronidae) of Turkey. *Journal of Insect Biodiversity* 4: 1–51.